

ISBN 89-950054-9-1 93520
89-950054-7-5 93520

Proceedings

CONTRIBUTED PAPERS – Vol. II



The 8th World Conference on Animal Production
June 28 – July 4, 1998
Seoul National University
Seoul, Korea

Organizing Committee of the 8th WCAP

(1) **Conference Presidents** : Hyun U. Kim and Kil S. Chung

(2) **Chairman** : Jong K. Ha

(3) **Secretary General** : Hong L. Choi **Secretary** : Mr. J. H. Kim

(4) **Organizing Subcommittees**

Scientific Program

Chairman : I. K. Paik

Vice-Chairman : C. W. Kang

Finance

Chairman : C. W. Kim

Vice-Chairman : D. J. Yoo

Publication

Chairman : J. K. Jung

Vice-Chairman : H. W. Lee

Home Stay Program

Chairman : Y. J. Choi

Vice-Chairman : T. S. Koh

Special & Social Program

Chairman : N. H. Lee

Vice-Chairman : Y. S. Son

Program Coordinating

Chairman : H. T. Lee

Vice-Chairman : C. K. Han

Tour & Transportation

Chairman : B. D. Cho

Vice-Chairman : H. Y. Park

Animal Life Science Exhibition

Chairman : K. M. Chung

Vice-Chairman : J. T. Yoon, H. H. Im, Y. S. Shin, Y. C. Kim, K. B. Lim

Publication Subcommittee

Chairman : J. K. Jung

Vice-Chairman : H. W. Lee

Members : S. H. Choi, J. B. Kim, K. S. Sohn, D. I. Jin, K. J. Jeon, S. C. Lee, J. D. Kim,
B. K. Lee, Y. H. Choi

Editorial Remark : All the contributed papers were photocopied from original manuscripts submitted to Publication Subcommittee without editing except those exceeding 2 pages. Some of authors did not observe Guidelines for Abstract Preparation, in which exact format and style of abstract were described. Thus, style and format of abstracts in this volume are not consistent.

Proceedings

CONTRIBUTED PAPERS

Vol. II

The 8th World Conference on Animal Production

June 28 - July 4, 1998

Seoul National University

Seoul, Korea

	Holstein) and Holstein beef cattle	744
PP1-13	V. Skrivanova, M. Skrivan, M. Marounek, P. Klein, J. Lastovkova and M. Koblirle (Czech Republic) Attempt to increase content of polyunsaturated fatty acids in meat of calves	746
PP1-14	D. H. Kim, B. Y. Park, Y. M. Yoo, J. M. Lee, I. B. Chung, Y. K. Kim, I. S. Kim and M. H. Lee (Korea) Characteristics of Technical Meat Quality Parameters between the PSE and normal pork classified by CIE L* Threshold	748
PP1-15	J. S. Ham, S. G. Jeong, Y. M. In, H. J. Park and Y. K. Kim (Korea) Making and Utilization of low cholesterol frozen egg yolk	750
PP1-16	B. Y. Park, Y. M. Yoo, J. M. Lee, D. H. Kim, K. S. Park and Y. K. Kim (Korea) Studies on the FFA, VBN changes and panal traits of chilled Hanwoo beef for storage period	752
PP1-17	J. M. Lee, Y. M. Yoo, B. Y. Park, J. H. Kim and Y. K. Kim. (Korea) Effect of market weight and sex on the carcass characteristic of commercial pig in Korea ...	754
PP1-18	B. Y. Park, Y. M. Yoo, J. M. Lee, D. H. Kim, Y. K. Kim and G. B. Park (Korea) Study on the incidence of Dark-cutting in Korean native cattle (Hanwoo)	756
PP1-19	J. H. Kim, B. Y. Park, Y. M. Yoo, J. M. Lee and Y. K. Kim (Korea) Effects of dietary oils on fatty acids composition of white and dark meat from spent hens	758
PP1-20	D. H. Kim, I. B. Chung, Y. K. Kim, J. W. Lee, I. S. Kim and M. H. Lee (Korea) Relationships between muscle conductance and pork quality factors	760
PP1-21	N. H. Lee, Y. B. Kim, C. S. Yoon and K. S. Seong (Korea) Study on the shelf-life on Vaccum Packaged Chilled Beef Produced in USA	762
PP1-22	I. J. Yoo, K. H. Jeon, W. M. Park, K. S. Kim and S. D. Lim (Korea) Study on suppression of bone crumble in retorted <i>Sangye-tang</i>	764
PP1-23	S. C. Huang and Y. C. Wu (Taiwan) Effect of calcium chloride on tenderization of spent layers meat	766
PP1-24	P. Y. Yueh, H. C. Ding and H. P. Su (Taiwan) Effect of different processing conditions on the stability of immunoglobulin Y in chicken eggs	768
PP1-25	Y. I. Choi, H. G. Cho and C. L. Lee (Korea) Incidence of PSE by several antemortem factors	770
PP1-26	Y. M. Yoo, B. Y. Park, J. M. Lee, J. H. Kim, Y. K. Kim and H. K. Park (Korea) Effect of backfat thickness on the characteristics of carcass components and beef quality of Hanwoo bulls and steers	772
PP1-27	C. N. Meghen, C. S. Scott, D. G. Bradley, D. E. MacHugh, R. T. Loftus and E. P. Cunningham (Ireland) DNA based traceability for the meat industry	774
PP1-28	C. S. Lee, S. K. Park, O. H. Kim, B. C. Sang and H. R. Oh (Korea) Meat identification of Korean native cattle(Hanwoo) and European cattles(Non-Hanwoo) by DNA polymorphic analysis	776

Section 10 : Milk Processing (PP2)

PP2-1	Agung Purnomoadi, F. Nakai, K. Higuchi, K. Ueda, O. Enishi, M. Kurihara, F. Terada and A. Abe (Japan) The use of near infrared spectra of urine for monitoring milk urea nitrogen in dairy cattle	780
PP2-2	J. K. Cho and Y. H. Yoon (Korea) Antimutagenic activity of <i>Lactobacillus</i> spp. from Kefir and Yoghurt and non-starter strains	782

The use of near infrared spectra of urine for monitoring milk urea nitrogen in dairy cattle

Agung Purnomoadi, F. Nakai*, K. Higuchi, K. Ueda,
O. Enishi, M. Kurihara, F. Terada and A. Abe

National Institute of Animal Industry, Tsukuba, Ibaraki 305-0901, Japan
**Tokushima pref. Livestock Exp. Sta., Tokushima 771-1310, Japan*

Introduction

Milk should be an ideal medium for evaluating the health and efficiency of production of dairy cattle, because the concentration of urea in milk can be used as an indicator of nutritional status of dairy cows due to the relationship between the balance and energy to protein intake (Pehrson, 1996). However, the previous study on measurement of milk urea nitrogen (MUN) by near infrared (NIR) spectroscopy of milk was failed because of the interference by dominant fat components and the fact that MUN is a trace component (Purnomoadi *et al.*, 1998). An ideal medium for assessing the nutritional status may be provided by urine. It is known that there is a significant relationship between milk urea and urinary nitrogen (Gonda and Lindberg, 1994), hence it may be worth trying to use urine for detecting nutritional status of dairy cows.

This study was done to investigate the most available wavelength of urine spectra for monitoring nutritional status in animals as shown by MUN. This study is a part of the work for developing a continuous system measurement for a live animal system using NIR spectra.

Materials and Methods

Four lactating Holstein cows were housed in a controlled climatic room over four weeks and were exposed to a temperature of 18°C for the first two weeks, and then to 28°C for the next two weeks. This elevation was performed so as to obtain the significant trend of MUN because if temperature is raised to over comfort zone, the cow will be energetically underfeeding due to the decreasing dry matter intake. Feed was given four times a day at 08:00, 10:00, 16:30 and 19:00 to meet the requirements of maintenance and milk production (MAFF, 1994). Milking was done twice a day in the morning and evening at 09:00 and 18:00, respectively. Milk yield was recorded from week-2 to week-4. The concentration of MUN was determined by Urease-Indophenol method (Wako pure chemical, Ltd., Japan). Urine samples were collected daily after morning milking. Samples were filtered to separate the solid contaminants.

NIR spectra of the urine samples were measured with a Pacific Scientific (Neotec) model 6500 instrument (Perstorp Analytical, MD) using a transmittance cell sample (thickness : 1 mm). Spectra were read over a range of wavelengths between 1100 and 2500 nm at 2 nm intervals. The spectra were then calculated on second derivative of $\log 1/T$, where T is transmittance. The spectra from daily urine samples (n=76) were averaged and analyzed to find the wavelength that had the strongest relationship with MUN. Analysis was done using linear regression from the 700 points of wavelength recorded.

Results and Discussion

The average of daily milk production, MUN and spectra of urine at 1550 nm during elevation of temperature from 18°C to 28°C are shown in figure 1. Milk production started to decrease at day-1 after elevating the room temperature. The decrease continued, stopped around day-6 and then steadied. The phenomenon of MUN level was contradictory to that of milk production. Though milk production steadied at day-6, MUN kept decreasing to the original level (before the temperature was elevated). If

MUN expresses the balance of energy and protein, it can be assumed that the recovery process to balance the protein to energy ratio still continues until day-12.




Figure 1. The trend of daily morning yield, MUN (○) and the NIR spectra of urine (●) at 1550 nm

Detection of daily increase in MUN using NIR spectra of urine using the wavelength of urine spectra at 1550 nm (U-1550) was highly correlated ($r = 0.92$). Both trends of MUN and U-1550 are also shown in figure 1. The trend of U-1550 is very similar to the trend of MUN level and seems to be more sensitive rather than that of MUN. The MUN level elevation from day-2 to day-6 seems to be one-day later compared to the U-1550. This is because the milk collected in the morning is the accumulation of milk secreted in the udder after evening milking until the next morning milking. The wavelength at 1550 nm in urine is considered to be the absorbance of N-H bond component (Murray and Williams, 1990).

This result not only confirms a significant relationship between milk urea and urinary nitrogen, but also strongly shows that the spectra of urine can be used to monitor the variation of MUN in order to manage the energy-protein balance status in dairy cattle.

References

- Gonda HL and Lindberg LE. 1994. Evaluation of Dietary Nitrogen Utilization in Dairy Cows Based on Urea Concentrations in Blood, Urine and Milk, and on Urinary Concentration of Purine Derivatives. *Acta Agric. Scand.* 44: 236-245.
- MAFF. 1994. Japanese Feeding Standard for Dairy Cattle. Central Assoc. Livestock Ind., Tokyo. 115.
- Murray I and Williams PC. 1990. Chemical Principles of Near -Infrared Technology, in: *Near-Infrared Technology in the Agricultural and Food Industries*, Ed. Williams PC and Norris KH. American Assoc. Cereal Chemist, Inc. St.Paul, Minnesota, USA. 17-34.
- Pehrson B. 1996. Milk Analysis as an Indicator of The Nutritional and Disease Status of Dairy Cows, in: *Recent Advances in Animal Nutrition*, Ed. Garnsworthy PC, Wiseman J and Haresign W. Nottingham Univ. Press. 113-133.
- Purnomoadi A, Batajoo KK, Ueda K, Nishida T, Kurihara M, Terada F and Abe A. 1998. The use of near infrared spectroscopy for predicting milk nitrogenous components. *Proc. The 94th Symposium of Jap. Soc. Zootech. Sci. Kobe Univ.* 67.