

DAFTAR PUSTAKA

1. Yoshinaga I.C, Sedey AL, Coulter DK, Mehl AL. Language of early and later identified children with hearing loss. *Pediatrics* 1998;102:1161-71.
2. American Academy of Pediatrics, Task force on newborn and infant screening, newborn and infant hearing loss: detection and intervention. *Pediatrics* 1999;103: 527-30.
3. Joint Committee on Infant Hearing, year 2000 position statement: principles and guidelines for early hearing detection and intervention. *Pediatrics* 2000;106:798-817.
4. Runjan L, Amir I, Suwento R, Mangunatmadja I. Skrining gangguan pendengaran pada neonatus risiko tinggi. *Sari Pediatri* 2005;6(4):149-54.
5. Thompson DC, McPhilips H, Davis RL, Lieu TA, Homer CJ, Helfand M. Universal newborn hearing screening. *JAMA* 2001;286:2000-10.
6. Suwento R. Gangguan pendengaran pada bayi dan anak. Dalam: Seminar Sehari Penatalaksanaan Gangguan Pendengaran Dan Ketulian. 17 Februari 2007; di Semarang; FK Undip; 2007. h. 1-12.
7. World Health Organization, situation review and update on deafness, hearing loss and intervention programmes: proposed plans of action for prevention and alleviation of hearing impairment in countries of the south-east asia region. New Delhi; WHO 2007.
8. Fortnum HM, Summerfield AQ, Marshall DH, Davis AC, Bamford JM. Prevalence of permanent childhood hearing impairment in the United Kingdom and implications for universal neonatal hearing screening: questionnaire based ascertainment study. *Br Med J* 2001;323:1-6.
9. Jardim IS, Matas CG, Carvallo RMM. Newborn hearing screening with transient evoked otoacoustic emissions and automatic auditory brainstem response. *Einstein* 2008;6:253-61.
10. Sukadi A. Hiperbilirubinemia. Dalam: Kosim MS, Yunanto A, Dewi R, Sarosa GI, Usman A, editor. *Buku Ajar Neonatologi*. Edisi pertama. Jakarta; Badan penerbit IDAI; 2008. h.147-69.

11. American Academy of Pediatrics, Provincial Committee for Quality Improvement. Practice parameter: management of hyperbilirubinemia in the healthy term newborn. *Pediatrics* 1994; 94: 558-65.
12. Rapin I. Hearing impairment. In: Swaiman K.F, Ashwal S, editors. *Pediatric Neurology Principles and Practice*. 3rd ed. New York. Mosby Inc; 1999. p.77-95
13. Cunningham M, Cox E O. Hearing assessment in infants and children: recommendations beyond neonatal screening. *Pediatrics* 2003; 111:436-40.
14. Mason JA, Herrmann KR. Universal infant hearing screening by automated auditory brainstem response measurement. *Pediatrics* 1998;101:221- 8.
15. Sharma P, Chhangani N.P, Meena K.R, Jora R, Sharma N, Gupta B.D. Brainstem evoked response audiometry (BAER) in neonates with hyperbilirubinemia. *Indian J Pediatr* 2006;73(5):413-6.
16. Amin SB, Ahlfors C, Orlando MS, Dalzell LE, Merle KS, Guillet R. Bilirubin and serial auditory brainstem response in premature infants. *Indian J Pediatr* 2006;73 (5):413-6.
17. Ahlfors CE, Parker AE. Unbound bilirubin concentration is associated with abnormal automated auditory brainstem response for jaundiced newborns. *Pediatrics* 2008; 121: 976-8.
18. MedlinePlus medical encyclopedia: Hearing loss. Diakses 28 Agustus 2008. Diambil dari URL: <http://www.nlm.nih.gov/medlineplus/ency/article/003044.html>
19. Schrijver I. Hereditary non-syndromic sensorineural hearing loss. *J Mol Diagn* 2004;6(4):275-84.
20. Suwento R. Deteksi dini gangguan pendengaran pada anak. Dalam: *Simposium Sehari Anakku Tidak Bisa Mendengar*. 18 Februari 2006; di Semarang. Semarang; FK Undip;2006. h.15- 27.
21. Deka RC, Sarin D. Congenital TORCH infection and hearing loss. In: Deka D, editor. *Congenital Intrauterine TORCH Infections*. 1st ed. New Delhi, Jaypee brothers; 2004. p.112-20.

22. Rivera LB, Boppana SB, Fowler KB, Britt WJ, Stagno S, Pass RF. Predictors of hearing loss in children with symptomatic congenital cytomegalovirus infection. *Pediatrics* 2002;110:762-7.
23. Richardson MP, Reid A, Tarlow MJ, Rudd PT. Hearing loss during bacterial meningitis. *Arch Dis Child* 1997;76:134-8.
24. Smith RJH, Bale JF, White KR. Sensorineural hearing loss in children. *Lancet* 2005;365:879-90.
25. Hain CT. Ototoxic medications. Di akses 28 Agustus 2008. Diambil dari URL: <http://www.tchain.com/otoneurology/disorders/bilat/ototoxins.html>
26. Volpe JJ. Viral, protozoan, and related intracranial infection. In: Volpe JJ , editor. *Neurology of the Newborn*. 4th ed. Philadelphia:W.B.Saunders Company;2001. p.717-73.
27. Maria A, Deorari A. Outcome of neonates born with vertically transmitted TORCH infections. In: Deka D, editor. *Congenital Intrauterine TORCH Infections*. 1st ed. New Delhi, Jaypee brothers; 2004. p.82-98.
28. Meyer C, Witte J, Hildmann A, Hennecke KH, Schunck KU, Maul K et al. Neonatal screening for hearing disorders in infants at risk: Incidence, risk factors, and follow up. *Pediatrics* 1999;104:900-4.
29. Chu K, Elimian A, Barbera J, Ogburn P, Spitzer A, Quirk JG. Antecedents of newborn hearing loss. *Obstet Gynecol* 2003; 101: 584-8.
30. Zamani A, Daneshjou K, Ameni A, Takand J. Estimating the incidence of neonatal hearing loss in high risk neonates. *Acta Medica Iranica* 2004; 42(3): 176- 80.
31. Beata ACS, Ioana ANCA. Etiopathogeny of congenital and early-onset hearing loss; detection and early intervention methods in infants and children. *Medica* 2008; 3: 45-51.
32. Widiastuti MI. BERA. Semarang; Laboratorium Ilmu Penyakit Saraf FK UNDIP RS Dokter Kariadi; 1992.
33. Abiratno SF. Auditory brainstem response (ABR) prinsip dasar, tehnik pemeriksaan dan penggunaannya dalam klinik. Dalam: Abiratno SF, editor. *Workshop Peranan OAE dan ABR Dalam Klinik*. 29 Maret 2003; di Surabaya; FK Unair; 2003.

34. Porter ML, Dennis BL. Hyperbilirubinemia in the term newborn. *AAFP* 2002; 65: 599-606.
35. Halamek LP., Stevenson DK. Neonatal jaundice and liver disease. In: Fanaroff AA, Martin RJ, Editors. *Neonatal-Perinatal Medicine; Diseases of the Fetus and Infant*, 6th ed. New York Mosby-Year Book Inc;1997.p.1345-62.
36. Dennery PA, Seidman DS, Stevenson DK. Neonatal hyperbilirubinemia. *New Eng J Med* 2001; 344(8):581-90.
37. Gomella TL, Cunningham MD, Eyal FG, Zenk KE. Hiperbilirubinemia. In: Gomella TL, editor. *Neonatology; Management Procedures, On-call Problems, Diseases and Drugs*. New York. Lange Medical Book/McGraw-Hill Co; 2004. p. 381-95.
38. Aminullah A. Ikterus dan hiperbilirubinemia pada neonatus. Dalam: Markum AH, Ismael S, Editor. *Buku Ajar Ilmu Kesehatan Anak*. Jakarta : FKUI; 1999. h. 313-7.
39. Soetomenggolo TS, Iman S. Kelainan toksik dan nutrisi. Dalam : Soetomenggolo TS, Ismael S, Editor. *Buku Ajar Neurologi Anak*, Cetakan ke 2. Jakarta: BP IDAI; 2000. h.541-3.
40. Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics* 2004;114:297-316.
41. Gregg RB, Wiorek LS, Arvedson JC. Pediatric audiology: a review. *Pediatr Rev* 2004; 25:224-34.
42. Shapiro SM. Bilirubin toxicity in the developing nervous system. *Ped Neurol* 2003; 29(5):410-21.
43. Carter K. Kernicterus in full-term infants-United states,1994-1998. *JAMA* 2001; 286: 299-300.
44. Shapiro SM, Nakamura H. Bilirubin and the auditory system. *J of Perinatol* 2001;21:S52-S55.
45. Volpe JJ. Bilirubin and brain injury. In: Volpe JJ,editor. *Neurology of The Newborn*. 4th ed. Philadelphia: W.B. Saunders Company; 2001. p. 521-46.

46. Alatas H, Karyomanggolo WT, Musa DA, Boediarso A, Oesman IN. Desain penelitian. Dalam: Sastroasmoro S, Ismael S, Editor. Dasar-dasar Metodologi Penelitian Klinis. Edisi 2. Jakarta: Sagung Seto; 2002. h. 79-96.
47. Taralan T, Soetomenggolo TS, Passat J, Agusman IS. Studi kohort. Dalam: Sastroasmoro S, Ismael S, Editor. Dasar- Dasar Metodologi Penelitian Klinis. Edisi 2. Jakarta: Sagung Seto; 2002. h. 97-109.
48. Maisels MJ. Neonatal hyperbilirubinemia. In: Klaus MH, Fanaroff AA, editors. Care of The High Risk Neonate, 5th ed. Philadelphia. W.B. Saunders Company; 2001. p.324-62.
49. Tioseco JA, Aly H, Milner J, Patel K, El-Mohandes AAE. Does gender affect neonatal hyperbilirubinemia in infants? *Pediatr Crit Care Med* 2005; 6: 171-4.
50. Boo NY, Oakes M, Lye MS, Said H. Risk factors associated with hearing loss in term neonates with hyperbilirubinaemia. *J Trop Pediatr* 1994;40:194-7.
51. Moore DR. Auditory development and the role of experience. *BMB* 2002;63:171-81.
52. Madden C, Rutter M, Hilbert L, Greinwald JH, Choo DI. Clinical and audiological features in auditory neuropathy, *Arch Otolaryngol Head Neck Surg* 2002;128:1026-30.
53. Fallon JB, Irvine DRF, Shepherd RK. Cochlear implants and brain plasticity. *Hear res* 2008;238:110-7.
54. Holster IR, Hoeve LJ, Wieringa MH, Willis-Lorrier RMS, Gier HHW. Evaluation of hearing loss after failed neonatal hearing screening. *J Pediatr* 2009;155:646-50.
55. Oysu C, Ulubil A, Aslan I, Baserer N. Incidence of cochlear involvement in hyperbilirubinemic deafness. *Ann Otol Rhinol Laryngol* 2002;111:1021- 4.
56. Boo NY, Rohani AJ, Asma A. detection of sensorineural hearing loss using automated auditory brainstem-evoked response and transient-evoked otoacoustic emission in term neonates with severe hyperbilirubinemia. *Singapore Med J* 2008; 49:209- 14.

57. Wennberg RP, Ahlfors CE, Bhutani VK, Johnson LH, Shapiro SM. Toward understanding kernicterus: a challenge to improve the management of jaundiced newborns. *Pediatrics* 2006;117;474-85.

-
1. Yoshinaga I.C, Sedey AL, Coulter DK, Mehl AL. Language of early and later identified children with hearing loss. *Pediatrics* 1998;102:1161-71.
 2. American Academy of Pediatrics, Task force on newborn and infant screening, newborn and infant hearing loss: detection and intervention. *Pediatrics* 1999;103: 527-30.
 3. Joint Committee on Infant Hearing, year 2000 position statement: principles and guidelines for early hearing detection and intervention. *Pediatrics* 2000;106:798-817.
 4. Runjan L, Amir I, Suwento R, Mangunatmadja I. Skrining gangguan pendengaran pada neonatus risiko tinggi. *Sari Pediatri* 2005;6(4):149-54.
 5. Thompson DC, McPhilips H, Davis RL, Lieu TA, Homer CJ, Helfand M. Universal newborn hearing screening. *JAMA* 2001;286:2000-10.
 6. Suwento R. Gangguan pendengaran pada bayi dan anak. Dalam: Seminar Sehari Penatalaksanaan Gangguan Pendengaran Dan Ketulian. Semarang;2007. h.1-12.
 7. World Health Organization, situation review and update on deafness, hearing loss and intervention programmes: proposed plans of action for prevention and alleviation of hearing impairment in countries of the south-east asia region. 2007.
 8. Fortnum HM, Summerfield AQ, Marshall DH, Davis AC, Bamford JM. Prevalence of permanent childhood hearing impairment in the United Kingdom and implications for universal neonatal hearing screening: questionnaire based ascertainment study. *BMJ* 2001;323:1-6.
 9. Jardim IS, Matas CG, Carvalho RMM. Newborn hearing screening with transient evoked otoacoustic emissions and automatic auditory brainstem response. *Einstein* 2008;6:253-61.
 10. Sukadi A. Hiperbilirubinemia. Dalam: Kosim MS, Yunanto A, Dewi R, Sarosa GI, Usman A, editor. *Buku Ajar Neonatologi*. Edisi pertama. Badan penerbit IDAI; 2008. h.147-69.

-
11. American Academy of Pediatrics, Provincial Committee for Quality Improvement. Practice parameter: management of hyperbilirubinemia in the healthy term newborn. *Pediatrics* 1994; 94: 558-65.
 12. Rapin I. Hearing impairment. In: Swaiman K.F, Ashwal S, editors. *Pediatric Neurology Principles and Practice*. 3rd ed. New York. Mosby Inc; 1999. p.77-95
 13. Cunningham M, Cox E O. Hearing assessment in infants and children: recommendations beyond neonatal screening. *Pediatrics* 2003; 111:436-40.
 14. Mason JA, Herrmann KR. Universal infant hearing screening by automated auditory brainstem response measurement. *Pediatrics* 1998;101:221- 8.
 15. Sharma P, Chhangani N.P, Meena K.R, Jora R, Sharma N, Gupta B.D. Brainstem evoked response audiometry (BAER) in neonates with hyperbilirubinemia. *Indian J Pediatr* 2006;73(5):413-6.
 16. Amin SB, Ahlfors C, Orlando MS, Dalzell LE, Merle KS, Guillet R. Bilirubin and serial auditory brainstem response in premature infants. *Indian J Pediatr* 2006;73 (5):413-6.
 17. Ahlfors CE, Parker AE. Unbound bilirubin concentration is associated with abnormal automated auditory brainstem response for jaundiced newborns. *Pediatrics* 2008; 121: 976-8.
 18. MedlinePlus medical encyclopedia: Hearing loss. Diambil dari URL: <http://www.nlm.nih.gov/medlineplus/ency/article/003044.html>.
 19. Schrijver I. Hereditary non-syndromic sensorineural hearing loss. *J Mol Diagn* 2004;6(4):275-84.
 20. Suwento R. Deteksi dini gangguan pendengaran pada anak. Dalam: *Simposium Sehari Anakku Tidak Bisa Mendengar*. Semarang;2006. h.15- 27.
 21. Deka RC, Sarin D. Congenital TORCH infection and hearing loss. In: Deka D, editor. *Congenital Intrauterine TORCH Infections*. 1st ed. New Delhi, Jaypee brothers; 2004. p.112-20.
 22. Rivera LB, Boppana SB, Fowler KB, Britt WJ, Stagno S, Pass RF. Predictors of hearing loss in children with symptomatic congenital cytomegalovirus infection. *Pediatrics* 2002;110:762-7.

-
23. Richardson MP, Reid A, Tarlow MJ, Rudd PT. Hearing loss during bacterial meningitis. *Arch Dis Child* 1997;76:134-8.
 24. Smith RJH, Bale JF, White KR. Sensorineural hearing loss in children. *Lancet* 2005;365:879-90.
 25. Hain CT. Ototoxic medications. Diambil dari URL:
<http://www.tchain.com/otoneurology/disorders/bilat/ototoxins.html>
 26. Volpe JJ. Viral, protozoan, and related intracranial infection. In: Volpe JJ, editor. *Neurology of the Newborn*. 4th ed. Philadelphia: W.B. Saunders Company; 2001. p.717-73.
 27. Maria A, Deorari A. Outcome of neonates born with vertically transmitted TORCH infections. In: Deka D, editor. *Congenital Intrauterine TORCH Infections*. 1st ed. New Delhi, Jaypee brothers; 2004. p.82-98.
 28. Meyer C, Witte J, Hildmann A, Hennecke KH, Schunck KU, Maul K et al. Neonatal screening for hearing disorders in infants at risk: Incidence, risk factors, and follow up. *Pediatrics* 1999;104:900-4.
 29. Chu K, Elimian A, Barbera J, Ogburn P, Spitzer A, Quirk JG. Antecedents of newborn hearing loss. *Obstet Gynecol* 2003; 101: 584-8.
 30. Zamani A, Daneshjou K, Ameni A, Takand J. Estimating the incidence of neonatal hearing loss in high risk neonates. *Acta Medica Iranica* 2004; 42(3): 176- 80.
 31. Beata ACS, Ioana ANCA. Etiopathogeny of congenital and early-onset hearing loss; detection and early intervention methods in infants and children. *Medica* 2008; 3: 45-51.
 32. Widiastuti MI. BERA. *Laboratorium Ilmu Penyakit Saraf FK UNDIP RS Dokter Kariadi Semarang*. 1992.
 33. Abiratno SF. Auditory brainstem response (ABR) prinsip dasar, tehnik pemeriksaan dan penggunaannya dalam klinik. Dalam: Abiratno SF, editor. *Workshop Peranan OAE dan ABR Dalam Klinik*; Surabaya; 2003.
 34. Porter ML, Dennis BL. Hyperbilirubinemia in the term newborn. *AAFP* 2002; 65: 599-606.

-
35. Halamek LP., Stevenson DK. Neonatal jaundice and liver disease. In: Fanaroff AA, Martin RJ, Editors. Neonatal-Perinatal Medicine; Diseases of the Fetus and Infant, 6th ed. New York Mosby-Year Book Inc;1997.p.1345-62.
 36. Dennery PA, Seidman DS, Stevenson DK. Neonatal hyperbilirubinemia. *New Eng J Med* 2001; 344(8):581-90.
 37. Gomella TL, Cunningham MD, Eyal FG, Zenk KE. Hiperbilirubinemia. In: Gomella TL, editor. Neonatology; Management Procedures, On-call Problems, Diseases and Drugs. New York. Lange Medical Book/McGraw-Hill Co; 2004. p. 381-95.
 38. Aminullah A. Ikterus dan hiperbilirubinemia pada neonatus. Dalam : Markum AH, Ismael S, Editor. Buku Ajar Ilmu Kesehatan Anak. Jakarta : FKUI; 1999. h. 313-7.
 39. Soetomenggolo TS, Iman S. Kelainan toksik dan nutrisi. Dalam : Soetomenggolo TS, Ismael S, Editor. Buku Ajar Neurologi Anak, Cetakan ke 2. Jakarta: BP IDAI; 2000. h.541-3.
 40. Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics* 2004;114:297-316.
 41. Gregg RB, Wiorek LS, Arvedson JC. Pediatric audiology: a review. *Pediatr Rev* 2004; 25:224-34.
 42. Shapiro SM. Bilirubin toxicity in the developing nervous system. *Ped Neurol* 2003; 29(5):410-21.
 - 43 . Carter K. Kernicterus in full-term infants-United states,1994-1998. *JAMA* 2001; 286: 299-300.
 44. Shapiro SM, Nakamura H. Bilirubin and the auditory system. *Journal of Perinatology* 2001;21:S52-S55.
 45. Volpe JJ. Bilirubin and brain injury. In: Volpe JJ,editor. *Neurology of The Newborn*. 4th ed. Philadelphia: W.B. Saunders Company; 2001. p. 521-46.

-
46. Alatas H, Karyomanggolo WT, Musa DA, Boediarso A, Oesman IN. Desain penelitian. Dalam: Sastroasmoro S, Ismael S, Editor. Dasar- dasar Metodologi Penelitian Klinis. Edisi 2. Sagung Seto. Jakarta; 2002. h. 79-96.
 47. Taralan T, Soetomenggolo TS, Passat J, Agusman IS. Studi kohort. Dalam: Sastroasmoro S, Ismael S, Editor. Dasar- Dasar Metodologi Penelitian Klinis. Edisi 2. Sagung Seto. Jakarta; 2002. h. 97-109.
 48. Maisels MJ. Neonatal hyperbilirubinemia. In: Klaus MH, Fanaroff AA, editors. Care of The High Risk Neonate, 5th ed. Philadelphia. W.B. Saunders Company; 2001. p.324-62.
 49. Tioseco JA, Aly H, Milner J, Patel K, El-Mohandes AAE. Does gender affect neonatal hyperbilirubinemia in infants? *Pediatr Crit Care Med* 2005; 6: 171-4.
 50. Boo NY, Oakes M, Lye MS, Said H. Risk factors associated with hearing loss in term neonates with hyperbilirubinaemia. *J Trop Pediatr* 1994;40:194-7.
 51. Moore DR. Auditory development and the role of experience. *BMB* 2002;63:171-81.
 52. Madden C, Rutter M, Hilbert L, Greinwald JH, Choo DI. Clinical and audiological features in auditory neuropathy, *Arch Otolaryngol Head Neck Surg* 2002;128:1026-30.
 53. Fallon JB, Irvine DRF, Shepherd RK. Cochlear implants and brain plasticity. *Hear res* 2008;238:110-7.
 54. Holster IR, Hoeve LJ, Wieringa MH, Willis-Lorrier RMS, Gier HHW. Evaluation of hearing loss after failed neonatal hearing screening. *J Pediatr* 2009;155:646-50.
 55. Oysu C, Ulubil A, Aslan I, Baserer N. Incidence of cochlear involvement in hyperbilirubinemic deafness. *Ann Otol Rhinol Laryngol* 2002;111:1021-4.
 56. Boo NY, Rohani AJ, Asma A. detection of sensorineural hearing loss using automated auditory brainstem-evoked response and transient-evoked otoacoustic emission in term neonates with severe hyperbilirubinemia. *Singapore Med J* 2008; 49:209- 14.

57 .Wennberg RP, Ahlfors CE, Bhutani VK, Johnson LH, Shapiro SM. Toward understanding kernicterus: a challenge to improve the management of jaundiced newborns. *Pediatrics* 2006;117;474-85.