

## DAFTAR PUSTAKA

1. Hadders-Algra M. Evaluation of motor function in young infants by means of the assessments of general Movements: a review. *Ped Phys Ther* 2001; 13 : 27-36.
2. Hadders-Algra M, Klip van den Nieuwendijk AWJ, Martijn A, Van Eykern LA. Assesment of general movements : toward better understanding of a sensitive method to evaluate brain functionin in young infants. *Dev Med Child Neurol*,1997; 39 :88-98.
3. Prechtl HFR.General movements assesment as a method of developmental neurology : new paradigmas and their consequences. *Dev Med Child Neurol* 2001; 43: 836-42
4. Prechtl HFR.Qualitative changes of spontaneus movements in fetus and preterm infants are a marker of neurological disfunction.*Early Hum Dev* 1990;23:151-9
5. Hadders-Algra M, Prechtl HFR.Developmental course of general movements in early infancy, I: descriptive analysis of change in form. *Early Hum Dev* 1992; 28: 201-14
6. Bos AF, Einspieler C, Prechtl HFR. Intrauterine growth retardation, general movements and neurodevelopmental outcome: a review. *Dev Med Child Neurol* 2001; 43: 61-8
7. Prechtl HFR,Ferrari F,Cioni G.predictive value of general movements in asphyxiated fullterm infants. *Early Hum Dev* 1993;35:91-120.
8. Soorani-Lunsing I, Woltl HA, Hadders-Algra M. Are moderate hyperbilirubinemia in healthy term neonate really safe for the brain ? *Pediatr Res* 2001 ; 50 : 701-5
9. Porter ML, Dennis BL. Hyperbilirubinemia in the Term Newborn. *Am Fam Phy* 2002. 65: 599-606.

10. Aminullah A. Ikterus dan hiperbilirubinemia pada neonatus. Dalam : Markum AH, Ismael S, editor. Buku ajar Ilmu Kesehatan Anak. Jakarta : FKUI ; 1999. h.313-7.
11. American Academy of Pediatrics, Provincial Committee for Quality Improvement. Practice parameter: management of hyperbilirubinemia in the healthy term newborn. *Pediatrics* 1994; 94: 558-65.
12. Gomella TL. Hyperbilirubinemia. In: Gomella TL, Cunningham MD, Eyal FG, Zenk KE, editors. *Neonatology management, procedures, on-call problems, diseases and drugs*. 5th Ed. New York: McGraw Hill; 2004. p.381-95.
13. Uhudiah U, Oktavia D. Pemberian terapi sinar berdasarkan penilaian klinis pada neonatus dengan hiperbilirubinemia. Dalam : Rusdidjas, Tjipta GD, Dimyati Y, editor. *Kongres Nasional VIII Perinasia & Simposium Internasional*. Di Medan: Oktober 5-8, Perinasia ; 2003. h.74-81.
14. Indiarso F. Tranfusi tukar pada neonatus dengan hiperbilirubinemia. Dalam : Rusdidjas, Tjipta GD, Dimyati Y, editor. *Kongres Nasional VIII Perinasia & Simposium International*. Medan: Oktober 5-8, Perinasia ; 2003. h.84-98.
15. Halamek LP, Stevenson DK. Neonatal jaundice and liver disease. In : Fanaroff AA, Martin RJ, editor. *Neonatal-perinatal medicine ; Diseases of the fetus and infant*. 6th Ed. New York: Mosby-Year Book Inc ; 1997. p.1345-62.
16. Hadders-Algra M, Stremmelaar EF, Martijn A. Quality of general movements and the development of minor neurological dysfunction at toddler and school age. *Clin Rehab* 2004; 18:287-9.
17. Groen SE, Blecourt ACE, Postema K, Hadders-Algra M. General movements in early infancy predict neuromotor development at 9 to 12 years of age. *Dev Med Child Neurol* 2005; 43: 731-8
18. Ferrari F, Cioni G, Einspieler C, Roversi MF, Bos AF, Paolicelli PB, et all. Cramped Syncronized General Movements in Preterm as an Early Marker for Cerebral Palsy. *Arch Pediatr Adolesc Med* 2002; 156 : 460-7

19. Witarini KA, Etika R, Suryawan A, Indarso F, Irwanto, et al. Evaluasi penggunaan Metode Prechtel untuk menilai kualitas gerakan spontan bayi muda sehat: pengalaman Surabaya. Dalam : Sadjimin T, Juffrie M, Julia M, Wibowo T, editor. Buku Naskah Lengkap PIT IKA III Yogyakarta. 6-9 Mei 2007. Yogyakarta: IDAI Pusat ; 2007: 261.
20. Murray RK. Porfirin dan Pigmen Empedu. In : Biokomia Harper . Edisi 24. Jakarta: Penerbit Buku Kedokteran EGC; 1997:352-66
21. Kliegman RM. Ikterus dan hiperbilirubinemia pada bayi baru lahir. In: Behrman, Kliegman, Arvin, editor. Nelson Textbook of Pediatric. 15th Edition. Philadelphia: WB Saunders Co;2000 : 610-16.
22. Shapiro SM. Bilirubin Toxicity in the Developing Nervous System. Ped Neuro 2003 ; 29 : 410-21.
23. Stevenson DK, Fanaroff AA, Maisels MJ, Young BW, Wong RJ, Vreman HJ, et al. Prediction of Hyperbilirubinemia in Near-term and Term infants. Pediatrics 2001 ; 108 (1) : 31-9.
24. Hansen T. Mechanism of Bilirubin Toxicity: Clinical Implication. Clinical Perinatology 2002; 29: 765-78.
25. Amin SB, Ahlfors C, Orlando MS, Dalzell E, Merle KS, Guillet R. Bilirubin and Serial Auditory Brainstem Responses in Premature Infants. Pediatrics 2001; 107 (4): 667-70.
26. Dennery PA, Seidman DS, Stevenson DK. Neonatal Hyperbilirubinemia. The new Eng J Med, February 2001(8):581-90
27. Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. Pediatrics 2004;114:297-316.
28. Soemantri Ag. Biomolecular of Red Cell Glucose-6-Phosphate Dehydrogenase Deficiency of Asia Population. In: Wandita S, Herini ES, Surjono. Editor: Asian Symposium In Neonatology G6PD Deficiency and Related Condition. Yogyakarta, Agustus 8 – 9, 2000: 1 – 27
29. Hadders-Algra M. General movements: a window for early identification of children at high risk for developmental disorders. JPediatr 2004;145: 512-8

30. Hadders-Algra M. The assessments of General Movements is a valuable technique for the detection of brain dysfunction in young infants. A review. *Acta Paediatr Suppl*; 1996; 416 : 39-43
31. Garcia JM, Gherpelli JLD, Leone CR. The role of spontaneous general movements assessment in the neurological outcome of cerebral lesion in preterm infants. *J Pediatr (Rio J)* 2004; 80(4):296-304.
32. Hadders-Algra M, Groothius AMC. Quality of general movements in infancy is related to the development of neurological dysfunction, attention deficit hyperactivity disorder and aggressive behavior. *Dev Med Child Neurol* 1999; 41: 381-91.
33. Ferrari F, Cioni G, Prechtl HFR. Qualitative changes of general movements in preterm infants with brain lesion. *Early Hum Dev* 1990; 23: 193-223.
34. Cioni G, Ferrari F, Prechtl HFR. Posture and motility in fullterm infants. *Early Hum* 1989; 18: 247-62
35. Einspieler C, Prechtl HFR, Bos AF, Ferrari F, Cioni G. Prechtl's Method on the Qualitative Assessment of General Movements in preterm, term, and young infants. London: Mac Keith Press; 2004: 19-28, 35-45, 67-71.
36. Hansen TWR, Tommarello S, Allen JW. Subcellular Localization of Bilirubin in Rat brain after *invivo iv* administration of [H] Bilirubin. *Ped Res* 2001; 49: 203-7.
37. Ostrow JD, Pacolo L, Shapiro SM, Tiribelli C. New Concept in Bilirubin Encephalopathy. *Eur J Clin Invest*. 2003; 33: 988-997.
38. Rodrigues CMP, Sola S, Castro RE, Laires PA, Brites D. Perturbation of membrane dynamics in nerve cells as an early event during bilirubin-induced apoptosis. *J Lipid Res* 2002; 43: 885-94.
39. Silva RF, Rodrigues CM, Brites D. Bilirubin-induced Apoptosis in Cultured Rat Neural Cells is Aggravated by Chenodeoxycholic Acid but Prevented by Ursodeoxycholic Acid. *J Hepatol* 2001; 34 (3): 402-8.
40. Rodrigues CMP, Sola S, Brites D. Bilirubin Induces Apoptosis via the Mitochondrial Pathway in Developing Rat Brain Neurons. *Hepatol* 2002 ; 35 : 1186-95.

41. Madiyono B, Moeslichan S, Sastroasmoro S, Budiman I, Purwanto SH. Perkiraan besar sample. Dalam: Sastroasmoro S, Ismael S, Penyunting. Dasar-dasar metodologi penelitian klinis, edisi 2. Jakarta: Sagung seto; 2002: 259-87.
42. Ip S, Chung M, Kulig J, O'Brien R, Sege R, Glickin S, et all. An-evidence based review of important issues concerning neonatal Hyperbilirubinemia. Pediatrics 2004 ; 114(1) : e130-e53.
43. Tioseco JA, Aly H, Milner J, Patel K, El-Mohandes AAE. Does gender affect neonatal hyperbilirubinemia in infants? Pediatr Crit Care Med 2005; 6: 171-4.
44. Molkenboer JF, Roumen FJ, Smits LJ, Nijhuis JG. Birth weight and neurodevelopmental outcome of children at 2 years of age after planned vaginal delivery for breech presentation at term. Am J Obstet Gynecol 2007;196(2):15-16.
45. Golfier F, Vaudoyer F, Ecochard R, Champion F, Audra P, Raudrant D. Planned vaginal delivery versus elective caesarean section in singleton term breech presentation: a study of 1116 cases. Eur J Obst Gyn Reprod Biol. 2001 Oct; 98(2):186-92.
46. Bouwstra H, Boersma ER, Boehm G, Dijck-Brouwer DAJ, Muskiet FAJ, Hadders Algra M. Exclusive Breastfeeding of Healthy Term Infants for at least 6 weeks improves neurological condition. J. Nutr. 2003 ; 133 : 4243-4245
47. Bouwstra H, Dijk-Stigter GR, Grootenhuis HMJ, Janssen-Plas FEM, van Belle, Hadders-Algra M, et all. Prevalence of abnormal general movements in three month old infants. Early Hum Dev 2009; 85: 399-403.
48. Hadders-Algra M. Putative neural substrates of normal and abnormal general movements. J Neurosci 2007; 31 : 1181-90
49. De Graaf-Peters VB, Hadders-Algra M. Ontogeny of the human central nervous system : what is happening when ? Early Hum Dev 2006;82 : 257-66.

50. Miller SP, Latal B, Clark H, Barnwell A, Glidden D, Barkovich J, dkk. Clinical Sign Predict 30-month Neurodevelopmental Outcome after Neonatal Encephalopathy. *Am J Obst and Gyn* 2004; 190: 93-9.
51. Moster D, Lie TR, Markestad T. Joint Association of Apgar Scores and Early Neonatal Symptoms with Minor Disabilities at School Age. *Arch Dis Child Fetal Neonatal* 2002 ; 86 : 16-21
52. Wang X, Rousset CI, hagberg H. Lipopolysacharide-induced infalamation and perinatal brain injury. *Semin Fetal Neonatal Med.* 2006; 11 : 343-53
53. Johnson MV, Hoon AH. Possible mechanism in infants for selective basal ganglia damage from asphyxia, kernicterus, or mitochondrial encephalopathies. *J Child Neurol* 2000;15(9): 588-91.
54. Groenendaal F, Grond J, Vries LS. Cerebral metabolism in severe neonatal hyperbilirubinemia. *Pediatrics* 2004; 114(1): 291-4.
55. Soetomenggolo TS, Iman S. Kelainan Toksik dan Nutrisi. Dalam: Soetomenggolo TS, Ismael S, editor. Buku ajar Neurologi Anak. Edisi-2. Jakarta : BP IDAI ; 2000: 541-3.
56. Connolly AM, Volpe II. Clinical feature of bilirubin encephalopathy. *Clin Perinatol* 1990; 17: 371-79.
57. Turkel SB. Autopsy findinga associated with neonatal hiperbilirubinemia *Clin Perinatol* 1990; 2: 381-94.
58. Newman TB, Klebanoff MA. Neonatal hyperbilirubinemia and long-term out- come: another look at the collaborative perinatal project. *Pediatrics* 1993; 92:651–57.
59. Grimmer I, Berger-Jones K, Bührer C, Brandl U, Obladen M. Late neurological sequela of non-hemolytic hyperbilirubinemia of healthy term neonates. *Acta Paediatr* 1999; 88:661– 663.
60. Maisels MJ, Newman TB. Predicting hyperbilirubinemia in newborns: the importance of timing. *Pediatrics* 1999; 103:493–95.
61. Wu PYK, Hodgman JE, Kirkpatrick BV, White NB, Phil M, Bryla DA. Metabolic aspects of phototherapy. *Pediatrics* 1985; 75(suppl): 427– 33.

62. Martinez JC, Maisels J, Otheguy L, Garcia H, Savorani M, Mogni B. Hyperbilirubinemia in the breast-fed newborn: a controlled trial of four interventions. *Pediatrics* 1993; 91:470–73.
63. Valaes T, Koliopoulos T, Koltsidopoulos A. The impact of phototherapy in the management of neonatal hyperbilirubinemia: a comparison of historical cohorts. *Acta Paediatr* 1996; 85:273–76.
64. Amato M, Donati F. Cerebral blood flow velocity in term infants treated with phototherapy. *Brain Dev* 1991; 13:417– 19.
65. Benders MJ, Van Bel F, Van de Bor M. Haemodynamic consequences of phototherapy in term infants. *Eur J Pediatr* 1999; 158:323–28.
66. Tan KL, Skurr BA, Yip YY. Phototherapy and the brain-stem auditory evoked response in neonatal hyperbilirubinemia. *J Pediatr* 1992; 120:306–8.
67. Seidman DS, Paz I, Stevenson DK, Laor A, Danon YL, Gale R. The effect of phototherapy for neonatal hyperbilirubinemia on long term cognitive performance. *Pediatr Res* 1991; 29:265A.
68. Bouwstra H, Dijk-Stigter GR, Grootenhuis M, Janssen-Plas FEM, van Belle A, Hadders-Algra M. Prevalence of abnormal general movements in three months old infants. *Early Hum Dev* 2009; 85: 399-403.
69. Constaninov JC, Adamson-Macedo EN, Mirmisan M, Fleisher BE. Movement, imaging and neurobehavioral assessment as predictors cerebral palsy in preterm infants. *J Perinatol* 2007; 27 : 225-9.
70. Maurizio Romeo DM, Guzzetta A, Scoto M, Cioni M, Patusi P, Mazzone D, et all. Early neurologic assessment in preterm infants: integration of traditional neurologic examination and observation of general movements. *Eur J Pediatr Neurol* 2008; 12: 183-9

- 
1. Hadders-Algra M. Evaluation of motor function in young infants by means of the assessments of general Movements: a review. *Ped Phys Ther* 2001; 13 : 27-36.
  - 2 Hadders-Algra M, Klip van den Nieuwendijk AWJ, Martijn A, Van Eykern LA. Assesment of general movements : toward better understanding of a sensitive method to evaluate brain functionin in young infants. *Dev Med Child Neurol*,1997; 39 :88-98.
  - 3 Prechtl HFR.General movements assesment as a method of developmental neurology : new paradigmas and their consequences. *Dev Med Child Neurol* 2001; 43: 836-42
  - 4 Prechtl HFR.Qualitative changes of spontaneus movements in fetus and preterm infants are a marker of neurological dysfunction.*Early Hum Dev* 1990;23:151-9
  - 5 Hadders-Algra M, Prechtl HFR.Developmental course of general movements in early infancy, I: descriptive analysis of change in form. *Early Hum Dev* 1992; 28: 201-14
  - 6 Bos AF, Einspieler C, Prechtl HFR. Intrauterine growth retardation, general movements and neurodevelopmental outcome: a review. *Dev Med Child Neurol* 2001; 43: 61-8
  - 7 Prechtl HFR,Ferrari F,Cioni G.predictive value of general movements in asphyxiated fullterm infants. *Early Hum Dev* 1993;35:91-120.
  - 8 Soorani-Lunsing I, Woltl HA, Hadders-Algra M. Are moderate hyperbilirubinemia in healthy term neonate really safe for the brain ? *Pediatr Res* 2001 ; 50 : 701-5
  - 9 Porter ML, Dennis BL. Hyperbilirubinemia in the Term Newborn. *American Family Physician* 2002. 65: p 599-606.
  - 10 Aminullah A. Ikterus dan hiperbilirubinemia pada neonatus. Dalam : Markum AH, Ismael S, editor. Buku ajar Ilmu Kesehatan Anak. Jakarta : FKUI ; 1999. h.313-7.
  - 11 American Academy of Pediatrics, Provincial Committee for Quality Improvement. Practice parameter: management of hyperbilirubinemia in the healthy term newborn.*Pediatrics* 1994; 94: 558-65.
  - 12 Gomella TL. Hyperbilirubinemia. In: Gomella TL, Cunningham MD, Eyal FG, Zenk KE,editors. *Neonatology management, procedures, on-call problems, diseases and drugs*. 5th Ed. New York: McGaw Hill; 2004. p.381-95.
  - 13 Uhudiah U, Oktavia D. Pemberian terapi sinar berdasarkan penilaian klinis pada neonatus dengan hiperbilirubinemia. Dalam : Rusdidjas, Tjipta GD, Dimyati Y, editor. *Kongres Nasional VIII Perinasia & Simposium Internasional*. Medan : Perinasia ; 2003. h.74-81.
  - 14 Indiarso F. Tranfusi tukar pada neonatus dengan hiperbilirubinemia. Dalam : Rusdidjas, Tjipta GD, Dimyati Y, editor. *Kongres Nasional VIII Perinasia & Simposium International*. Medan : Perinasia ; 2003. h.84-98.
  - 15 Halamek LP, Stevenson DK. Neonatal jaundice and liver disease. In : Fanaroff AA, Martin RJ, editor. *Neonatal-perinatal medicine ; Diseases of the fetus and infant*. 6th Ed. New York: Mosby-Year Book Inc ; 1997. p.1345-62.
  - 16 Hadders-Algra M, Stremmelaar EF, Martjin A. Quality of general movements and the development of minor neurological dysfunction at toddler and school age. *Clnical Rehabilitation* 2004; 18:287-9.
  - 17 Groen SE, Blecourt ACE, Postema K, Hadders-Algra M. General movements in early infancy predict neuromotor development at 9 to 12 years of age. *Dev Med Child Neurol* 2005; 43: 731-8
  - 18 Ferrari F,Cioni G, Einspieler C, Roversi MF, Bos AF , Paolicelli PB, et all. Cramped Syncronized General Movements in Preterm as an Early Marker for Cerebral Palsy. *Arch Pediatr Adolesc Med* 2002; 156 : 460-7
  - 19 Witarini KA, Etika R, Suryawan A, Indarso F,Irwanto,et al.Evaluasi penggunaan Metode Prechtl untuk menilai kualitas gerakan spontan bayi muda sehat: pengalaman

- 
- Surabaya. Dalam : Sadjimin T, Juffrie M, Julia M, Wibowo T, editor. Buku Naskah Lengkap PIT IKA III Yogyakarta. Mei 2007. Yogyakarta: IDAI Pusat ; 2007: h.261-127.
- 20 Murray RK. Porfirin dan Pigmen Empedu. In : Biokomia Harper . Edisi 24. Jakarta: Penerbit Buku Kedokteran EGC; 1997. p. 352-66
- 21 Kliegman RM. Ikterus dan hiperbilirubinemia pada bayi baru lahir. In: Behrman, Kliegman, Arvin, editor. Nelson Textbook of Pediatric. 15th Edition. Philadelphia : WB Saunders Co ; 2000. p.610-16.
- 22 Shapiro SM. Bilirubin Toxicity in the Developing Nervous System. Pediatric Neurology 2003 ; 29 : 410-21.
- 23 Stevenson DK, Fanaroff AA, Maisels MJ, Young BW, Wong RJ, Vreman HJ, et al. Prediction of Hyperbilirubinemia in Near-term and Term infants. Pediatrics 2001 ; 108 (1) : 31-9.
- 24 Hansen T. Mechanism of Bilirubin Toxicity : Clinical Implication. Clinical Perinatology 2002 ; 29 : 765-78.
- 25 Amin SB, Ahlfors C, Orlando MS, Dalzell E, Merle KS, Guillet R. Bilirubin and Serial Auditory Brainstem Responses in Premature Infants. Pediatrics 2001; 107 (4).p. 667-70.
- 26 Dennery PA, Seidman DS, Stevenson DK. Neonatal Hyperbilirubinemia. The new England Journal of Medicine, February 2001 (8), Vol 344:581-590
- 27 Subcommittee on Hyperbilirubinemia. Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. Pediatrics 2004;114:297-316.
- 28 Soemantri Ag. Biomolecular of Red Cell Glucose-6-Phosphate Dehydrogenase Deficiency of Asia Population. In : Wandita S, Herini ES, Surjono. Editor: Asian Symposium In Neonatology G6PD Deficiency and Related Condition. Yogyakarta, Agustus 8 – 9, 2000. p. 1 – 27
- 29 Hadders-Algra M. General movements:a window for early identification of children at high risk for developmental disorders.JPediatr 2004;145: 512-8
- 30 Hadders-Algra M. The assessments of General Movements is a valueable techique for the detection of brain dysfunction in young infants. A review. Acta Paediatr Suppl; 1996; 416 : 39-43
- 31 Garcia JM, Gherpelli JLD,Leone CR. The role of spontaneous general movements assesment in the neurological outcome of cerebral lesion in preterm infants. JPediatr ( RioJ) 2004;80(4):296-304.
- 32 Hadders-Algra M, Groothius AMC.Quality of general movements in infancy is related to the development of neurological dysfunction, attention deficit hyperactivity disorder and aggressive behavior. Dev Med Child Neurol 1999;41: 381-91.
- 33 Ferrari F,Cioni G,Prechtl HFR.Qualitative changes of general movements in preterm infants with brain lession. Early Hum Dev 1990;23:193-223.
- 34 Cioni G, Ferrari F, Prechtl HFR. Posture and motility in fullterm infants. Early Hum 1989; 18: 247-62
- 35 Einspieler C, Prechtl HFR,Bos AF,Ferrari F,Cioni G. Prechtl's Method on the Qualitative Assesment of General Movements in preterm,term, and young infants.London: Mac Keith Press; 2004.p: 19-28, 35-45,67-71.
- 36 Hansen TWR, Tommarello S, Allen JW. Subcellular Localization of Bilirubin in Rat brain after invivo iv administration of [H] Bilirubin. Pediatric Research 2001; 49: 203-7.
- 37 Ostrow JD, Paolo L, Shapiro SM, Tiribelli C. New Concept in Bilirubin Encephalopathy. EurJClin Invest. 2003 ; 33 : 988-997.

- 
- 38 Rodrigues CMP, Sola S, Castro RE, Laires PA, Brites D. Perturbation of membrane dynamics in nerve cells as an early event during bilirubin-induced apoptosis. *J Lipid Research* 2002 ; 43 : 885-94.
- 39 Silva RF, Rodrigues CM, Brites D. Bilirubin-induced Apoptosis in Cultured Rat Neural Cells is Aggravated by Chenodeoxycholic Acid but Prevented by Ursodeoxycholic Acid. *J Hepatology* 2001 ; 34 (3) : 402-8.
- 40 Rodrigues CMP, Sola S, Brites D. Bilirubin Induces Apoptosis via the Mitochondrial Pathway in Developing Rat Brain Neurons. *Hepatol* 2002 ; 35 : 1186-95.
- 41 Madiyono B, Moeslichan S, Sastroasmoro S, Budiman I, Purwanto SH. Perkiraan besar sample. Dalam: Sastroasmoro S, Ismael S, Penyunting. Dasar-dasar metodologi penelitian klinis, edisi 2. Jakarta: Sagung seto; 2002. h.259-287.
- 42 Ip S, Chung M, Kulig J, O'Brien R, Sege R, Glickman S, dkk. An-evidence based review of important issues concerning neonatal Hyperbilirubinemia. *Pediatrics* 2004 ; 114(1) : e130-e53.
- 43 Tioseco JA, Aly H, Milner J, Patel K, El-Mohandes AAE. Does gender affect neonatal hyperbilirubinemia in infants? *Pediatr Crit Care Med* 2005; 6: 171-4.
- <sup>44</sup> Molkenboer JF, Roumen FJ, Smits LJ, Nijhuis JG. Birth weight and neurodevelopmental outcome of children at 2 years of age after planned vaginal delivery for breech presentation at term. *Am J Obstet Gynecol* 2007;196(2):15-16.
- <sup>45</sup> Golfier F, Vaudoyer F, Ecochard R, Champion F, Audra P, Raudrant D. Planned vaginal delivery versus elective caesarean section in singleton term breech presentation: a study of 1116 cases. *Eur J Obstet Gynecol Reprod Biol*. 2001 Oct;98(2):186-92.
- 46 Miller SP, Latal B, Clark H, Barnwell A, Glidden D, Barkovich J, dkk. Clinical Sign Predict 30-month Neurodevelopmental Outcome after Neonatal Encephalopathy. *American Journal of Obstetrics and Gynecology* 2004; 190: 93-9.
- 47 Moster D, Lie TR, Markestad T. Joint Association of Apgar Scores and Early Neonatal Symptoms with Minor Disabilities at School Age. *Arch Dis Child Fetal Neonatal* 2002 ; 86 : 16-21
- 49 Hadders-Algra M. Putative neural substrates of normal and abnormal general movements. *J Neuropediatr* 2007; 31 : 1181-1190
- 50 De Graaf-Peters VB, Hadders-Algra M. Ontogeny of the human central nervous system : what is happening when ? *Early Hum Dev* 2006;82 : 257-266.
- 51 Johnson MV, Hoon AH. Possible mechanism in infants for selective basal ganglia damage from asphyxia, kernicterus, or mitochondrial encephalopathies. *J Child Neurology* 2000 ; 15(9) : 588-91.
- 52 Groenendaal F, Grond J, Vries LS. Cerebral metabolism in severe neonatal hyperbilirubinemia. *Pediatrics* 2004 ; 114(1) : 291-4.
- 53 Soetomenggolo TS, Iman S. Kelainan Toksik dan Nutrisi. Dalam : Soetomenggolo TS, Ismael S, editor. Buku ajar Neurologi Anak. Edisi-2. Jakarta : BP IDAI ; 2000: 541-3.
- 54 Connolly AM, Volpe JJ 1990. Clinical feature of bilirubin encephalopathy. *Clin Perinatol* 17: 371-79
- 55 Turkel SB. 1990. Autopsy findings associated with neonatal hyperbilirubinemia *Clin Perinatol* 2: 381-94
- 56 Newman TB, Klebanoff MA 1993. Neonatal hyperbilirubinemia and long-term outcome: another look at the collaborative perinatal project. *Pediatrics* 92:651–657

- 
- 57 Grimmer I, Berger-Jones K, Bührer C, Brandl U, Obladen M 1999 Late neurological sequela of non-hemolytic hyperbilirubinemia of healthy term neonates. *Acta Paediatr* 88:661– 663
- 58 Maisels MJ, Newman TB 1999 Predicting hyperbilirubinemia in newborns: the importance of timing. *Pediatrics* 103:493– 495
- 59 Wu PYK, Hodgman JE, Kirkpatrick BV, White NB, Phil M, Bryla DA 1985 Metabolic aspects of phototherapy. *Pediatrics* 75(suppl):427– 433
- 60 Martinez JC, Maisels J, Otheguy L, Garcia H, Savorani M, Mogni B, Martinez JC 1993 Hyperbilirubinemia in the breast-fed newborn: a controlled trial of four interventions. *Pediatrics* 91:470 – 473.
- 61 Valaes T, Koliopoulos T, Koltsidopoulos A 1996 The impact of phototherapy in the management of neonatal hyperbilirubinemia: a comparison of historical cohorts. *Acta Paediatr* 85:273–276 .
- 62 Amato M, Donati F 1991 Cerebral blood flow velocity in term infants treated with phototherapy. *Brain Dev* 13:417– 419
- 63 Benders MJ, Van Bel F, Van de Bor M 1999 Haemodynamic consequences of phototherapy in term infants. *Eur J Pediatr* 158:323–328.
- 64 Tan KL, Skurr BA, Yip YY 1992 Phototherapy and the brain-stem auditory evoked response in neonatal hyperbilirubinemia. *J Pediatr* 120:306–308.
- 65 Seidman DS, Paz I, Stevenson DK, Laor A, Danon YL, Gale R 1991 The effect of phototherapy for neonatal hyperbilirubinemia on long term cognitive performance. *Pediatr Res* 29:265A