

## DAFTAR PUSTAKA

1. Thompson DC, McPhilips H, Davis RL, Lieu TL, Homer CJ, Helfand M. Universal Newborn Hearing Screening. *JAMA* 2001; 286:2000-10.
2. Runjan L, Amir I, Suwento R. Skrining gangguan pendengaran pada neonatus risiko tinggi. *Sari Pediatri* 2005 Maret; 6(4):149-54.
3. Suwento R. Gangguan pendengaran pada bayi dan anak. Dalam seminar sehari penatalaksanaan gangguan pendengaran dan ketulian; Semarang; 2007 Februari.h.1-12.
4. World Health Organization, situation review and update on deafness, hearing loss and intervention programmes: proposed plans of action for prevention and alleviation of hearing impairment in countries of the south-east asia region. 2007.
5. Fortnum H M, Summerfield A Q, Marshall DH, Davis AC, Bamford JM. Prevalence of permanent childhood hearing impairment in the United Kingdom and implications for universal neonatal hearing screening: questionnaire based ascertainment study. *BMJ* 2001;323:1-6.
6. Anonim. Rekapitulasi angka kelahiran bayi berdasarkan rekam medis RSUP Dr. Kariadi Semarang tahun 2008. Tidak dipublikasikan.
7. Rapin I. Hearing impairment. Dalam: Swaiman K.F, Ashwal S, penyunting. *Pediatric Neurology Principles and Practice*. Edisi ke-1. New York: Mosby Inc, 1999. h.77- 95
8. Yoshinaga-Itano C, Sedey AL, Coulter DK, Mehl AL. Language of early and later identified children with hearing loss. *Pediatrics* 1998;102:1161-71.
9. American Academy of Pediatrics. Task force on newborn and infant screening, newborn and infant hearing loss:detection and intervention. *Pediatrics* 1999;103:527-30.
10. Joint Committee on Infant Hearing. Year 2007 position statement: principles and guidelines for early hearing detection and intervention programs. *Pediatrics* 2007;120:898-921.

11. Dharmasetiawani N. Asfiksia dan resusitasi bayi baru lahir. Dalam: Kosim MS, Yunanto A, Dewi R, Sarosa GI, Usman A, penyunting. Buku Ajar Neonatologi. Edisi ke-1. Jakarta: IDAI, 2008. h.103-25.
12. Khreisat WH, Habahbeh Z. Risk factors of birth asphyxia: A study at Prince Ali Ben Al-Hussein Hospital. *Pakistan Journal of Medical Science* 2005; 21:30-2.
13. Menkes JH, Sarnat HB. Perinatal asphyxia and trauma. Edisi ke-6. Philadelphia: Lippincott Williams & Wilkins, 2000.h.401-66
14. Perlman JM. Markers of asphyxia and neonatal brain injury. *N Engl J Med* 1999, 341:363-5.
15. Koyama S, Kaga K, Sakata H, Iino Y, Kodera K. Pathological findings in the temporal bone of newborn infants with neonatal asphyxia. *Acta Oto-Laryngologica* 2005;125:1028-32.
16. Jiang ZD, Yin R, Shao XM, Wilkinson AR. Brain-stem auditory impairment during the neonatal period in term infants after asphyxia: dynamic changes in brain-stem auditory evoked response to clicks of different rates. *Clin Neurophysiol* 2004;115:1605-15.
17. Meyer C, Witte J, Hildmann A, Hennecke KH, Schunck KU, Maul K dkk. Neonatal screening for hearing disorders in infants at risk: Incidence, risk factors, and follow up. *Pediatrics* 1999;104:900-4.
18. Cunningham M, Cox EO. Hearing assessment in infants and children : Recommendations beyond neonatal screening. *Pediatrics* 2003;111:436-40.
19. Stearn N, Swanepoel DW. Identifying hearing loss by means of iridology. *African Journal of Traditional, Complimentary and Alternative Medicines* 2007;4(2):205-14.
20. Kilic I, Karahan H, Kurt T, Ergin H, Sahiner T. Brainstem evoked response audiometry and risk factors in premature infants. *Marmara Medical J* 2007;20(1):21-8.
21. Zang Z, Wilkinson AR, Jiang ZD. Distortion product otoacoustic emissions at 6 months in term infants after perinatal hypoxia-ischaemia or with a low apgar score. *Eur J Pediatr* 2008; 167:575–578.

22. Liston SI, Duval AJ. Dasar-dasar otolaringologi. Dalam: Effendi H, penyunting. Buku Ajar Penyakit THT. Edisi 6. Jakarta: EGC; 1997.h.27-38.
23. Christobal R, Oghalai JS. Hearing loss in children with ver low birth weight: current review of epidemiology and pathophysiology. Arch. Dis. Child. Fetal Neonatal 2008,93:462-8.
24. Lasky RE, Williams AL. The development of the auditory system from conception to term. NeoReviews 2005;6(3): 141-52.
25. Hain TC. Hearing loss. Diunduh dari URL: <http://www.dizziness-and-balance.com/disorders/hearing/hearing.html> tanggal 11 Agustus 2009.
26. Suwento R. Deteksi dini gangguan pendengaran pada anak. Dalam: Simposium sehari anakku tidak bisa mendengar; Semarang; 2006 Februari.h.15- 27.
27. Gifford KA, Holmes MG, Bernstein HH. Hearing loss in children. Pediatrics in Review. 2009; 30:207-16.
28. Deka RC, Sarin D. Congenital TORCH infection and hearing loss. Dalam: Deka D, penyunting. Congenital Intrauterine TORCH Infections.Edisi ke-1. New Delhi: Jaypee brothers, 2004.h.112-20.
29. Rivera LB, Boppana SB, Fowler KB, Britt WJ, Stagno S, Pass RF. Predictors of hearing loss in children with symptomatic congenital cytomegalovirus infection. Pediatrics 2002; 110:762-7.
30. Richardson MP, Reid A, Tarlow MJ, Rudd PT. Hearing loss during bacterial meningitis. Arch Dis Child 1997;76:134-8.
31. Hain TC. Ototoxic medications. Diunduh dari URL: <http://www.tchain.com/otoneurology/disorders/bilat/ototoxins.html> tanggal 10 Agustus 2009.
32. de Hoog M, van den Anker JN. Therapeutic drug monitoring of aminoglycoside in neonates. Clin Pharmacokinet 2009;48(5):343-5.
33. Maria A, Deorari A. Outcome of neonatus born with vertical transmitted TORCH infections. 1<sup>st</sup> Ed. New Delhi. Jaypee brothers; 2004. h. 82-98.

34. Zamani A, Daneshjou K, Ameni A, Takand J. Estimating the incidence of neonatal hearing loss in high risk neonates. *Acta Medica Iranica* 2004; 42(3):176-80.
35. Kolegium Ilmu Kesehatan Telinga Hidung Tenggorok, Bedah Kepala dan Leher. Modul telinga, gangguan pendengaran. Edisi 1. 2008.h.18-23.
36. Abiratno SF. Auditory brainstem response (ABR) prinsip dasar, teknik pemeriksaan dan penggunaannya dalam klinik. Dalam: Abiratno SF, penyunting. *Workshop Peranan OAE dan ABR Dalam Klinik*; Surabaya; 2003.
37. WHO. *Perinatal Mortality : a listing of available information*. FRH/MSM. Geneva: WHO; 1996.
38. Kosim MS. *Asfiksia neonatorum*. SMF Kes. Anak RSUP Dr. Kariadi / Bag. IKA FK UNDIP Semarang. 2005
39. American Academy of Pediatrics, Committee on Fetus and Newborn, American College of Obstetricians and Gynecologists and Committee on Obstetric Practice. The apgar score. *Pediatrics* 2006;117(4):1444-7.
40. Sills JH. Perinatal asphyxia. Dalam: Gomella TL, Cunningham MD, Eyal FG, Zenk KE, penyunting. *Neonatology; Management. Procedures, On-Call Problems, Diseases and Drugs*. Edisi ke-5. New York: Lange Medical Book/McGraw-Hill Co, 2004. h.512-23.
41. American Heart Association and American Academy of Pediatrics. *Textbook of neonatal resuscitation*. Kattwinkel J, penyunting. Edisi ke-5. New York: McGraw-Hill, 2006.h.1-14.
42. Aurora S, Snyder EY. Perinatal asphyxia. Dalam : Cloherty J, Eichenwald EC, Stark AR, penyunting. *Manual of Neonatal Care*. Edisi ke-5. New York: Lippincott, William & Wilkins, 2004. h.536-55
43. Hill A, Volpe JJ. Hypoxic-ischemic cerebral injury in the newborn. Dalam: Swaiman KF, Ashwal S, penyunting. *Pediatric Neurology Principles and Practice*. Edisi ke-3. New York: Mosby Inc, 1999. h.191-204
44. Sävman K. The brain and resuscitation. *NeoReviews* 2008;9:513-9.

45. Perkin RM, Ashwal S. Hypoxic-ischemic encephalopathy in infants and older children. Dalam: Swaiman KF, Ashwal S, penyunting. *Pediatric Neurology Principles and Practice*. Edisi ke-3. New York: Mosby Inc, 1999. h.915-44
46. Johnston MV, Trescher WH, Ishida A, Nakajima W. Neurobiology of hypoxic-ischemic injury in the developing brain. *Pediatric Research* 2001;49(6):735-41.
47. Agius LM. Cytokine and trophic biology as cellular transcriptional susceptibility in neuronal ischemia. *International Journal of Molecular Medicine and Advance Sciences* 2005;1(2):121-9.
48. Ferriero DM. Neonatal brain injury. *N Engl J Med* 2004;35(1):1985-95.
49. Arpino C, D'Argenzo L, Ticconi C, Di Paolo A, Stellin V, Lopez V, et.al. Brain damage in preterm infants: etiological pathways. *Ann Ist Super Sanità* 2005;41(2):229-37.
50. St. Clair EG, Anderson SJ, Oltvai ZN. Bcl-2 counters apoptosis by bax heterodimerization-dependent and independent mechanisms in the T-cell lineage. *The Journal of Biological Chemistry* 1997; 272(46):29347–55.
51. Webster KA. Mitochondrial death channels. *American Scientist* 2009;97(5):73-88.
52. Maślińska D, Laure-Kamionowska M, Kaliszek A, Makarewicz D. Proinflammatory cytokines in injured rat brain following perinatal asphyxia. *Folia Neuropathol* 2002;40(4):177-82.
53. Hahn JS. Clinical manifestation of hypoxic-ischemic encephalopathy. Diunduh dari URL: <http://neuroresident.stanford.edu/Teaching/PediNeuro/HIE.clinical.chap.JH2001.html> tanggal 4 Januari 2009.
54. Wennberg RP, Ahlfors CE, Bhutani VK, Johnson LH, Shapiro SM. Toward understanding kernicterus: A challenge to improve the management of jaundiced newborns. *Pediatrics* 2006;117;474-85.
55. Amin SB, Ahlfors C, Orlando MS, Dalzell E, Merle KS, Guillet R. Bilirubin and serial auditory brainstem responses in premature infants. *Pediatrics* 2001;107(4):667-70.

56. Hille ETM, van Straaten HLM, Verkerk PH. Prevalence and independent risk factors for hearing loss in NICU infants. *Acta Pædiatrica* 2007; 96:1155-8.
57. Ghazali MV, Sastromihardjo S, Soedjarwo SR, Soelaryo T, Pramulyo H. Studi kohort. Dalam: Sastroasmoro S, Ismael S, penyunting. *Dasar-dasar Metodologi Penelitian Klinis*. Edisi ke-2. Jakarta: Sagung Seto, 2002. h.97-109.
58. Morley C, Davis P. Surfactant treatment for premature lung disorders: A review of best practices in 2002. *Paediatric Respiratory Reviews* 2004:299-304
59. Flint EF. Severe childhood deafness in Glasgow, 1965-1979. *J Laryngol Otol*. 1983;97(5):421-5.
60. Jiang ZD. Maturation of peripheral and brainstem auditory function in the first year following perinatal asphyxia: a longitudinal study. *J Speech Lang Hear Res* 1998;41(1):83-93.
61. Schmutzhard J, Glueckert R, Sergi C, Schwentner I, Abraham I, Schrott-Fischer A. Does perinatal asphyxia induce apoptosis in the inner ear?. *Hear Res* 2009;250(1-2):1-9
62. Fallon JB, Irvine DRF, Shepherd RK. Cochlear implants and brain plasticity. *Hear Res* 2008;238:110-7.
63. Departemen Kesehatan Republik Indonesia. *Pedoman pelaksanaan stimulasi, deteksi dan intervensi dini tumbuh kembang anak*. 2006.h.15-39.
64. Weichbold V, Nekahm-Heis D, Welzl-Mueller K. Universal newborn hearing screening and postnatal hearing loss. *Pediatrics* 2006;117(4):631-36.
65. Holster IR, Hoeve LJ, Wieringa MH, Willis-Lorrier RMS, Gier HHW. Evaluation of hearing loss after failed neonatal hearing screening. *J Pediatr* 2009;155:646-50.
66. du Plessis AJ. The role of systemic hemodynamic disturbances in prematurity-related. *J Child Neurol* 2009;24:1127-40.
67. Jiang ZD, Brosi DM, Wilkinson AR. Differences in impaired brainstem conduction between neonatal chronic lung disease and perinatal asphyxia. *Clin Neurophysiol* 2009;121(5):725-33.

68. Ohl C, Dornier L, Czajka C, Chobaut JC, Tavernier L. Newborn hearing screening on infants at risk. *Int J Pediatr Otorhinolaryngol* 2009;73(12):1691-5.
69. Bashirudin J, Soetirto I. Gangguan pendengaran akibat bising. Dalam : Buku Ajar Ilmu Kesehatan Telinga Hidung Tenggorok Kepala Leher. Edisi ke-6. Jakarta: Balai Penerbit FK UI, 2007. h.49-52.
70. Berg AL, Chavez CT, Serpanos YC. Monitoring noise levels in a tertiary neonatal intensive care unit. *CICSD* 2010; 37:69–72.
71. Kolegium Ilmu Kesehatan Telinga Hidung Tenggorok, Bedah Kepala dan Leher. Modul THT komunitas, preventif dan promotif. 2008.h.9-10.

---