

# *Helicobacter pylori* infection appears essential for stomach carcinogenesis: Observations in Semarang, Indonesia

Shinkan Tokudome,<sup>1,6</sup> Witjitra D. Samsuria,<sup>2</sup> Soeripto,<sup>3</sup> F. X. Ediati Triningsih,<sup>3</sup> Sadao Suzuki,<sup>1</sup> Akihiro Hosono,<sup>1</sup> Teguh Triono,<sup>3</sup> Indra Wijaya,<sup>4</sup> Sarjadi,<sup>4</sup> Ika P. Miranti,<sup>4</sup> Reza Ghadimi<sup>1</sup> and Malcolm A. Moore<sup>5</sup>

<sup>1</sup>Nagoya City University Graduate School of Medical Sciences, Mizuho-ku, Nagoya 467-8601, Japan; <sup>2</sup>Tugurejo Hospital, Semarang, Indonesia; <sup>3</sup>Faculty of Medicine, Gadjah Mada University, Yogyakarta, Indonesia; <sup>4</sup> Faculty of Medicine, Diponegoro University, Semarang, Indonesia; and <sup>5</sup>APJCP Editorial Office, Asian Pacific Organization for Cancer Prevention, c/o National Cancer Institute, Bangkok, Thailand

(Received July 7, 2005/Revised September 26, 2005/Accepted September 30, 2005/Online publication December 8, 2005)

The gastric cancer incidence in Semarang, Indonesia, is exceedingly low: only approximately 1/100th of the level in Japan. To elucidate the reason, we carried out an ecological study recruiting 69 male and 102 female participants from the general populace in January 2005. Positive urea breath tests were 0% for both men and women, and *Helicobacter pylori* (*H. pylori*) IgG antibodies were found in 2% (0–5, 95% confidence interval) of men and 2% (0–4) of women, significantly lower than the 62% (58–65) and 57% (53–60), respectively, in Japan. Furthermore, there were no positive findings with the pepsinogen tests in Semarang, again significant in comparison with the 23% (22–25) and 22% (20–23) in Japan. Variation in smoking levels and consumption of NaCl, vegetables and fruit were found, but not to an extent that would allow explanation of the major differences in gastric cancer incidence. We may conclude that the very low prevalence of *H. pylori* infection and thus chronic atrophic gastritis account for the rarity of stomach cancer in Semarang, Indonesia. (*Cancer Sci* 2005; 96: 873–875)

Since 2002, we have been conducting a collaborative epidemiologic appraisal of host and environmental factors for stomach and colorectal cancer in several South-east Asian countries. Ecological and case-control studies are now being carried out in Hanoi and Ho Chi Minh City, Vietnam; Khon Kaen, Thailand; and Yogyakarta and Semarang, Indonesia, in order to take advantage of the major variation in cancer incidence among these geographical areas and also with data for Japan. Stomach cancer incidence rates in Hanoi, Ho Chi Minh City, Khon Kaen, Yogyakarta and Semarang are approximately 1/2, 1/4, 1/10, 1/50 and 1/100 those prevailing in Japan, respectively: the annual age-adjusted incidence rates for Semarang were 0.6/10<sup>5</sup> for men and 0.3/10<sup>5</sup> for women during 1990–1999, and the respective figures for Japan in 1995 were 67/10<sup>5</sup> and 27/10<sup>5</sup>.<sup>(1,2)</sup>

Gastric cancer may be caused by environmental or lifestyle risks, host genetic polymorphisms, as well as aging.<sup>(3)</sup> Many laboratory studies have pointed to roles for carcinogenic substances, including amine pyrolysate products and nitrosamines; however, grilled or barbecued meat and fish are categorized as possible risk factors in humans.<sup>(3)</sup> A probable risk factor for stomach cancer is salt or salty foods, which act

synergistically with *Helicobacter pylori* (*H. Pylori*) infection in the development of stomach cancer in experimental animals<sup>(4)</sup> and humans.<sup>(5)</sup> Convincing preventive factors are vegetables and fruit and refrigeration, and a probable preventive factor is vitamin C. The International Agency for Research on Cancer (IARC) has concluded that gastric cancer is a smoke-related malignancy.<sup>(6)</sup> *H. pylori* is a definite carcinogen,<sup>(7–9)</sup> and is accepted to be a major factor for chronic atrophic gastritis (CAG),<sup>(5)</sup> a precursor lesion for stomach cancer.<sup>(10)</sup>

We here report the results of an ecological study of stomach cancer with reference to *H. pylori* infection and pepsinogen tests as a marker of CAG,<sup>(11,12)</sup> along with smoking habits and excretion of sodium and potassium<sup>(13)</sup> as markers of intake of salt or salted foods, and vegetables and fruit, respectively.<sup>(14)</sup> The study took place in Semarang, located in the central-north of Java island facing the Java sea, and was compared with published values for Japanese people.<sup>(15)</sup>

## Subjects and Methods

In January 2005, we randomly recruited 69 male and 102 female participants from the general populace, mostly Javanese, in the city of Semarang. Mean ages were 57.4 ± 10.9 (SD) for men and 49.2 ± 9.8 for women. Written informed consent was obtained from the study participants and the protocol was approved by the Internal Review Boards of Nagoya City University, Japan, and Diponegoro University and Tugurejo Hospital, Semarang, Indonesia. The subjects were requested to respond to lifestyle questions, including smoking habits, and food frequency questionnaires, and were interviewed by health nurses at a local hospital. Bodyweight and height were measured, and oral mucous membranes, overnight-fasting blood, breath, second morning voiding urine and feces were sampled from each participant.

For the urea breath test (UBT), UBiT-IR300 kits (Otsuka Pharmaceutical, Tokyo, Japan) were used with ≥ 2.5‰ as positive. Serum antibodies for *H. pylori* were examined by enzyme immunoassay (Kyowa Medics, Tokyo, Japan) and

<sup>6</sup>To whom correspondence should be addressed.  
E-mail: tokudome@med.nagoya-cu.ac.jp

**Table 1. *Helicobacter pylori*-related markers and pepsinogen (PG) tests in the populace in Semarang compared with Japan**

Test	Semarang		Japan	
	Men (n = 69)	Women (n = 102)	Men	Women
Positive urea breath test (%) <sup>†</sup>	0	0	NA	NA
Positive serum <i>H. pylori</i> IgG (%) <sup>‡</sup>	2 (0–5) <sup>§</sup>	2 (0–4)	62 (58–65) <sup>¶</sup>	57 (53–60)
Positive pepsinogen test (%) <sup>††</sup>	0	0	23 (22–25) <sup>**</sup>	22 (20–23)

<sup>†</sup>Positive test was defined as  $\geq 2.5\%$  by UBIT-IR300 kits. <sup>‡</sup>Positive test was defined as ELISA value  $\geq 2.3$  by enzyme immunoassay. <sup>§</sup>Age-adjusted prevalence (95% confidence interval). <sup>¶</sup>Values are cited from reference number (17). <sup>\*\*</sup>Positive test was defined as PGI  $\leq 70$  ng/mL by chemical luminescence enzyme immunoassay and PGI/PGII  $\leq 3.0$ . <sup>\*\*</sup>Values are cited from reference number (12).

**Table 2. Smoking rates and urinary excretion of salt and potassium in the populace in Semarang compared with Japan**

	Semarang		Japan	
	Men	Women	Men	Women
Smoking rate (%)	55.0 (22.3–87.8) <sup>†</sup>	0	49.5 (47.2–51.7) <sup>†</sup>	18.1 (16.8–19.5)
Salt (g/day)	11.1 (9.8–12.5)	10.4 (9.9–11.0)	12.9 (12.7–13.1) <sup>§</sup>	11.2 (11.1–11.4)
Potassium (g/day)	2.3 (2.2–2.5)	2.1 (2.1–2.2)	2.5 (2.5–2.5) <sup>§</sup>	2.4 (2.3–2.4)

<sup>†</sup>Age-adjusted mean (95% confidence interval). <sup>‡</sup>Values were cited from reference number (18). <sup>§</sup>Values were cited from reference number (15).

ELISA values  $\geq 2.3$  were defined as positive. Serum pepsinogen (PG) I and PGII were measured by chemical luminescence enzyme immunoassay (Eiken Chemicals, Tokyo, Japan) with cut off points of PGI  $\leq 70$  ng/mL and PGI/PGII  $\leq 3.0$ ,<sup>(11)</sup> as a non-invasive surrogate for the Sydney classification of CAG.<sup>(16)</sup> For all of these parameters, age-adjustment was made adopting the world population,<sup>(1)</sup> as standard, for comparison with Japanese figures.

Second morning voiding urine excretions of sodium, as a marker of intake of salt or salty foods, and potassium, as a marker of consumption of vegetables and fruit, were analyzed by electrode assays, and creatinine was analyzed by an enzymatic method. The daily levels were then estimated with adjustment for creatinine,<sup>(13)</sup> and compared with the consumption values of Japanese people.<sup>(15)</sup>

## Results

As shown in Table 1, age-adjusted positive rates for UBT were 0% for both men and women in Semarang. *H. pylori* IgG antibodies were found in only 2% (0–5) of men and 2% (0–4) of women, and were significantly lower than the 62% (58–65) and 57% (53–60), respectively, in Japan.<sup>(17)</sup> Positive findings for the PG tests were 0% in both sexes, again significantly lower than the 23% (22–25) and 22% (20–23) reported for Japanese people.<sup>(12)</sup>

Smoking rates were rather higher for men in Semarang than in Japan but lower for women (Table 2).<sup>(18)</sup> Salt excretions were calculated to be 11.1 g/day (9.8–12.5 g/day) for men and 10.4 g/day (9.9–11.0 g/day) for women in Semarang, and only slightly lower than the consumption of 12.9 g/day (12.7–13.1 g/day) for men and 11.2 g/day (11.1–11.4 g/day) for women in Japan.<sup>(15)</sup> Values for potassium excretions were 2.3 g/day (2.2–2.5 g/day) for men and 2.1 g/day (2.1–2.2 g/day) for women, again close to the consumption of 2.5 g/day (2.5–2.5 g/day) and 2.4 g/day (2.3–2.4 g/day) in Japan.

## Discussion

Our findings of negative or very low prevalence of UBT and *H. pylori* IgG antibodies in both men and women in Semarang, Indonesia are in clear contrast to the case in Japan and, presumably, are directly related to the null PG testing. Smoking was found to be more common for men in Semarang than in Japan, but less common for women. There were only minor differences in estimated intake values for salt, and for vegetables and fruit between people in Semarang and Japan. No evidence was obtained that the marked variation in cancer incidence could be explained on the basis of low consumption of salt or salty foods, or far greater vegetable and fruit consumption. We can assume that refrigeration is less prevalent in Semarang than in Japan. Whether host genetic polymorphisms associated with cellular immunity for chronic inflammation,<sup>(19)</sup> or differences in *H. pylori* strains regarding infectivity, virulence or inflammatory potency,<sup>(5,20)</sup> may be important factors in Semarang remains to be clarified. However, the most plausible interpretation of our results is that the rarity of gastric cancer in Semarang is attributable to a low prevalence of *H. pylori* infection along with the non-existence of CAG, the well-established precursor for stomach cancer.<sup>(5,7–10)</sup>

The precise reason why *H. pylori* prevalence is very low in Javanese people in Semarang and Yogyakarta is unknown.<sup>(21)</sup> It seems paradoxical given the fact that *H. pylori* is transmitted via fecal–oral, oral–oral or water-borne routes, because most people in Semarang still use well water and the sanitary environment conditions remain basic.<sup>(5)</sup> The low positivity in Javanese people is, however, compatible with the low prevalence reported for Malay people in Malaysia who share ethnic traits.<sup>(22)</sup> We should scrutinize whether genetic polymorphisms of Javanese people might confer immunity against *H. pylori* infection. It is also conceivable that specific environmental factors on Java island, including dietary habits, may be effective in preventing *H. pylori* infection.

While ecological studies are generally regarded as providing low-rank evidence, and the number of recruited subjects was not, strictly speaking, randomly selected and not sufficiently large to be representative of the Semarang populace, our data do provide simple and compelling evidence that the low stomach cancer incidence in Semarang is due to the very low prevalence of *H. pylori* infection and CAG, suggesting that *H. pylori* is a definite and essential factor for the onset of stomach cancer. Further, large scale studies taking into account local medical practice, such as gastric cancer screen-

ing and cancer registration, including its completeness and accuracy, now need to be carried out for confirmation.

## Acknowledgments

Supported in part by a Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Culture, Sports, Science and Technology, Japan. We thank Dr M. Ikeda, Ms N. Nakanishi and Ms M. Watanabe for their statistical and technical assistance.

## References

- IARC/IACR. *Cancer Incidence in Five Continents*, Vol. VIII. Lyon: IARC, 2002.
- Oshima A, Tsukuma H, Ajiki W. Cancer registration in Japan. *Asian Pac J Cancer Prev* 2001; **2** (IACR Suppl.): 31–5.
- World Cancer Research Fund/American Institute for Cancer Research. *Food, Nutrition and the Prevention of Cancer: a Global Perspective*. Washington, DC: World Cancer Research Fund/American Institute for Cancer Research, 1997.
- Nozaki K, Shimizu N, Inada K *et al*. Synergistic promoting effects of *Helicobacter pylori* infection and high-salt diet on gastric carcinogenesis in Mongolian gerbils. *Jpn J Cancer Res* 2002; **93**: 1083–9.
- Graham KS, Graham DY. *H. pylori-Associated Gastrointestinal Diseases*, 2nd edn. Newtown: Handbooks in Health Care, 2002.
- IARC. *Monographs on the Evaluation of Carcinogenic Risks to Humans. Tobacco Smoke and Involuntary Smoking*, Vol. 83. Lyon: IARC, 2004.
- IARC. *Monographs on the Evaluation of Carcinogenic Risks to Humans. Schistosomes, Liver Flukes and Helicobacter pylori*, Vol. 61. Lyon: IARC, 1994.
- Uemura N, Okamoto S, Yamamoto S *et al*. *Helicobacter pylori* infection and the development of gastric cancer. *N Engl J Med* 2001; **345**: 784–9.
- Brenner H, Arndt V, Stegmalter C, Ziegler H, Rothenbacher D. Is *Helicobacter pylori* infection a necessary condition for noncardia gastric cancer? *Am J Epidemiol* 2004; **159**: 252–8.
- Correa P. A human model of gastric carcinogenesis. *Cancer Res* 1988; **48**: 3554–60.
- Miki K, Morita M, Sasajima M, Hoshina R, Kanda E, Urita Y. Usefulness of gastric cancer screening using the serum pepsinogen test method. *Am J Gastroenterol* 2003; **98**: 735–9.
- Watase H, Inagaki T, Yoshikawa I, Furihata S, Watanabe Y, Miki K. Five years follow up study of gastric cancer screening using the pepsinogen test method in Adachi city. *J Jpn Assoc Cancer Detect Diag* 2004; **11**: 77–81.
- Kawasaki T, Itoh K, Uezono K, Sasaki H. A simple method for estimating 24h urinary sodium and potassium excretion from second morning voiding urine specimen in adults. *Clin Exp Pharmacol Physiol* 1993; **20**: 7–14.
- Science and Technology Agency, Japan. *Follow-up of Standard Tables of Food Composition in Japan*. Tokyo: Ishiyaku Shuppan, 1992. (In Japanese.)
- Ministry of Health, Labour and Welfare, Japan. *The National Nutrition Survey in Japan, 2002*. Tokyo: Daiichi Shuppan, 2004. (In Japanese.)
- Dixon MF, Genta RM, Yardley JH, Correa P. Classification and grading of gastritis. The updated Sydney System. International Workshop on the Histopathology of Gastritis, Houston, 1994. *Am J Surg Pathol* 1996; **20**: 1161–81.
- Kikuchi S, Nakajima T, Kobayashi O *et al*. Effect of age on the relationship between gastric cancer and *Helicobacter pylori*. *Jpn J Cancer Res* 2000; **91**: 774–9.
- Japan Tobacco [website on the internet]. [Cited 15 August 2005.] Available from URL: <http://www.health-net.or.jp/tobacco/product/pd090000.html>
- El-Omar EM, Carrington M, Chow WH *et al*. Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 2000; **404**: 398–402.
- Covacci A, Telford JL, Del Giudice G, Parsonnet J, Rappuoli R. *Helicobacter pylori* virulence and genetic geography. *Science* 1999; **284**: 1328–33.
- Tokudome S, Soeripto, Triningsih FXE *et al*. Rare *Helicobacter pylori* infection as a factor for the very low stomach cancer incidence in Yogyakarta, Indonesia. *Cancer Lett* 2005; **219**: 57–61.
- Miwa H, Go MF, Sato N. *H. pylori* and gastric cancer: the Asian enigma. *Am J Gastroenterol* 2002; **97**: 1106–12.