

DAFTAR PUSTAKA

1. Passie, Torsten., Seifert, Juergen., Schneider, Udo and M. Emrich, Hinderk. The Pharmacology of Psilocybin. [homepage on internet]. Available from: <http://files.shroomery.org/cms/4529203-pharmacologyofpsilocybin.pdf> (accessed 2002)
2. Setyabudi, Nugroho. Mabuk Jamur Mushroom, Mahasiswa Undip Ngamuk & Tewas. [homepage on internet]. Available from: <http://jogja.okezone.com/read/2012/11/17/512/719612/mabuk-jamur-mushroom-mahasiswa-undip-ngamuk-tewas> (accessed 2012)
3. Supradja, IGNA. Hasil wawancara/survei di Bali (Kuta, Sanur dan sekitarnya) tentang Penggunaan Jamur Panaeolus ini (“ Magic Mushroom”) dalam hidangan “Special Ommelet”, 1985
4. Walker, Jim I. Different Psiloscybin Species. [homepage on internet]. Available from: http://www.erowid.org/plants/mushrooms/mushrooms_info8.shtml (accessed 1994)
5. Gartz, Jochen. Extraction and Analysis of Indole Derivatives From Fungal Biomass. [homepage on internet]. Available from: http://www.erowid.org/plants/mushrooms/mushrooms_journal1.shtml (accessed 1994)
6. Hasler F, Grimberg U, Benz MA, Huber T, Vollenweider FX. "Acute psychological and physiological effects of psilocybin in healthy humans: a

- double-blind, placebo-controlled dose-effect
study". *Psychopharmacology* 172 (2); 2004. p.145–56
7. Erowid. Psilocybin, Psilocin, and Magic Mushrooms Dose [homepage on internet]. Available from:
http://www.erowid.org/plants/mushrooms/mushrooms_dose.shtml
8. Gartz J. "New aspects of the occurrence, chemistry and cultivation of European hallucinogenic mushrooms". *Supplemento agli Annali dei Musei Civici di Rovereto Sezione Archeologica, Storia e Scienze Naturali* 8; 1992. p.107–24
9. Gartz J. Narrenschwaemme. Psychotrope Pilze di Europa di Europa. Herausforderung sebuah Wertsystem und Forschung. Editions Heuwinkel. Genf / Neuallschwill, 1993
10. Beug MW, Bigwood J. Psilosibin and psilosin level in twenty generations multiple species of wild mushrooms in the Pacific Northwest. USAJ Ethnopharm 5; 1982. p.271-289
11. Gartz J. Biotransformasi derivatif tryptamine dalam budaya miselium Psilocybe. J. Dasar Microbiol 29; 1989. p.347-352
12. Gartz J. Biotransformasi tryptamine dalam miselia berbuah cubensis Psilocybe. Planta Med 55; 1989. p.249-250
13. Sottolano SM, Lurie IS. Quantitation of psilocybin in hallucinogenic mushrooms using high performance liquid chromatography. J. Forensik Sci 28; 1983. p.931-935

14. Casale JF. An organic-water extraction method for the isolation and identification of psilocin from hallucinogenic mushrooms. *J. Forensik Sci* 30; 1985. p.247-250
15. Kysilka R, Wurst M. A new extraction procedure for psilocybin and psilocin determination in mushroom samples. *Planta Med* 56; 1990. p.327-328
16. Wurst M, Kysilka R, Koza T. Analysis and indole alkaloid isolation mushroom by high performance liquid chromatography. *J. Chromatogr* 593; 1992. p.201-208
17. Hasler F, Bourquin D, Brenneisen R, Vollenweider FX. Renal excretion profiles of psilocin following oral administration of psilocybin: a controlled study in man. *J Pharm Biomed Anal* 30; 2002. p.331-9
18. Carter OL, Pettigrew JD, Burr DC, Alais D, Hasler F, Vollenweider FX. Psilocybin impairs high-level but not low-level motion perception. *Neuroreport* 15; 2004. p.1947-1951.
19. Vollenweider FX, Vontobel P, Hell D, Leenders KL. 5-HT modulation of dopamine release in basal ganglia in psilocybin-induced psychosis in man-a PET study with [11C]raclopride. *Neuropsychopharmacology* 20; 1999. p.424-33
20. Carter OL, Pettigrew JD, Burr DC, Alais D, Hasler F, Vollenweider FX. Psilocybin impairs high-level but not low-level motion perception. *Neuroreport* 15; 2004. p.1947-51

21. Carter OL, Pettigrew JD, Hasler F, Wallis GM, Liu GB, Hell D, Vollenweider FX. Modulating the rate and rhythmicity of perceptual rivalry alternations with the mixed 5-HT2A and 5-HT1A agonist psilocybin, 2005
22. Erowid. Psilocybin Mushrooms. [homepage on internet]. Available from: http://www.erowid.org/plants/mushrooms/mushrooms_effects.shtml (accessed 1997)
23. Gouzoulis-Mayfrank E, Schreckenberger M, Sabri O, Arning C, Thelen B, Spitzer M, Kovar KA, Hermle L, Bull U, Sass H. Neurometabolic effects of psilocybin, 3,4-methylenedioxymethamphetamine (MDMA) and d-methamphetamine in healthy volunteers. A double-blind, placebo-controlled PET study with [18F]FDG. *Neuropsychopharmacology* 20; 1999. p.565-81
24. Vollenweider FX, Leenders KL, Scharfetter C, Maguire P, Stadelmann O, Angst J. Positron emission tomography and fluorodeoxyglucose studies of metabolic hyperfrontality and psychopathology in the psilocybin model of psychosis. *Neuropsychopharmacology* 16; 1997. p.357-72
25. Vollenweider FX, Vollenweider-Scherpenhuyzen MF, Babler A, Vogel H, Hell D. "Psilocybin induces schizophrenia-like psychosis in humans via a serotonin-2 agonist action". *NeuroReport* 9 (17); 1998. p.3897–902
26. Carhart-Harris RL, Nutt DJ. "User perceptions of the benefits and harms of hallucinogenic drug use: a web-based questionnaire study". *Journal of Substance Abuse* 15 (4); 2010. p.283–300

27. Van Amsterdam J, Opperhuizen A, van den Brink W. "Harm potential of magic mushroom use: a review". *Regulatory Toxicology and Pharmacology* 59 (3); 2011. p.423–9.
28. Simeon D. "Depersonalisation disorder: a contemporary overview". *CNS Drugs* 18 (6); 2004. p.343–54
29. Peden NR, Pringle SD. Hallucinogenic fungi. *Lancet* 1; 1982. p.396-7
30. Strassman RJ. Adverse reactions to psychedelic drugs. A review of the literature. *J Nerv Ment Dis* 172; 1984. p.577-95
31. Nichols CD, Sanders-Bush E. Molecular genetic responses to lysergic acid diethylamide include transcriptional activation of MAP kinase phosphatase-1, C/EBP-beta and ILAD-1, a novel gene with homology to arrestins. *J Neurochem* 90; 2004. p.576-84
32. Strassman RJ, Qualls CR, Uhlenhuth EH, Kellner R. Dose-response study of N,Ndimethyltryptamine in humans. II. Subjective effects and preliminary results of a new rating scale. *Arch Gen Psychiatry* 51; 1994. P.98-108
33. World health organization. General guidelines for methodologies on research and evaluation of traditional medicine [homepage on internet]. Geneva: WHO; 2001
34. Reigner B, Blesch K. Estimating the starting dose for entry into humans: principles and practice. *European Journal of Clinical Pharmacology* 57(12); 2002. p.835-845

Lampiran 1. Ethical clearance

	<p>KOMISI ETIK PENELITIAN KESEHATAN (KEPK) FAKULTAS KEDOKTERAN UNIVERSITAS DIPONEGORO DAN RSUP dr KARIADI SEMARANG Sekretariat : Kantor Dekanat FK Undip Lt.3 Jl. Dr. Soetomo 18. Semarang Telp. 024-8311523/Fax. 024-8446905</p>	 RSUP Dr. KARIADI
<p><u>ETHICAL CLEARANCE</u> No.223 /EC/FK/RSDK/2013</p>		
<p>Komisi Etik Penelitian Kesehatan Fakultas Kedokteran Universitas Diponegoro/ RSUP. dr. Kariadi Semarang, setelah membaca dan menelaah USULAN Penelitian dengan judul :</p>		
<p style="text-align: center;">PENGARUH PEMBERIAN EKSTRAK JAMUR PSILOCYBE CUBENSIS DOSIS BERTINGKAT TERHADAP AKTIVITAS MOTORIK MENCIT SWISS WEBSTER DENGAN METODE ROTAROD MANUAL</p>		
<p>Peneliti Utama : Orieza Sativa Novitaloka</p>		
<p>Pembimbing : 1. dr. Sigid Kirana Lintang Bima, Sp.F 2. dr. Tuntas Dhanardhono, M.Si.Med</p>		
<p>Penelitian : Dilaksanakan di Laboratorium Forensik FK Undip Cabang Semarang untuk pengandangan hewan dan uji pada hewan coba</p>		
<p>Setuju untuk dilaksanakan, dengan memperhatikan prinsip-prinsip yang dinyatakan dalam Deklarasi Helsinki 1975, yang diamended di Seoul 2008 dan Pedoman Nasional Etik Penelitian Kesehatan (PNEPK) Departemen Kesehatan RI 2011</p>		
<p>Pada laporan akhir peneliti harus melampirkan cara pemeliharaan & dekapitasi hewan coba</p>		
<p>Semarang, 24 Mei 2013</p>		
 <p>KOMISI ETIK PENELITIAN KESEHATAN FAKULTAS KEDOKTERAN UNDIP/RSUP DR. KARIADI KETUA, Prof.Dr.dr.Suprihati, M.Sc, Sp.THT-KL(K) NIP. 19500621197703 2 001</p>		

Lampiran 2. Cara perhitungan dosis

1. Besar dosis jamur *Psilocybe cubensis* dosis ringan pada manusia 0,25 gram - 1 gram, dosis sedang dosisnya adalah 1 gram - 2,5 gram dan dosis berat adalah 2,5 gram - 5 gram.
2. Rumus yang biasa di gunakan adalah Human Equivalent Doses (HED dalam mg / kg) = Dosis Hewan (mg / kg) × Hewan Km ÷ Km Manusia.
3. Maka, di dapatkan dosis hewan dengan besar KM pada mencit adalah 3, yaitu:

- 1) Dosis ringan:

$$\text{PAD: } \frac{0,06 \times 37}{3} = 0,74 \sim 0,75 \text{ g/kg}$$

- 2) Dosis sedang:

$$\text{PAD: } \frac{0,12 \times 37}{3} = 1,48 \sim 1,5 \text{ g/kg}$$

- 3) Dosis berat:

$$\text{PAD: } \frac{0,24 \times 37}{3} = 2,96 \sim 3 \text{ g/kg}$$

Lampiran 3. Hasil analisis pengamatan aktivitas motorik mensit swiss webster dengan metode rotarod manual

Explore

kelompok mencit

Case Processing Summary

	kelompok mencit	Cases					
		Valid		Missing		Total	
		N	Percent	N	Percent	N	Percent
skor aktivitas motorik	kelompok kontrol	5	100,0%	0	,0%	5	100,0%
	kelompok perlakuan dosis rendah 0,75gr/kg	5	100,0%	0	,0%	5	100,0%
	kelompok perlakuan dosis sedang 1,5gr/kg	5	100,0%	0	,0%	5	100,0%
	kelompok perlakuan dosis tinggi 3gr/kg	5	100,0%	0	,0%	5	100,0%

Descriptives

		kelompok mencit		Statistic	Std. Error
skor aktivitas motorik	kelompok kontrol	Mean		444,00	3,114
		95% Confidence Interval for Mean	Lower Bound	435,35	
			Upper Bound	452,65	
		5% Trimmed Mean		443,83	
		Median		442,00	
		Variance		48,500	
		Std. Deviation		6,964	
		Minimum		437	
		Maximum		454	
		Range		17	
		Interquartile Range		13	
		Skewness		,725	,913
		Kurtosis		-,933	2,000
kelompok perlakuan dosis rendah 0,75gr/kg		Mean		349,20	5,398
		95% Confidence Interval for Mean	Lower Bound	334,21	
			Upper Bound	364,19	
		5% Trimmed Mean		349,50	
		Median		353,00	
		Variance		145,700	
		Std. Deviation		12,071	
		Minimum		331	
		Maximum		362	
		Range		31	
		Interquartile Range		22	
		Skewness		-,877	,913
		Kurtosis		,223	2,000
kelompok perlakuan dosis sedang 1,5gr/kg		Mean		305,00	7,855
		95% Confidence Interval for Mean	Lower Bound	283,19	
			Upper Bound	326,81	
		5% Trimmed Mean		304,94	
		Median		305,00	
		Variance		308,500	
		Std. Deviation		17,564	
		Minimum		283	
		Maximum		328	
		Range		45	
		Interquartile Range		33	
		Skewness		,091	,913
		Kurtosis		-,924	2,000
kelompok perlakuan dosis tinggi 3gr/kg		Mean		231,40	4,523
		95% Confidence Interval for Mean	Lower Bound	218,84	
			Upper Bound	243,96	
		5% Trimmed Mean		231,61	
		Median		233,00	
		Variance		102,300	
		Std. Deviation		10,114	
		Minimum		217	
		Maximum		242	
		Range		25	
		Interquartile Range		19	
		Skewness		-,608	,913
		Kurtosis		-,855	2,000


Tests of Normality

	kelompok mencit	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
skor aktivitas motorik	kelompok kontrol	,213	5	,200*	,935	5	,634
	kelompok perlakuan dosis rendah 0,75gr/kg	,224	5	,200*	,948	5	,720
	kelompok perlakuan dosis sedang 1,5gr/kg	,134	5	,200*	,991	5	,984
	kelompok perlakuan dosis tinggi 3gr/kg	,174	5	,200*	,954	5	,763

*. This is a lower bound of the true significance.

a. Lilliefors Significance Correction

skor aktivitas motorik



T-Test

Group Statistics

		N	Mean	Std. Deviation	Std. Error Mean
skor aktivitas motorik	kelompok mencit kelompok kontrol kelompok perlakuan dosis rendah 0,75gr/kg	5 5	444,00 349,20	6,964 12,071	3,114 5,398

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						95% Confidence Interval of the Difference	
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	Lower		Upper
						94,800	6,232	80,429	109,171	94,800
skor aktivitas motorik	1,554 Equal variances assumed 15,211 Equal variances not assumed 15,211	,248 6,397	15,211 6,397	8 94,800	,000 ,000	94,800 94,800	6,232 6,232	79,777 79,777	109,823 109,823	

T-Test

Group Statistics

		N	Mean	Std. Deviation	Std. Error Mean
skor aktivitas motorik	kelompok mencit kelompok kontrol kelompok perlakuan dosis sedang 1,5gr/kg	5 5	444,00 305,00	6,964 17,564	3,114 7,855

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						95% Confidence Interval of the Difference	
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	Lower		Upper
						139,000	8,450	119,515	158,485	139,000
skor aktivitas motorik	2,888 Equal variances assumed 16,450 Equal variances not assumed 16,450	,128 5,227	16,450 5,227	8 139,000	,000 ,000	8,450 8,450	117,560 117,560	160,440 160,440		


T-Test

Group Statistics

		N	Mean	Std. Deviation	Std. Error Mean
	kelompok mencit	5	444,00	6,964	3,114
skor aktivitas motorik	kelompok kontrol kelompok perlakuan dosis tinggi 3gr/kg	5	231,40	10,114	4,523

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
						Lower		Upper	
skor aktivitas motorik	,811	,394	38,712	8	,000	212,600	5,492	199,936	225,264
			38,712	7,097	,000	212,600	5,492	199,650	225,550

Lampiran 4. Dokumentasi penelitian

Gambar 7. Proses penelitian

- a. Rotarod manual yang digunakan
- b. Pengandangan dan proses adaptasi mencit selama 7 hari
- c. Serbuk jamur *Psilocybe cubensis*
- d. Proses pembuatan infusa

Lampiran 5. Biodata mahasiswa**Identitas**

Nama : Orieza Sativa Novitaloka
NIM : G2A009192
Tempat/tanggal lahir : Semarang/14 November 1992
Jenis kelamin : Perempuan
Alamat : Perum Rumpun Diponegoro Jl. Bangau IV/54 Semarang
Nomor HP : 085641401992
e-mail : orieza.sativa@yahoo.com

Riwayat Pendidikan Formal

1. SD : SD Negeri Jomblang 02 Semarang Lulus tahun: 2004
2. SMP : SMP Negeri 9 Semarang Lulus tahun: 2007
3. SMA : SMA Negeri 1 Semarang Lulus tahun: 2009
4. S1 : Fakultas Kedokteran Universitas Diponegoro Masuk tahun:2009