

## DAFTAR PUSTAKA

1. M. Sholeh kosim , dkk. Buku Ajar Neonatologi. Ikatan Dokter Anak Indonesia. Jakarta . 2012
2. Blackburn ST, penyunting. Bilirubin metabolisme. Maternal, fetal, & neonatal fisiologi, a clinical perspective. Edisi ke-3. Saunders. Missouri; 2007
3. Wong RJ, Stevenson DK, Ahlfors CE, Vreman HJ. Neonatal Jaundice : Bilirubin Physiology and Clinical Chemistry. *NeoReviews* 2007; 8; 58-67.
4. M. Sholeh Kosim, dkk. Hubungan Hiperbilirubinemia dan Kematian Pasien yang Dirawat di NICU RSUP Dr Kariadi Semarang. *Sari Pediatri* [Internet]. 2007
5. Indarso F. Tranfusi tukar pada neonatus dengan hiperbilirubinemia. Dalam: Makalah lengkap kongres nasional VII perinasia & simposium internasional, 2003. H84-98
6. Etika R, Harianto A, Indarso F, Damanik SM. Hiperbilirubinemia pada neonatus. Dalam: Permono B, Kaspan F, Soegianto S, Soejoso DA, Narendra MB, Noer MS, penyunting. *Continuing education ilmu kesehatan anak*, 2004
7. Lissauer Tom dan Fanaroff Avroy. *At a Glance Neonatology*. Penerbit Erlangga. Jakarta. 2009
8. Porter ML, Dennis BL, Hyperbilirubinemia in the term newborn. *Am Fam Phy* 2002; 65;599-606
9. Sarici SU, Serdar MA, Korkmaz A, Erdem G, Oran O, Tekinalp G, dkk. Incident, course, and prediction of hyperbilirubinemia in near term and term newborn. *Pediatrics* 2004: 113:775-80
10. Mac Mahon JR, Stevenson DK, Oski FA. Bilirubin metabolism. Dalam : Taeusch HW, Ballard RA, editor. *Avery's disease of the newborn*. Edisi ke-7 .Philadelphia : WB Saunders Company, 1998;h-995-1002.

11. Maisels MJ. Jaundice. Dalam : Avery GB, Flecher MA, Mac Donald MG , Penyunting. Neonatology & management of the newborn. Edisi ke 5. Baltimore: Lippincot William & Wilkins , 1999; h-765-819
12. Halamek LP, Stevenson DK. Neonatal jaundice and liver disease, Dalam ; Fanaroff AA, Martin RJ, penyunting. Neonatal –perinatal madicine. Disease of the fetus in infant. Edisi ke 7. St louis; Mosby inc, 2002 ; h.1309-50
13. S. A. K. Indriyani, I Wayan Retayasa, Achmad Surjono,Purnomo Suryantoro. Percentage birth weight loss and hyperbilirubinemia during the first week of life in term newborns. Paediatrica Indonesiana [Internet]. 2003.
14. Surjono A. Manajemen hiperbilirubinemia pada Bayi Baru Lahir . Preceeding of the Oda-Day Seminar of Management of Icterus in neonatus;2003 May 24; Yogyakarta, Indonesia. Yogyakarta: FKUGM, 2003
15. Meyer Thomas C. A Study Of Serum Bilirubin Levels In Relation to Kericterus and Prematurity. ... 1955
16. Newman TB, Liljestrang P, Jeremy RJ, Ferriero DM, Wu YW, Hudes ES, et al, for the Jaundice and Infant Feeding Study Team. Outcomes among newborns with total serum bilirubin levels of 25 mg per deciliter or more. N Engl J Med . 2006;354:1889–900
17. Stoll B, Kleigman R. Jaundice and hyperbilirubinemia in the Neewborn. In Behrman R, Kleigman R, Jenson H, eds. *Nelson Book of Pediatrics*. 17 ed. Philadelphia: W.B Saunders 2004 ; 592-8
18. Martin C, Cloherty J. Neonatal hyperbilirubinemia. In; Cloherty J, Eichenwald E, Stark A, eds. *Manual of Neonatal care*. 5 ed. USA ; lippincott wiliam & wilkins 2004:185-221
19. Stirrat G. The immune system.In Hytten F, Chamberlain G ed: *Clinical Physiologyin Obstetrics*. London. Blackwell; 1991.101
20. Sarwono Prawiroharjo. Ilmu kebidanan. Edisi keempat. PT. Bina Pustaka Sarwono Prawiroharjo.Jakarta. 2010

21. World Health of Organization Collaborative Study Team on the Role of Breastfeeding on the Prevention of infant mortality: Effect of breastfeeding on infant and child mortality due to infection disease in less development countries : A pooled analysis. *Lancet*. 2000; 355; 452
22. Linn S, Schoenbaum SC, Monson RR, Rosner B, Stubblefield PG, Ryan KJ: Epidemiology of neonatal hyperbilirubinemia. *Pediatrics* 1985;75:770-4.
23. University of Maryland. Amniotic fluid. Available from ; URL; <http://www.umm.edu/ency/article/002220.htm>
24. Pillitteri, Adele. Mathernal and Child health Care Nurshing. Edisi ke 6. Lippincott Williams & Wilkins. Philadelphia.2010
25. March of Dimes. Amniotic fluid Abnormalities. Available from : URL ; [http://www.marchofdimes.com/professionals/14332\\_4536.asp](http://www.marchofdimes.com/professionals/14332_4536.asp)
26. Williams W. Amniotic Fluid : Objectives with narration and illustratrations. Available from: URL : [http://www.clt.astate.edu/wwilliam/cls\\_1521.urinalysis.htm](http://www.clt.astate.edu/wwilliam/cls_1521.urinalysis.htm)
27. Fretts RC. Etiology and prevention of stillbirth. *Am J Obstet Gynecol* 2005
28. Laroia N. Birth trauma. 2006. Diunduh dari : <http://www.emedicine.com/.htm>
29. Capeles EL, MEAD PB. Management of preterm premature of membrane. Lack of national concensus. *Am J Obstet Gynecol* 1987; 11: 157.
30. Vadillo ortega F, Gonzales avila G, Karchmer S, Cruz NM, Ayala Ruiz A, Lama MS. Collagen metabolism in premature rupture on amniotic membranes. *Obstet Gynecol* 1990; 75;84-8
31. Cecilia Maria Draque, Adriana Sañudo, Clovis de Araujo Peres and Maria Fernanda, Transcutaneous Bilirubin in Exclusively Breastfed Healthy Term Newborns Up to 12 Days of life. *Pediatrics*, 2011
32. Satrio Wibowo. Perbandingan kadar bilirubin neonatus dengan dan tanpa defisiensi Glucose-6-Phosphate Dehydrogenase, Infeksi dan tidak Infeksi [Internet]. 2007

33. M. Sholeh Kosim, dkk. Infeksi Neonatal Akibat Air Ketuban Keruh. Sari Pediatri . Available from: URL : <http://www.idai.or.id/saripediatri/abstrak.asp?q=620> . 2009
34. Rinawati Rohsiswatmo. Buku Indonesia Menyusui. Ikatan Dokter Anak Indonesia. Jakarta . 2009
35. Ga'rtner U, Goeser T, Wolkoff AW. Effect of fasting on the uptake of bilirubin and sulfobromophthalein by the isolated perfused rat liver. *Gastroenterology*. 1997
36. Giovanna Bertini, MD; Carlo Dani, MD; Michele Tronchin, PhD, dkk. Is Breastfeeding Really Favoring Early Neonatal Jaundice?. *PEDIATRICS* Vol. 107 No. 3 . 2001
37. Wu PYK, Hodgman JE, Kirkpatrick BV, et al. Metabolic aspects of phototherapy. *Pediatrics*. 1985
38. R Keren, V K Bhutani, X Luan, et al. Identifying newborns at risk of significant hyperbilirubinaemia: a comparison of two recommended approaches. Available from: URL : [www.archdischild.com](http://www.archdischild.com) . 2004
39. Williams W. Amniotic fluid: Objectives with narration and illustrations[Internet].2006[Update 2008 Oct 8; cited 2008 July 19].Available from: [www.CLS-1521](http://www.CLS-1521)
40. Armawati Ni Made. Kejadian hiperbilirubinemia berdasarkan pemberian minum bayi di ruang bayi RSUD PTPN XII tahun 2010. *Jurnal Penelitian Kesehatan Suara Forikes*. 2011
41. Utomo Martono Tri, Risk factor of Neonatal Sepsis : A Preliminary Study in Dr, Soetomo Hospital. Indonesian Journal of tropical and Infectious Disease. 2010.
42. Moelberg P, Johnson C, brown TS. Leukocytosis in labor : what are its implication ? .Available from: URL : <http://www.ncbi.nlm.nih.gov/pubmed>. 1994.
43. Deeksha Pandey, Saurav Bhagai, Vidhi Vanya, et al. Leukocyte count on admission as a predicto of clinical outcome in the expectant management for

women with preterm prelabor rupture of membranes. Sri Lanka Journal of Obstetrics and Gynaecology. 2011.

44. Murase Y, Tomoda Y, Imai N. Studies on meconium-like substances in the urine of mother and neonate during perinatal period. Available from: URL : <http://www.ncbi.nlm.nih.gov/pubmed/4078409> . 1985.

## LAMPIRAN 2. SURAT IJIN PENELITIAN



**RUMAH SAKIT  
HERMINA PANDANARAN**

Jl. Pandanaran No. 24 Semarang  
Telp. (024) 8442525; Fax. (024) 8450818

---

Semarang, 22 April 2013

Nomor : 224 /PERG/RSHPN/IV/2013  
Lamp. : -  
Perihal : Jawaban Permohonan Ijin Peminjaman Berkas Rekam Medik

Kepada Yth.  
Dekan Fakultas Kedokteran UNDIP  
Di -  
Tempat

Dengan hormat,  
Dasar :

- Surat dari Fak Kedokteran Undip tanggal 22 Maret 2013, No. 1261/UN7.3.4/D1/PP/2013, perihal Permohonan Ijin Peminjaman Rekam Medik.

Kami sampaikan bahwa kami memberikan ijin kepada mahasiswa tersebut dibawah ini untuk meminjam data rekam medik di RS Hermina Pandanaran.

No	Nama Mahasiswa	NRP	Judul / Topik
1	Sisca Rahardina	G2A 009 030	Hubungan Frekuensi Kencing dan Frekuensi Defekasi dengan Perubahan Berat Badan pada Neonatus
2	Ita Conita	G2A 009 029	Hubungan Penurunan Berat Badan dengan Kadar Bilirubin pada Neonatus Hari Ketiga Pasca Lahir
3	Nailul Khusna	G2A 009 156	Faktor Resiko Hiperbilirubinemia pada Neonatus Golongan Darah A atau B dan Ibu Golongan Darah O
4	Nur Ade Oktaviantil	G2A 009 153	Perbedaan Rerata Kadar Bilirubin pada Neonatus yang Mendapat ASI Eksklusif dan Tidak Eksklusif
5	Winda Ningsih	G2A 009 034	Hubungan Penurunan Berat Badan Neonatus dengan Kadar Bilirubin Hari Ketiga dan Kadar Akhir Minggu Pertama
6	Rizky Amalia Putri	G2A 009 087	Faktor Resiko Hiperbilirubinemia pada Neonatus

Pembimbing : Dr. dr. M. Mexitalla S, Sp.A (K)

Demikian kami sampaikan. Atas perhatian dan kerjasamanya diucapkan terima kasih.

Hormat kami,  
Direktur RS Hermina Pandanaran

dr. Hadi Wibowo, MMR 

## LAMPIRAN 4. HASIL UJI STATISTIK

### 1. Hiperbilirubin

#### a. Data karakteristik

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Yes	24	4,7	4,7	4,7
No	483	95,3	95,3	100,0
Total	507	100,0	100,0	

### 2. ASI eksklusif

#### a. Data karakteristik

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Yes	354	69,8	69,8	69,8
No	153	30,2	30,2	100,0
Total	507	100,0	100,0	

#### b. Uji normalitas

Excl Breastfeeding	Kolmogorov-Smirnov(a)			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
hiperbilirubin Yes	,539	354	,000	,153	354	,000
No	,533	153	,000	,325	153	,000

a Lilliefors Significance Correction

#### c. Crosstab

		hiperbilirubin		Total	
		Yes	No	Yes	
Excl Breastfeeding	Yes	Count	10	344	354
		Expected Count	16,8	337,2	354,0
		% of Total	2,0%	67,9%	69,8%
	No	Count	14	139	153
		Expected Count	7,2	145,8	153,0
		% of Total	2,8%	27,4%	30,2%
Total		Count	24	483	507
		Expected Count	24,0	483,0	507,0
		% of Total	4,7%	95,3%	100,0%

## d. Uji Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	9,478(b)	1	,002		
Continuity Correction(a)	8,128	1	,004		
Likelihood Ratio	8,582	1	,003		
Fisher's Exact Test				,005	,003
Linear-by-Linear Association	9,460	1	,002		
N of Valid Cases	507				

a Computed only for a 2x2 table

b 0 cells (,0%) have expected count less than 5. The minimum expected count is 7,24.

## e. Risk Estimate

	Value	95% Confidence Interval	
	Lower	Upper	Lower
Odds Ratio for Excl Breastfeeding (Yes / No)	,289	,125	,665
For cohort hiperbilirubin = Yes	,309	,140	,680
For cohort hiperbilirubin = No	1,070	1,014	1,128
N of Valid Cases	507		

## 3. Ketuban pecah dini (KPD)

## a. Data karakteristik

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Yes	44	8,7	8,7	8,7
No	463	91,3	91,3	100,0
Total	507	100,0	100,0	

## b. Uji Normality

Ketuban Pecah Dini	Kolmogorov-Smirnov(a)			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
hiperbilirubin Yes	,537	44	,000	,276	44	,000
No	,541	463	,000	,210	463	,000

a Lilliefors Significance Correction



## c. Crosstab

			hiperbilirubin		Total
			Yes	No	Yes
Ketuban Pecah Dini	Yes	Count	3	41	44
		Expected Count	2,1	41,9	44,0
		% of Total	,6%	8,1%	8,7%
	No	Count	21	442	463
		Expected Count	21,9	441,1	463,0
		% of Total	4,1%	87,2%	91,3%
Total	Count	24	483	507	
	Expected Count	24,0	483,0	507,0	
	% of Total	4,7%	95,3%	100,0%	

## d. Uji Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,464(b)	1	,496		
Continuity Correction(a)	,096	1	,757		
Likelihood Ratio	,416	1	,519		
Fisher's Exact Test				,454	,347
Linear-by-Linear Association	,463	1	,496		
N of Valid Cases	507				

a Computed only for a 2x2 table

b 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,08.

## e. Risk Estimate

	Value	95% Confidence Interval	
	Lower	Upper	Lower
Odds Ratio for Ketuban Pecah Dini (Yes / No)	1,540	,441	5,382
For cohort hiperbilirubin = Yes	1,503	,467	4,841
For cohort hiperbilirubin = No	,976	,899	1,060
N of Valid Cases	507		

#### 4. Ketuban keruh

##### a. Data karakteristik

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Yes	92	18,1	18,1	18,1
	No	415	81,9	81,9	100,0
	Total	507	100,0	100,0	

##### b. Uji Normality

	ketuban keruh	Kolmogorov-Smirnov(a)			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
hiperbilirubin	Yes	,540	92	,000	,171	92	,000
	No	,541	415	,000	,226	415	,000

a Lilliefors Significance Correction

##### c. Crosstab

			hiperbilirubin		Total
			Yes	No	Yes
ketuban keruh	Yes	Count	3	89	92
		Expected Count	<b>4,4</b>	<b>87,6</b>	<b>92,0</b>
		% of Total	,6%	17,6%	18,1%
	No	Count	21	394	415
		Expected Count	19,6	395,4	415,0
		% of Total	4,1%	77,7%	81,9%
Total	Count	24	483	507	
	Expected Count	24,0	483,0	507,0	
	% of Total	4,7%	95,3%	100,0%	

##### d. Uji Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,541(b)	1	,462		
Continuity Correction(a)	,215	1	,643		
Likelihood Ratio	,591	1	,442		
Fisher's Exact Test				<b>,595</b>	<b>,338</b>
Linear-by-Linear Association	,540	1	,463		
N of Valid Cases	507				

a Computed only for a 2x2 table

b 1 cells (25,0%) have expected count less than 5. The minimum expected count is 4,36.

## e. Risk Estimate

	Value	95% Confidence Interval	
	Lower	Upper	Lower
Odds Ratio for ketuban keruh (Yes / No)	,632	,185	2,167
For cohort hiperbilirubin = Yes	<b>,644</b>	<b>,196</b>	<b>2,115</b>
For cohort hiperbilirubin = No	1,019	,975	1,064
N of Valid Cases	507		

5. Infeksi ibu

## a. ibu infeksi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Yes	60	11,8	11,8	11,8
	No	447	88,2	88,2	100,0
	Total	507	100,0	100,0	

## b. Uji Normality

ibu infeksi		Kolmogorov-Smirnov(a)			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
hiperbilirubin	Yes	,540	60	,000	,175	60	,000
	No	,541	447	,000	,222	447	,000

a Lilliefors Significance Correction

## c. Crosstab

			hiperbilirubin		Total
			Yes	No	Yes
ibu infeksi	Yes	Count	2	58	60
		Expected Count	2,8	57,2	60,0
		% of Total	,4%	11,4%	11,8%
	No	Count	22	425	447
		Expected Count	21,2	425,8	447,0
		% of Total	4,3%	83,8%	88,2%
Total	Count	24	483	507	
	Expected Count	24,0	483,0	507,0	
	% of Total	4,7%	95,3%	100,0%	

## d. Uji Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,296(b)	1	,586		
Continuity Correction(a)	,049	1	,826		
Likelihood Ratio	,324	1	,569		
Fisher's Exact Test				1,000	,443
Linear-by-Linear Association	,295	1	,587		
N of Valid Cases	507				

a Computed only for a 2x2 table

b 1 cells (25,0%) have expected count less than 5. The minimum expected count is 2,84.

## e. Risk Estimate

	Value	95% Confidence Interval	
	Lower	Upper	Lower
Odds Ratio for ibu infeksi (Yes / No)	,666	,153	2,907
For cohort hiperbilirubin = Yes	,677	,163	2,808
For cohort hiperbilirubin = No	1,017	,966	1,070
N of Valid Cases	507		

## LAMPIRAN 5. DOKUMENTASI



Proses pencarian dokumen rekam medis



Proses pencarian data subyek penelitian



Proses pencarian data subyek penelitian



Proses input data subyek penelitian ke dalam komputer



Penulis beserta karyawan rekam medis

## **LAMPIRAN 6. DAFTAR RIWAYAT HIDUP**

### **Identitas**

Nama : Rizky Amalia Putri  
NIM : G2A 009 087  
Tempat/tanggal lahir : Gresik, 08 Desember 1990  
Jenis kelamin : Perempuan  
Alamat : Jl. Merak Raya no. 33 Rewwin Kel. Wedoro ,  
Kec. Waru , Kab. Sidoarjo Jawa Timur  
Nomor Telepon/HP : (031) 8532872 / 083849692293  
Alamat email : rizkyamaliaputri\_87@yahoo.com

### **Riwayat Pendidikan Formal**

1. SD : SD Al – Hikmah Surabaya
2. SMP : SMP Al – Hikmah Surabaya, lulus tahun : 2006
3. SMA : SMA Negeri 15 Surabaya, lulus tahun : 2009
4. FK UNDIP : Masuk tahun : 2009

### **Keanggotaan organisasi**

1. Asisten Biologi FK Undip Tahun 2011
2. Asisten Biologi FK Undip Tahun 2012
3. Sie logistik Scientific Fair FK Undip Tahun 2011
4. Sie logistik Penerimaan Mahasiswa Baru FK Undip Tahun 2010
5. Team Angklung FK Undip Tahun 2012