

DAFTAR PUSTAKA

1. Rudan I, Boschi-Pinto C, Bilolav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. Bull World Health Organ 2008;86:408-16.
2. Wardlaw T, Johansson EW, Hodge M. Pneumonia: The Forgotten Killer of Children. UNICEF's Division of Communication. New York: WHO Press; 2006.
3. Said, M., 2010, *Pneumonia Penyebab Utama Mortalitas Anak Balita Di Indonesia*, [Http://www.idai.or.id](http://www.idai.or.id).
4. Ag. Soermantri D. *Buku Ajar Ilmu Kesehatan Anak*. Semarang : Fakultas Kedokteran Universitas diponegoro
5. Subhi R, Adamson M, Campbell H, Weber M, Smith K, Duke T, et al. The prevalence of hypoxaemia among ill children in developing countries: A systematic review. Lancet Infect Dis 2009;9:219-27.
6. Abdulkadir MB, Ibraheem RM, Gobir AA, Johnson WBR. Hypoxaemia as a measure of disease severity in young hospitalised Nigerian children with pneumonia : A cross-sectional study. 2015;9(2). doi:10.7196/SAJCH.901.
7. Husian NOSA, Salah E, Algasim S, Mhamoud A. Prevalence of hypoxemia in under-five children with pneumonia in an emergency pediatrics hospital in Sudan. Indian J Crit Care Med. 2015;19(4):203. doi:10.4103/0972-5229.154549.
8. Jain DL, Sarathi V, Jawalekar S. Predictors of treatment failure in hospitalized children [3-59 months] with severe and very severe

- pneumonia. Indian Pediatr 2013;50:787-9.
9. Marcdante.karen J kliegman. rober. M. *Buku Ajar Ilmu Kesehatan Anak Nelson*. keenam. (Tharemapalan S, ed.). Singapore: elsevier Pte Ltd; 2014.
 10. Dahlan Z. *Pneumonia. Dalam: Sudoyo AW, Setiyohadi B,Alwi I, Simadibrata M, Setiati S, Editor. Buku Ajar Ilmu Penyakit Dalam Jilid III*. v. Jakarta: Interna Publishing; 2009.
 11. Levine OS, O'Brien KL, Deloria-Knoll M, et al. The pneumonia etiology research for child health project: A 21st century childhood pneumonia etiology study. *Clin Infect Dis*. 2012;54(SUPPL. 2):93-102.
doi:10.1093/cid/cir1052.
 12. Sectish T PC. *Pneumonia. In: Kliegman RM, Behrman RE, Jenson HB, Stanton BF, Eds. Nelson Textbook of Pediatrics*. 18th ed. Philadelphia: Saunders Elsevier; 2008.
 13. Bennet NJ, Domachowske J. Pediatric pneumonia. Available from:
<http://emedicine.medscape.com/article/967822-overview>. Diakses 31 Januari 2016.
 14. Chart booklet: Integrated Management of Childhood Illness (IMCI). Geneva, World Health Organization/The United Nation Children's Fund (UNICEF), 2008.
 15. Integrated Management of Childhood Illness (IMCI). WHO recommendations on the management of diarrhoea and pneumonia in HIV-infected infants and children. Geneva: World Health Organization; 2010.
 16. Sunyatatingkamto, Iskandar Z, Alan RT, Budiman I, Surjono A, Wibowo

- T et al. The role of indoor air pollution and other factors in the incidence of pneumonia in children under-five. *Pediatrica Indonesiana*. 2004;44:25-9.
17. Huong P Le, Hien PT, Lan NT, et al. First report on prevalence and risk factors of severe atypical pneumonia in Vietnamese children aged 1 – 15 years. 2014. doi:10.1186/1471-2458-14-1304.
 18. Sulaeman, Endang Sutisna, 2011. *Manajemen Kesehatan Teori dan Praktik di Puskesmas*. Bandung : PT Remaja Rusdakarya.
 19. WHO. 2011. World Pneumonia Day
http://www.who.int/maternal_child_adolescent/news_events/events/2011/world_pneumonia_day/en/. Diakses tanggal 29 Januari 2016.
 20. DepKes RI. Riskesdas. Profil Kesehatan Indonesia 2004. 2006.
 21. Broor S, Pandey RM, Ghosh M, Maitreyi RS, Lodha R, Singhal T, et al. Risk factors for severe acute lower respiratory tract infection in children under-five. *Indian Pediatr*. 2001;38:1361-9.
 22. Graham NMH. The epidemiology of acute respiratory infections in children and adults: A global perspective. *Epidemiol Rev*. 1990;22:149-78.
 23. Nataprawira HM, Alwi EH, Adriani N. Faktor Risiko Morbiditas dan Mortalitas Pneumonia Berat pada Anak Usia Balita. *Major Kedokt Indones*. 2009;60:443-447.
 24. Fakunle GA, Ana GR, Ayede AI. Environmental risk factors for acute respiratory infections in hospitalized children under 5 years of age in Ibadan, Nigeria. *Paediatr Int Child Health*. 2014;34(2):120-124.
doi:10.1179/2046905513Y.0000000107.

25. WHO. Pelayanan kesehatan anak di rumah sakit. *Kesehatan*. 2013;2(pelayanan masyarakat):2. [http://gooleknah.nmetreisnbs.2013;2\(pelayanan masyarakat\):2](http://gooleknah.nmetreisnbs.2013;2(pelayanan masyarakat):2).
26. Basnet S, Adhikari RK, Gurung CK. Hypoxemia in children with pneumonia and its clinical predictors. *Indian J Pediatr*. 2006;73(9):777-781. doi:10.1007/BF02790384.
27. Regelmann WE, Hill HR, Cates KL, Quie PG. Immunology of the newborn. In: Feigin RD, Cherry JD, eds. *Textbook of Pediatric Infectious Diseases*. 3rd ed. Philadelphia: WB Saunders Company, Harcourt Brace Jovanovich Inc, 1992 876–887.
28. De Graaff JC, Bijker JB, Kappen TH, van Wolfswinkel L, Zuithoff NP, Kalkman CJ. Incidence of intraoperative hypoxemia in children in relation to age. *Anesth Analg* 2013;117:169-75.
<http://www.ncbi.nlm.nih.gov/pubmed/23687233>.
29. Lozano, J.M. Epidemiology of hypoxemia in children with acute lower respiratory tract infection. *Int J Tuberc Lung Dis* 2001; 5(6): 496-504.
30. Graham SM, English M, Hazir T, Enarson P, Duke T. Challenges to improving case management of childhood pneumonia at health facilities in resource-limited settings. *Bull World Health Organ*. 2008;86:349-55.
31. Rudan I, Boschi-Pinto C, Bilonlav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. *Bull World Health Organ*. 2008;86(5):408-416. doi:10.2471/BLT.07.048769.
32. Zhang Q, Guo Z, Bai Z MN. A 4 year prospective study to determine risk factors for severe community acquired pneumonia in children in southern


- China. Pediatr Pulmonol. 2012 Jul 6. doi: 10.1002/ ppul.22608. [Epub ahead of print].
33. Tiewsoh K, Lodha R, Pandey RM, Broor S, Kalaivani M KS. Factors determining the outcome of children hospitalized with severe pneumonia. BMC pediatrics. 2009;9:15.
 34. Wiens MO, Pawluk S, Kissoon N, et al. Pediatric post-discharge mortality in resource poor countries: a systematic review. PLoS One 2013; 8: e66698.
 35. Rytter MJ, Kolte L, Briend A, Friis H CV. The immune system in children with malnutrition—a systematic review. PLoS One 2014; 9: e105017.
 36. Chisti MJ, Tebruegge M, La Vincente S, Graham SM, Duke T. Pneumonia in severely malnourished children in developing countries—mortality risk, aetiology and validity of WHO clinical signs: a systematic review. Trop Med Int Health 2009; 14: 1173–89.
 37. Reed C, Madhi SA, Klugman KP, et al. Development of the respiratory index of severity in children (RISC) score among young children with respiratory infections in South Africa. PLoS One 2012; 7: e27793.
 38. Victora CG, Kirkwood BR, Ashworth A et al. (1999) Potential interventions for prevention of childhood pneumonia in developing countries: improving nutrition. The American Journal of Clinical Nutrition 70, 309–320.
 39. Subhi, R, Adamson, M, Campbell, H, Weber, M, Smith, K, and Duke, T. The prevalence of hypoxaemia among ill children in developing countries:

- a systematic review. *Lancet Infect Dis.* 2009; 9: 219–227.
40. Djelantik IG, Gessner BD, Sutanto A, Steinhoff M, Linehan M, Moulton LH, et al. Case fatality proportions and predictive factors for mortality among children hospitalized with severe pneumonia in a rural developing country setting. *J Trop Pediatr.* 2003;49.
41. West TE, Goetghebuer T, Milligan P, Mulholland EK, Weber MW. Long-term morbidity and mortality following hypoxaemic lower respiratory tract infection in Gambian children. *Bull World Health Organ.* 1999;77(2):144-148.
- <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2557604&tool=pmcentrez&rendertype=abstract>.
42. Ramachandran P, Nedunchelian K, Vengatesan A, Suresh S. Risk factors for mortality in community acquired pneumonia among children aged 1-59 months admitted in a referred hospital. *Indian Pediatr.* 2012;48:889-95.
43. Factors R, Hospitalized C. Risk Factors for Mortality in Children Hospitalized with Pneumonia. *Indian Pediatr.* 2012.
doi:10.1002/ppul.22608.
44. *Sylvia , A , Lorraine , MW. Patofisiologi:konsep Klinis Proses-Proses Penyakit. Buku Kedokteran EGC, 2014.*

Lampiran 1. Ethical clearance



Lampiran 2. Surat izin penelitian



Lampiran 3. Form penelitian

FORM DATA PENELITIAN

IDENTITAS PASIEN

No CM :
 Nama Penderita :
 Umur / Tgl Lahir :
 Jenis Kelamin :
 Alamat :

 Tanggal Masuk :
 Tanggal Keluar :

IDENTITAS ORANG TUA

Ayah
 Nama :
 Umur :
 Pendidikan :
 Pekerjaan :

 Ibu
 Nama :
 Umur :
 Pendidikan :
 Pekerjaan :

ANTROPOMETRI

BB Lahir	:	Kg
PB Lahir	:	cm
BB Sekarang	:	Kg
PB / TB Sekarang	:	cm
BB Bulan Lalu	:	Kg
WAZ	:	
HAZ	:	
WHZ	:	
Status Gizi	:	

Keluhan Utama	
----------------------	--

Gejala Respiratori	Indikator	Onset
- Batuk	Ya / Tidak	
- Sesak	Ya / Tidak	
- Sianosis	Ya / Tidak	
- Nyeri Dada	Ya / Tidak	

Gejala Lain	Indikator	Onset
-Demam		
-		
-		

Riwayat Imunisasi	
--------------------------	--

Riwayat Makan dan Minum	Keterangan
ASI Ekslusif	Ya / Tidak
MP ASI	Ya / Tidak

Riwayat Pertumbuhan dan Perkembangan	
Riwayat Penyakit Terdahulu	

Lampiran 4. Analisis statistik SPSS

Case Summaries

Umur

N	Mean	Std. Deviation	Median	Minimum	Maximum
78	12.42	11.566	7.00	2	53

Frequencies

Frequency Table

SaO2

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid < 95	26	33.3	33.3	33.3
>= 95	52	66.7	66.7	100.0
Total	78	100.0	100.0	

Jenis kelamin

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Laki-laki	43	55.1	55.1	55.1
Perempuan	35	44.9	44.9	100.0
Total	78	100.0	100.0	

Status gizi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Baik	39	50.0	50.0	50.0
Kurang	13	16.7	16.7	66.7
Buruk	26	33.3	33.3	100.0
Total	78	100.0	100.0	

ASI Eksklusif

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ya	55	70.5	70.5	70.5
	Tidak	23	29.5	29.5	100.0
	Total	78	100.0	100.0	

Derajat pneumonia

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Pneumonia	17	21.8	21.8	21.8
	Berat	52	66.7	66.7	88.5
	Sangat berat	9	11.5	11.5	100.0
	Total	78	100.0	100.0	

Antibiotik

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tunggal	38	48.7	48.7	48.7
	Kombinasi	40	51.3	51.3	100.0
	Total	78	100.0	100.0	

Respon terapi

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Tidak	28	35.9	35.9	35.9
	Membaik	50	64.1	64.1	100.0
	Total	78	100.0	100.0	

Lama rawat

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	>= 7 hari	23	29.5	29.5	29.5
	< 7 hari	55	70.5	70.5	100.0
	Total	78	100.0	100.0	

Status of Discharge

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Meninggal	5	6.4	6.4	6.4
	Sembuh	73	93.6	93.6	100.0
	Total	78	100.0	100.0	

Komorbid

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ada	44	56.4	56.4	56.4
	Tidak	34	43.6	43.6	100.0
	Total	78	100.0	100.0	

Explore

SaO₂

Case Summaries

Umur


SaO ₂	N	Mean	Std. Deviation	Median	Minimum	Maximum
< 95	26	14.12	12.962	8.00	2	53
=> 95	52	11.58	10.836	7.00	2	50
Total	78	12.42	11.566	7.00	2	53

Tests of Normality

SaO ₂	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Umur	< 95	.220	26	.002	.839	26
	=> 95	.202	52	.000	.810	52

a. Lilliefors Significance Correction

Umur



NPar Tests**Mann-Whitney Test****Ranks**

SaO2	N	Mean Rank	Sum of Ranks
Umur < 95	26	42.56	1106.50
>= 95	52	37.97	1974.50
Total	78		

Test Statistics^a

	Umur
Mann-Whitney U	596.500
Wilcoxon W	1974.500
Z	-.845
Asy mp. Sig. (2-tailed)	.398

a. Grouping Variable: SaO2

Ranks

	Respon terapi	N	Mean Rank	Sum of Ranks
Umur	Tidak	28	40,36	1130,00
	Membuat	50	39,02	1951,00
	Total	78		

Test Statistics^a

	Umur
Mann-Whitney U	676,000
Wilcoxon W	1951,000
Z	-,251
Asymp. Sig. (2-tailed)	,802

a. Grouping Variable: Respon terapi

Ranks

	Lama rawat	N	Mean Rank	Sum of Ranks
	>= 7 hari	23	40,85	939,50
Umur	< 7 hari	55	38,94	2141,50
	Total	78		

Test Statistics^a

	Umur
Mann-Whitney U	601,500
Wilcoxon W	2141,500
Z	-,341
Asymp. Sig. (2-tailed)	,733

a. Grouping Variable: Lama rawat

Ranks

	Status of Discharge	N	Mean Rank	Sum of Ranks
	Meninggal	5	42,70	213,50
Umur	Sembuh	73	39,28	2867,50
	Total	78		

Test Statistics^a

	Umur
Mann-Whitney U	166,500
Wilcoxon W	2867,500
Z	-,327
Asymp. Sig. (2-tailed)	,744
Exact Sig. [2*(1-tailed Sig.)]	,752 ^b

a. Grouping Variable: Status of Discharge

b. Not corrected for ties.

Crosstabs

SaO2 * Jenis kelamin

Crosstab

			Jenis kelamin		Total
			Laki-laki	Perempuan	
SaO2 < 95	Count		15	11	26
	Expected Count		14.3	11.7	26.0
	% within Jenis kelamin		34.9%	31.4%	33.3%
	Count		28	24	52
	Expected Count		28.7	23.3	52.0
	% within Jenis kelamin		65.1%	68.6%	66.7%
Total	Count		43	35	78
	Expected Count		43.0	35.0	78.0
	% within Jenis kelamin		100.0%	100.0%	100.0%

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.104 ^b	1	.747		
Continuity Correction ^a	.006	1	.936		
Likelihood Ratio	.104	1	.747		
Fisher's Exact Test				.812	.469
Linear-by-Linear Association	.102	1	.749		
N of Valid Cases	78				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11. 67.

SaO₂ * Status gizi

Crosstab

		Status gizi			Total
		Baik	Kurang	Buruk	
SaO ₂	< 95	Count	12	3	26
		Expected Count	13.0	4.3	26.0
		% within Status gizi	30.8%	23.1%	33.3%
	>= 95	Count	27	10	52
		Expected Count	26.0	8.7	52.0
		% within Status gizi	69.2%	76.9%	66.7%
	Total	Count	39	13	78
		Expected Count	39.0	13.0	78.0
		% within Status gizi	100.0%	100.0%	100.0%

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)
Pearson Chi-Square	1.673 ^a	2	.433
Likelihood Ratio	1.680	2	.432
Linear-by -Linear Association	.786	1	.375
N of Valid Cases	78		

a. 1 cells (16.7%) have expected count less than 5. The minimum expected count is 4.33.

SaO₂ * ASI Eksklusif**Crosstab**

		ASI Eksklusif		Total
		Ya	Tidak	
SaO ₂	< 95	Count	18	26
		Expected Count	18.3	26.0
		% within ASI Eksklusif	32.7%	33.3%
	>= 95	Count	37	52
		Expected Count	36.7	52.0
		% within ASI Eksklusif	67.3%	66.7%
	Total	Count	55	78
		Expected Count	55.0	78.0
		% within ASI Eksklusif	100.0%	100.0%

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.031 ^b	1	.861		
Continuity Correction ^a	.000	1	1.000		
Likelihood Ratio	.031	1	.861		
Fisher's Exact Test				1.000	.530
Linear-by-Linear Association	.030	1	.862		
N of Valid Cases	78				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.

SaO₂ * Derajat pneumonia

Crosstab

			Derajat pneumonia			Total
			Pneumonia	Berat	Sangat berat	
SaO ₂	< 95	Count	2	21	3	26
		Expected Count	5.7	17.3	3.0	26.0
		% within Derajat pneumonia	11.8%	40.4%	33.3%	33.3%
	>= 95	Count	15	31	6	52
		Expected Count	11.3	34.7	6.0	52.0
		% within Derajat pneumonia	88.2%	59.6%	66.7%	66.7%
	Total	Count	17	52	9	78
		Expected Count	17.0	52.0	9.0	78.0
		% within Derajat pneumonia	100.0%	100.0%	100.0%	100.0%

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)
Pearson Chi-Square	4.722 ^a	2	.094
Likelihood Ratio	5.372	2	.068
Linear-by-Linear Association	2.372	1	.124
N of Valid Cases	78		

a. 1 cells (16.7%) have expected count less than 5. The minimum expected count is 3.00.

SaO₂ * Antibiotik

Crosstab

		Antibiotik		Total	
		Tunggal	Kombinasi		
SaO ₂	< 95	Count	12	26	
		Expected Count	12.7	26.0	
		% within Antibiotik	31.6%	33.3%	
	>= 95	Count	26	52	
		Expected Count	25.3	52.0	
		% within Antibiotik	68.4%	66.7%	
Total		Count	38	78	
		Expected Count	38.0	78.0	
		% within Antibiotik	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.103 ^b	1	.749		
Continuity Correction ^a	.006	1	.936		
Likelihood Ratio	.103	1	.749		
Fisher's Exact Test				.813	.468
Linear-by -Linear Association	.101	1	.750		
N of Valid Cases	78				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 12.67.

SaO₂ * Respon terapi

Crosstab

			Respon terapi		Total
			Tidak	Membaike	
SaO ₂	< 95	Count	13	13	26
		Expected Count	9.3	16.7	26.0
		% within Respon terapi	46.4%	26.0%	33.3%
	>= 95	Count	15	37	52
		Expected Count	18.7	33.3	52.0
		% within Respon terapi	53.6%	74.0%	66.7%
Total		Count	28	50	78
		Expected Count	28.0	50.0	78.0
		% within Respon terapi	100.0%	100.0%	100.0%

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.371 ^b	1	.066		
Continuity Correction	2.514	1	.113		
Likelihood Ratio	3.317	1	.069		
Fisher's Exact Test				.083	.057
Linear-by-Linear Association	3.327	1	.068		
N of Valid Cases	78				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 9.

33.

SaO₂ * Lama rawat

Crosstab

			Lama rawat		Total
			>= 7 hari	< 7 hari	
SaO ₂	< 95	Count	12	14	26
		Expected Count	7.7	18.3	26.0
		% within Lama rawat	52.2%	25.5%	33.3%
	>= 95	Count	11	41	52
		Expected Count	15.3	36.7	52.0
		% within Lama rawat	47.8%	74.5%	66.7%
Total	Count	23	55	78	
	Expected Count	23.0	55.0	78.0	
	% within Lama rawat	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	5.210 ^b	1	.022		
Continuity Correction ^a	4.077	1	.043		
Likelihood Ratio	5.055	1	.025		
Fisher's Exact Test				.034	.023
Linear-by-Linear Association	5.143	1	.023		
N of Valid Cases	78				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.

Risk Estimate

	Value	95% Confidence Interval	
		Lower	Upper
Odds Ratio for SaO ₂ (< 95 / >= 95)	3.195	1.154	8.847
For cohort Lama rawat = >= 7 hari	2.182	1.117	4.260
For cohort Lama rawat = < 7 hari	.683	.466	1.001
N of Valid Cases	78		

SaO₂ * Status of Discharge

Crosstab

		Status of Discharge		Total	
		Meninggal	Sembuh		
SaO ₂	< 95	Count	2	26	
		Expected Count	1.7	26.0	
		% within Status of Discharge	40.0%	33.3%	
	>= 95	Count	3	52	
		Expected Count	3.3	52.0	
		% within Status of Discharge	60.0%	66.7%	
Total		Count	5	78	
		Expected Count	5.0	78.0	
		% within Status of Discharge	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.107 ^b	1	.744		
Continuity Correction ^a	.000	1	1.000		
Likelihood Ratio	.104	1	.747		
Fisher's Exact Test				1.000	.543
Linear-by-Linear Association	.105	1	.745		
N of Valid Cases	78				

a. Computed only for a 2x2 table

b. 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.

67.

SaO₂ * Komorbid

Crosstab

		Komorbid		Total	
		Ada	Tidak		
SaO ₂	< 95	Count	17	9	
		Expected Count	14.7	11.3	
		% within Komorbid	38.6%	26.5%	
	>= 95	Count	27	25	
		Expected Count	29.3	22.7	
		% within Komorbid	61.4%	73.5%	
Total		Count	44	34	
		Expected Count	44.0	34.0	
		% within Komorbid	100.0%	100.0%	

Chi-Square Tests

	Value	df	Asy mp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.277 ^b	1	.258		
Continuity Correction ^a	.789	1	.375		
Likelihood Ratio	1.293	1	.255		
Fisher's Exact Test				.335	.188
Linear-by-Linear Association	1.261	1	.261		
N of Valid Cases	78				

a. Computed only for a 2x2 table

b. 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.33.

Jenis kelamin * Respon terapi

Crosstabulation

		Respon terapi		Total
		Tidak	Membaik	
Jenis kelamin	Count	17	26	43
	% within Jenis kelamin	39,5%	60,5%	100,0%
	Count	11	24	35
	% within Jenis kelamin	31,4%	68,6%	100,0%
	Count	28	50	78
	% within Jenis kelamin	35,9%	64,1%	100,0%
Total				

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	,551 ^a	1	,458		
Continuity Correction ^b	,255	1	,614		
Likelihood Ratio	,554	1	,457		
Fisher's Exact Test				,487	,308
Linear-by-Linear Association	,544	1	,461		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 12,56.

b. Computed only for a 2x2 table

Jenis kelamin * Lama rawat

Crosstab

		Lama rawat		Total
		>= 7 hari	< 7 hari	
Jenis kelamin	Count	11	32	43
	% within Jenis kelamin	25,6%	74,4%	100,0%
	Count	12	23	35
	% within Jenis kelamin	34,3%	65,7%	100,0%
	Count	23	55	78
	% within Jenis kelamin	29,5%	70,5%	100,0%
Total				

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	,703 ^a	1	,402		
Continuity Correction ^b	,347	1	,556		
Likelihood Ratio	,701	1	,402		
Fisher's Exact Test				,459	,277
Linear-by-Linear Association	,694	1	,405		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 10,32.

b. Computed only for a 2x2 table

Jenis kelamin * Status of discharge

Crosstab

		Status of Discharge		Total
		Meninggal	Sembuh	
Jenis kelamin	Laki-laki	Count	4	43
		% within Jenis kelamin	9,3%	90,7%
	Perempuan	Count	1	35
		% within Jenis kelamin	2,9%	97,1%
Total		Count	5	78
		% within Jenis kelamin	6,4%	93,6%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	1,336 ^a	1	,248		
Continuity Correction ^b	,478	1	,489		
Likelihood Ratio	1,448	1	,229		
Fisher's Exact Test				,372	,250
Linear-by-Linear Association	1,319	1	,251		
N of Valid Cases	78				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,24.

b. Computed only for a 2x2 table

Status gizi * Lama rawat

Crosstab

		Lama rawat		Total
		>= 7 hari	< 7 hari	
Status gizi	Baik	Count	9	39
	Baik	% within Status gizi	23,1%	76,9%
	Kurang	Count	5	13
	Kurang	% within Status gizi	38,5%	61,5%
	Buruk	Count	9	26
	Buruk	% within Status gizi	34,6%	65,4%
Total		Count	23	78
		% within Status gizi	29,5%	70,5%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,603 ^a	2	,449
Likelihood Ratio	1,606	2	,448
Linear-by-Linear Association	1,110	1	,292
N of Valid Cases	78		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 3,83.

Status gizi * Status of discharge

Crosstab

		Status of Discharge		Total
		Meninggal	Sembuh	
Status gizi	Baik	Count	1	39
	Baik	% within Status gizi	2,6%	97,4%
	Kurang	Count	1	13
	Kurang	% within Status gizi	7,7%	92,3%
	Buruk	Count	3	26
	Buruk	% within Status gizi	11,5%	88,5%
Total		Count	5	78
		% within Status gizi	6,4%	93,6%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2,137 ^a	2	,344
Likelihood Ratio	2,196	2	,333
Linear-by-Linear Association	2,102	1	,147
N of Valid Cases	78		

a. 3 cells (50,0%) have expected count less than 5. The minimum expected count is ,83.

Status gizi * Respon terapi

Crosstab

		Respon terapi		Total
		Tidak	Membaik	
Status gizi	Baik	Count	18	39
	Baik	% within Status gizi	46,2%	53,8% 100,0%
	Kurang	Count	3	13
Status gizi	Kurang	% within Status gizi	23,1%	76,9% 100,0%
	Buruk	Count	7	26
	Buruk	% within Status gizi	26,9%	73,1% 100,0%
Total		Count	28	78
		% within Status gizi	35,9%	64,1% 100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3,621 ^a	2	,164
Likelihood Ratio	3,671	2	,159
Linear-by-Linear Association	2,739	1	,098
N of Valid Cases	78		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 4,67.

ASI eksklusif * Lama rawat

Crosstab

		Lama rawat		Total
		>= 7 hari	< 7 hari	
ASI Eksklusif	Count	16	39	55
	% within ASI	29,1%	70,9%	100,0%
	Eksklusif			
	Count	7	16	23
	% within ASI	30,4%	69,6%	100,0%
	Eksklusif			
Total	Count	23	55	78
	% within ASI	29,5%	70,5%	100,0%
	Eksklusif			

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	,014 ^a	1	,906		
Continuity Correction ^b	,000	1	1,000		
Likelihood Ratio	,014	1	,906		
Fisher's Exact Test				1,000	,555
Linear-by-Linear Association	,014	1	,906		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 6,78.

b. Computed only for a 2x2 table

ASI eksklusif * Status of discharge

Crosstab

		Status of Discharge		Total
		Meninggal	Sembuh	
ASI Eksklusif	Count	2	53	55
	% within ASI	3,6%	96,4%	100,0%
	Eksklusif			
	Count	3	20	23
	% within ASI	13,0%	87,0%	100,0%
	Eksklusif			
Total	Count	5	73	78
	% within ASI	6,4%	93,6%	100,0%
	Eksklusif			

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	2,392 ^a	1	,122		
Continuity Correction ^b	1,081	1	,298		
Likelihood Ratio	2,150	1	,143		
Fisher's Exact Test				,149	,149
Linear-by-Linear Association	2,362	1	,124		
N of Valid Cases	78				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 1,47.

b. Computed only for a 2x2 table

ASI eksklusif * Respon terapi**Crosstab**

		Respon terapi		Total
		Tidak	Membalik	
ASI Eksklusif	Ya	Count	21	55
		% within ASI	38,2%	61,8%
		Eksklusif		100,0%
	Tidak	Count	7	23
Total		% within ASI	30,4%	69,6%
		Eksklusif		100,0%
		Count	28	78
		% within ASI	35,9%	64,1%
		Eksklusif		100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	,423 ^a	1	,515		
Continuity Correction ^b	,153	1	,695		
Likelihood Ratio	,429	1	,512		
Fisher's Exact Test				,609	,351
Linear-by-Linear Association	,418	1	,518		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 8,26.

b. Computed only for a 2x2 table

Derajat pneumonia * Lama rawat**Crosstab**

		Lama rawat		Total
		>= 7 hari	< 7 hari	
Derajat pneumonia	Pneumonia	Count	3	17
		% within Derajat pneumonia	17,6%	82,4%
	Berat	Count	15	52
		% within Derajat pneumonia	28,8%	71,2%
	Sangat berat	Count	5	9
		% within Derajat pneumonia	55,6%	44,4%
Total		Count	23	78
		% within Derajat pneumonia	29,5%	70,5%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4,098 ^a	2	,129
Likelihood Ratio	3,918	2	,141
Linear-by-Linear Association	3,583	1	,058
N of Valid Cases	78		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 2,65.

Derajat pneumonia * Status of discharge**Crosstab**

		Status of Discharge		Total
		Meninggal	Sembuh	
Derajat pneumonia	Pneumonia	Count	1	17
		% within Derajat pneumonia	5,9%	94,1%
	Berat	Count	4	52
		% within Derajat pneumonia	7,7%	92,3%
	Sangat berat	Count	0	9
		% within Derajat pneumonia	0,0%	100,0%
Total		Count	5	78
		% within Derajat pneumonia	6,4%	93,6%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	,767 ^a	2	,682
Likelihood Ratio	1,335	2	,513
Linear-by-Linear Association	,155	1	,694
N of Valid Cases	78		

a. 3 cells (50,0%) have expected count less than 5. The minimum expected count is ,58.

Derajat pneumonia * Respon terapi**Crosstab**

		Respon terapi		Total
		Tidak	Membaik	
Derajat pneumonia	Pneumonia	Count	5	17
		% within Derajat pneumonia	29,4%	70,6%
		Count	18	52
	Berat	% within Derajat pneumonia	34,6%	65,4%
		Count	5	9
	Sangat berat	% within Derajat pneumonia	55,6%	44,4%
Total		Count	28	78
		% within Derajat pneumonia	35,9%	64,1%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1,859 ^a	2	,395
Likelihood Ratio	1,795	2	,408
Linear-by-Linear Association	1,405	1	,236
N of Valid Cases	78		

a. 1 cells (16,7%) have expected count less than 5. The minimum expected count is 3,23.

Antibiotik * Lama rawat**Crosstab**

		Lama rawat		Total
		>= 7 hari	< 7 hari	
Antibiotik	Tunggal	Count	9 23,7%	29 76,3% 100,0%
	Kombinasi	Count	14 35,0%	26 65,0% 100,0%
	Total	Count	23 29,5%	55 70,5% 100,0%
		% within Antibiotik		

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1,200 ^a	1	,273		
Continuity Correction ^b	,718	1	,397		
Likelihood Ratio	1,208	1	,272		
Fisher's Exact Test				,326	,199
Linear-by-Linear Association	1,185	1	,276		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 11,21.

b. Computed only for a 2x2 table

Antibiotik * Status of discharge**Crosstab**

		Status of Discharge		Total
		Meninggal	Sembuh	
Antibiotik	Tunggal	Count	2 5,3%	36 94,7% 100,0%
	Kombinasi	Count	3 7,5%	37 92,5% 100,0%
	Total	Count	5 6,4%	73 93,6% 100,0%
		% within Antibiotik		

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	,163 ^a	1	,687		
Continuity Correction ^b	,000	1	1,000		
Likelihood Ratio	,164	1	,686		
Fisher's Exact Test				1,000	,525
Linear-by-Linear Association	,160	1	,689		
N of Valid Cases	78				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,44.

b. Computed only for a 2x2 table

Antibiotik * Respon terapi**Crosstab**

		Respon terapi		Total
		Tidak	Membaik	
Antibiotik	Tunggal	Count	11	38
		% within Antibiotik	28,9%	71,1% 100,0%
	Kombinasi	Count	17	40
		% within Antibiotik	42,5%	57,5% 100,0%
Total		Count	28	78
		% within Antibiotik	35,9%	64,1% 100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	1,555 ^a	1	,212		
Continuity Correction ^b	1,022	1	,312		
Likelihood Ratio	1,565	1	,211		
Fisher's Exact Test				,244	,156
Linear-by-Linear Association	1,536	1	,215		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 13,64.

b. Computed only for a 2x2 table

Komorbid * lama rawat**Crosstab**

		Lama rawat		Total
		>= 7 hari	< 7 hari	
Komorbid	Count	17	27	44
	% within Komorbid	38,6%	61,4%	100,0%
	Count	6	28	34
	% within Komorbid	17,6%	82,4%	100,0%
	Count	23	55	78
	% within Komorbid	29,5%	70,5%	100,0%
Total				

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4,064 ^a	1	,044		
Continuity Correction ^b	3,117	1	,077		
Likelihood Ratio	4,215	1	,040		
Fisher's Exact Test				,050	,037
Linear-by-Linear Association	4,012	1	,045		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 10,03.

b. Computed only for a 2x2 table

Komorbid * Status of discharge**Crosstab**

		Status of Discharge		Total
		Meninggal	Sembuh	
Komorbid	Count	2	42	44
	% within Komorbid	4,5%	95,5%	100,0%
	Count	3	31	34
	% within Komorbid	8,8%	91,2%	100,0%
	Count	5	73	78
	% within Komorbid	6,4%	93,6%	100,0%
Total				

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	,585 ^a	1	,444		
Continuity Correction ^b	,089	1	,765		
Likelihood Ratio	,580	1	,446		
Fisher's Exact Test				,648	,378
Linear-by-Linear Association	,578	1	,447		
N of Valid Cases	78				

a. 2 cells (50,0%) have expected count less than 5. The minimum expected count is 2,18.

b. Computed only for a 2x2 table

Komorbid * Respon terapi

Crosstab

		Respon terapi		Total
		Tidak	Membaiik	
Komorbid	Ada	Count	17	27
	Ada	% within Komorbid	38,6%	61,4%
	Tidak	Count	11	23
	Tidak	% within Komorbid	32,4%	67,6%
Total		Count	28	50
		% within Komorbid	35,9%	64,1%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2- sided)	Exact Sig. (1- sided)
Pearson Chi-Square	,329 ^a	1	,566		
Continuity Correction ^b	,113	1	,737		
Likelihood Ratio	,330	1	,565		
Fisher's Exact Test				,638	,370
Linear-by-Linear Association	,325	1	,569		
N of Valid Cases	78				

a. 0 cells (0,0%) have expected count less than 5. The minimum expected count is 12,21.

b. Computed only for a 2x2 table

Descriptives

Respon terapi		Statistic	Std. Error
Tidak	Mean	12,11	2,117
	95% Confidence Interval for Mean	Lower Bound 16,45	7,76
	5% Trimmed Mean	10,71	
	Median	8,00	
	Variance	125,507	
	Std. Deviation	11,203	
	Minimum	2	
	Maximum	53	
	Range	51	
	Interquartile Range	13	
Umur	Skewness	2,094	,441
	Kurtosis	5,686	,858
	Mean	12,60	1,679
	95% Confidence Interval for Mean	Lower Bound 15,97	9,23
	5% Trimmed Mean	11,44	
Membuat	Median	7,00	
	Variance	140,980	
	Std. Deviation	11,873	
	Minimum	2	
	Maximum	50	
	Range	48	
	Interquartile Range	15	
	Skewness	1,336	,337
	Kurtosis	1,146	,662

Tests of Normality

	Respon terapi	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Umur	Tidak	,183	28	,017	,787	28	,000
	Membuat	,221	50	,000	,825	50	,000

a. Lilliefors Significance Correction

Descriptives

		Lama rawat		Statistic	Std. Error
Umur	>= 7 hari	Mean		12,87	2,499
		95% Confidence Interval for Mean	Lower Bound	7,69	
			Upper Bound	18,05	
		5% Trimmed Mean		11,43	
		Median		7,00	
		Variance		143,664	
		Std. Deviation		11,986	
	< 7 hari	Minimum		2	
		Maximum		53	
		Range		51	
		Interquartile Range		16	
		Skewness		1,827	,481
		Kurtosis		4,487	,935
		Mean		12,24	1,550
< 7 hari	>= 7 hari	95% Confidence Interval for Mean	Lower Bound	9,13	
			Upper Bound	15,34	
		5% Trimmed Mean		11,07	
		Median		7,00	
		Variance		132,110	
		Std. Deviation		11,494	
		Minimum		2	
	< 7 hari	Maximum		50	
		Range		48	
		Interquartile Range		13	
		Skewness		1,469	,322
		Kurtosis		1,602	,634

Tests of Normality

	Lama rawat	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Umur	>= 7 hari	,223	23	,004	,792	23	,000
	< 7 hari	,195	55	,000	,817	55	,000

a. Lilliefors Significance Correction

Descriptives

		Status of Discharge		Statistic	Std. Error
Umur	Meninggal	Mean		12,60	4,632
		95% Confidence Interval for Mean	Lower Bound	-,26	
		Mean	Upper Bound	25,46	
		5% Trimmed Mean		12,33	
		Median		7,00	
		Variance		107,300	
		Std. Deviation		10,359	
	Sembuh	Minimum		3	
		Maximum		27	
		Range		24	
		Interquartile Range		19	
		Skewness		,766	,913
		Kurtosis		-1,771	2,000
		Mean		12,41	1,370
		95% Confidence Interval for Mean	Lower Bound	9,68	
		Mean	Upper Bound	15,14	
		5% Trimmed Mean		11,11	
		Median		7,00	
		Variance		137,107	
		Std. Deviation		11,709	
		Minimum		2	

Tests of Normality

	Status of Discharge	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Umur	Meninggal	,306	5	,143	,871	5	,269
	Sembuh	,195	73	,000	,814	73	,000

a. Lilliefors Significance Correction

Analisis multivariat

Dependent Variable Encoding

Original Value	Internal Value
< 7 hari	0
=> 7 hari	1

Categorical Variables Codings

		Frequency	Parameter coding	
			(1)	(2)
Derajat pneumonia	Pneumonia	17	1,000	,000
	Berat	52	,000	1,000
Komorbid	Sangat berat	9	,000	,000
	Ada	44	1,000	
Antibiotik	Tidak	34	,000	
	Tunggal	38	1,000	
SaO2	Kombinasi	40	,000	
	< 95	26	1,000	
	=> 95	52	,000	

Block 0: Beginning Block

Classification Table^{a,b}

	Observed	Predicted			Percentage Correct	
		Lama rawat		< 7 hari		
		< 7 hari	=> 7 hari			
Step 0	Lama rawat	< 7 hari	55	0	100,0	
		=> 7 hari	23	0	,0	
Overall Percentage					70,5	

a. Constant is included in the model.

b. The cut value is ,500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)
Step 0 Constant	-,872	,248	12,327	1	,000	,418

Variables not in the Equation

	Score	df	Sig.
Step 0 Variables	SaO2(1)	5,210	,022
	Pneumonia	4,098	,129
	Pneumonia(1)	1,466	,226
	Pneumonia(2)	,031	,861
	Antibiotik(1)	1,200	,273
	Komorbid(1)	4,064	,044
	Overall Statistics	13,632	,018

Block 1: Method = Backward Stepwise (Likelihood Ratio)**Omnibus Tests of Model Coefficients**

		Chi-square	df	Sig.
Step 1	Step	14,052	5	,015
	Block	14,052	5	,015
	Model	14,052	5	,015
Step 2 ^a	Step	-3,661	2	,160
	Block	10,391	3	,016
	Model	10,391	4	,034
Step 3 ^a	Step	-1,989	1	,158
	Block	8,403	2	,015
	Model	8,403	2	,015

a. A negative Chi-squares value indicates that the Chi-squares value has decreased from the previous step.

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	80,555 ^a	,165	,235
2	84,216 ^b	,125	,178
3	86,205 ^b	,102	,145

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than ,001.

b. Estimation terminated at iteration number 4 because parameter estimates changed by less than ,001.

Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	2,503	7	,927
2	,593	6	,997
3	,078	2	,962

Contingency Table for Hosmer and Lemeshow Test

		Lama rawat = < 7 hari		Lama rawat = >= 7 hari		Total
		Observed	Expected	Observed	Expected	
Step 1	1	6	6,557	1	,443	7
	2	9	8,574	1	1,426	10
	3	8	7,587	1	1,413	9
	4	8	8,304	2	1,696	10
	5	7	7,426	3	2,574	10
	6	5	5,259	3	2,741	8
	7	7	5,553	2	3,447	9
	8	2	2,583	3	2,417	5
Step 2	9	3	3,156	7	6,844	10
	1	8	8,248	1	,752	9
	2	14	13,378	2	2,622	16
	3	3	3,172	1	,828	4
	4	13	13,198	4	3,802	17
	5	3	3,202	2	1,798	5
	6	6	6,176	4	3,824	10

	7	5	4,382	3	3,618	8
	8	3	3,244	6	5,756	9
	1	22	21,755	3	3,245	25
Step 3	2	19	19,245	8	7,755	27
	3	6	6,245	3	2,755	9
	4	8	7,755	9	9,245	17

Classification Table^a

	Observed	Predicted			Percentage Correct	
		Lama rawat		< 7 hari		
		< 7 hari	>= 7 hari			
Step 1	Lama rawat	< 7 hari	50	5	90,9	
		>= 7 hari	14	9	39,1	
Overall Percentage					75,6	
Step 2	Lama rawat	< 7 hari	52	3	94,5	
		>= 7 hari	17	6	26,1	
Overall Percentage					74,4	
Step 3	Lama rawat	< 7 hari	47	8	85,5	
		>= 7 hari	14	9	39,1	
Overall Percentage					71,8	

a. The cut value is ,500

Variables in the Equation

	B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I.for EXP(B)		
							Lower	Upper	
Step 1 ^a	SaO2(1)	1,143	,577	3,925	1	,048	3,135	1,012	9,708
	Pneumonia			3,530	2	,171			
	Pneumonia(1)	-1,535	1,019	2,271	1	,132	,215	,029	1,587
	Pneumonia(2)	-1,600	,868	3,400	1	,065	,202	,037	1,106
	Antibiotik(1)	-,937	,585	2,567	1	,109	,392	,124	1,233
	Komorbid(1)	1,143	,604	3,579	1	,059	3,135	,960	10,24
	Constant	-,194	,928	,044	1	,834	,824		4

	SaO2(1)	1,053	,541	3,780	1	,052	2,865	,992	8,280
Step 2 ^a	Antibiotik(1)	-,765	,551	1,930	1	,165	,465	,158	1,369
	Komorbid(1)	1,151	,581	3,916	1	,048	3,160	1,011	9,877
	Constant	-1,630	,531	9,415	1	,002	,196		
	SaO2(1)	1,085	,532	4,155	1	,042	2,958	1,043	8,393
Step 3 ^a	Komorbid(1)	,994	,560	3,146	1	,076	2,702	,901	8,103
	Constant	-1,903	,507	14,071	1	,000	,149		

a. Variable(s) entered on step 1: SaO2, Pneumonia, Antibiotik, Komorbid.

Model if Term Removed



Variable	Model Log Likelihood	Change in -2 Log Likelihood	df	Sig. of the Change
Step 1	SaO2	-42,283	4,011	,045
	Pneumonia	-42,108	3,661	,160
	Antibiotik	-41,630	2,706	,100
	Komorbid	-42,198	3,842	,050
Step 2	SaO2	-44,006	3,796	,051
	Antibiotik	-43,102	1,989	,158
	Komorbid	-44,223	4,230	,040
Step 3	SaO2	-45,196	4,188	,041
	Komorbid	-44,776	3,348	,067

Variables not in the Equation

		Score	df	Sig.
Step 2 ^a	Pneumonia	3,890	2	,143
	Variables	Pneumonia(1)	,087	,768
		Pneumonia(2)	1,306	,253
Step 3 ^b	Overall Statistics	3,890	2	,143
	Pneumonia	3,159	2	,206
	Variables	Pneumonia(1)	,175	,676
		Pneumonia(2)	,903	,342
	Antibiotik(1)	1,966	1	,161
Overall Statistics		5,432	3	,143

a. Variable(s) removed on step 2: Pneumonia.

b. Variable(s) removed on step 3: Antibiotik.

Lampiran 5. Dokumentasi

Lampiran 6. Biodata Mahasiswa**Identitas**

Nama : Aldora Putri Tammy
NIM : 22010112130131
Tempat/tanggal lahir : Padang, 20 April 1994
Jenis kelamin : Perempuan
Alamat : Jl. Bulusan Utara Raya No.01 , Tembalang, Semarang
Nomor Telepon : -
Nomor HP : 082226511913
e-mail : tammy.al doraputri@yahoo.com

Riwayat Pendidikan Formal

- | | | | |
|--------|------------------------------|-------------|--------|
| 1. SD | : SDN 06 Batusangkar | Lulus tahun | : 2006 |
| 2. SMP | : SMPN 1 Pariaman | Lulus tahun | : 2009 |
| 3. SMA | : SMAN 1 Padang Panjang | Lulus tahun | : 2012 |
| 4. S1 | : Pendidikan Dokter FK UNDIP | Masuk tahun | : 2012 |