

Shrimp Paste “Terasi” Volatile Compounds From Northern Coast of Central Java

by Agus Trianto

Submission date: 29-Aug-2018 03:36PM (UTC+0700)

Submission ID: 994475895

File name: C7-Shrimp_Paste_Terasi_Volatile_Compounds.pdf (356.77K)

Word count: 3061

Character count: 17114

SHRIMP PASTE “TERASI” VOLATILE COMPOUNDS FROM NORTHERN COAST OF CENTRAL JAVA

Retno Murwani^{a*}, Hera Surya Adhi Putra^b, Henggar Widiyanto^b, Agus Trianto^{a,b}, Ambariyanto^{a,b}

^aNatural Product Laboratory, Centre of Research and Services Diponegoro University, Indonesia

^bDepartment Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia

10
Article history

Received

11 November 2015

Received in revised form


18 December 2015

Accepted

23 February 2016

*Corresponding author
rmurwani@gmail.com

Graphical abstract



Abstract

A study was conducted to identify the composition of volatile compounds from traditional fermented shrimp called 'terasi'. Terasi samples were collected from six regions of northern coast of Central Java, Indonesia namely Pekalongan, Batang, Kendal, Demak, Jepara, and Pati. Mass spectral interpretation showed that terasi from these regions could be identified to contain a total of 102 volatile compounds. Terasi from Pekalongan, Batang, Kendal, Demak, Jepara, and Pati, each contained nine, 21, 10, 29, 12, and 21 volatile compounds respectively. There were four similar volatile compounds from Demak, Jepara, and Pati samples, and two distinctive off odor in all six regions.

Keywords: Fermented shrimp, northern coast of Central Java, terasi, volatile compounds

© 2016 Penerbit UTM Press. All rights reserved

1.0 INTRODUCTION

Fermented shrimp or shrimp paste well known as terasi is one of traditional processing of fishery products in Indonesia. Terasi is made from fermented shrimp with or without salt addition. However fish or a mixture of shrimp and fish can also be used depending on the availability or season. Terasi has reddish brown to blackish brown color depending on the raw materials and the naturally occurring microbes which involve during fermentation. Fresh terasi has a strong distinctive smell or odor which upon heating or cooking terasi releases volatile compounds. Exposure to such food odor produces appetizing response which can promote food intake [1, 2]. Terasi has been used for centuries as a flavor enhancer in many kinds of dishes and the released odor produce appetite for consuming the associated terasi flavored foods which can lead to satiety. Therefore terasi has been playing significant role in ensuring adequate food intake for centuries. Terasi from different production regions have similar but not the same odor as a result of processing

variety such as length of sun exposure for drying and of fermentation time. It is the aim of this study to identify terasi volatile compounds produced at the centre of traditional terasi production in northern coast of Central Java i.e. Pekalongan, Batang, Kendal, Demak, Jepara, and Pati.

2.0 EXPERIMENTAL

Terasi were collected from six regions in the northern coast of Central Java namely Demak Jepara, Pati (Juwana), Pekalongan, Batang and Kendal. Terasi was collected from traditional small scale producers. Extraction of volatile compounds was carried out by mixing terasi sample with distilled water (1:5) and placed in Lickens-Nickerson apparatus. An internal standard of 0.1 mL was added. The sample was heated in a boiling water bath for 2 h after which it was extracted with 50 mL of diethyl ether at 40 °C. After extraction the remaining water was eliminated by addition of sodium sulphate anhydrous [3].

The extract was concentrated using Vigreux column to obtain 1 mL volume [4]. Finally it was further concentrated by nitrogen gas to obtain a final volume of 0.1 mL to 0.2 mL and 0.1 μ L was injected into GC-MS (Hewlett Packard). GC-MS was equipped with an HP-Wax column of 30 mm long, 0.25 mm diameter, and 2.56 mm thickness. Helium as the gas carrier was flowed at 1 mm min⁻¹ at 3 bar (1 bar = 100 000 Pa) The interface and injection temperature was 230 °C. Temperature program was set at initial temperature of 70 °C for 1 min and 10 °C per min increase until the final temperature of 230 °C was achieved. The electron energy of mass spectrometry was 1.2 kV, Mr range of 33 to 400, and 0.5 second interval [5]. Mass Spectral Interpretation was done with the aid of a computer program to compare the patterns of mass spectra of compounds with a pattern of mass spectra in the National Institute of Standards and Technology mass spectra library collection. The collection has more than 62 000 patterns of mass spectra [6].

3.0 RESULTS AND DISCUSSION

The chromatogram of GC-MS results were identified based on the retention time and the area of the peaks. Mass spectral interpretation of the peaks were identified by comparing them to pattern of mass spectra in the NIST mass spectra library collection. Table 1 showed the results of each class of volatile compounds in each sampling location. Terasi from Demak, Jepara, Pati, Pekalongan, Batang and Kendal

could be identified to contain a total of 102 volatile compounds. The identified volatiles can be grouped into hydrocarbon, carbonyls, sulfur, alcohols, nitrogens, fatty acid, and ester compounds which showed similarities with others studying volatiles chemicals of shrimp or fish paste [7-9]

Terasi from Demak, Jepara, Pati, Pekalongan, Batang and Kendal each contained 29, 12, 21, nine, 21, and 10 volatile compounds, respectively. Terasi from Demak showed 10 hydrocarbon, four carbonyls, one sulphur, seven alcohols, five nitrogens, a fatty acid, and an ester. Terasi from Jepara showed four hydrocarbon, a sulphur, two alcohols, two nitrogen, two fatty acids, and an ester compounds. Terasi from Pati showed seven hydrocarbons, a carbonyl, two sulphurs, four alcohols, five nitrogens, and a fatty acid compounds. Terasi from Pekalongan showed four hydrocarbon, one carbonyls, three alcohols, and a nitrogen compounds.

Terasi from Batang showed four hydrocarbon, three carbonyls, two sulfur, five alcohols, five nitrogen, a fatty acid, and an ester compounds. Terasi from Kendal showed two hydrocarbon, a carbonyl, three alcohols, three nitrogen, and a fatty acid compounds. Terasi from Demak, Jepara, and Pati showed four same volatile compounds i.e hexadecane, 1,11-dodecadiene, dimethyl trisulfide, and indole, while Terasi from Pekalongan, Batang and Kendal showed three same compounds i.e Hexadecane, phenols, and indoles. The hexadecane and indole which give terasi off odor consistently found in samples from the six regions.



Figure 1 Map of the the six regions where terasi samples were collected (●)

Table 1 Volatile compounds of terasi samples from Demak, Jepara, Pati, Pekalongan, Batang, and Kendal

Volatile compounds detected and identified by GC-MS	Terasi from six regions of northern coast of Central Java					
	Demak	Jepara	Pati	Pekalongan	Batang	Kendal
Hydrocarbons						
1 Hexadecane	+	+	+	+	+	+
2 1,4-Octadiene	-	-	-	+	-	-
3 7-hexadecane	-	+	+	+	+	-
4 1,11-Dodecadiene	+	+	+	+	-	-
5 2-Methylbutylidene 2-phenylethyl	-	-	+	-	+	+
6 1-Docen-3-yne	-	-	-	-	+	-
7 Cyclododecane	-	+	-	-	-	-
8 3-Tetradecene	+	-	+	-	-	-
9 1.3-cyclooctadecene	+	-	-	-	-	-
10 1-Pentadecene	+	-	+	-	-	-
11 Cyclotetradecene	+	-	-	-	-	-
12 Cyclododecene	+	-	-	-	-	-
13 1,2-dimethyl cyclooctane	+	-	-	-	-	-
14 Bicyclo [2.2.1] heptane	+	-	-	-	-	-
15 1.4-cyclononadiene	-	-	+	-	-	-
16 2- methoxy naphthalene	+	-	-	-	-	-
SubTotal	10	4	7	4	4	2
Carbonyl						
17 5z, 8z, 11z-tetra-decatrien -2-one	+	-	-	+	+	-
18 Butyl hidroxytoluene	-	-	-	-	+	+
19 2-Hydroxy-4, 6-Cyclooctadien-1-one	-	-	-	-	+	-
20 2-Undecanon	+	-	-	-	-	-
21 2-Tridecanon	+	-	-	-	-	-
22 1-Finil-Etenon	+	-	-	-	-	-
23 Benzaldehyde	-	-	+	-	-	-
24 3,4,5-methyl Cyclopentene -1-on	-	-	+	-	-	-
SubTotal	4		1	1	3	1
Sulphur						
25 Dimethyl trisulfide	+	+	+	-	+	-
26 Ethane, 1,2-bis (Methylthio)	-	-	-	-	+	-
27 Dinonil trisulfide, isomer mix	-	-	+	-	-	-
SubTotal	1	1	2	0	2	0
15 Alcohol						
28 Phenol, 2,6-bis (1,1-dimethylethyl)	+	+	+	+	+	+
29 Phenol	+	+	+	+	+	+
30 1-Pentadecanol	-	-	-	+	-	-
31 3-Heptanol, 3,5-dimethyl	-	-	-	-	+	-
32 1,3-Benzenediol, 5-methyl	-	-	-	-	+	-
33 1-Tetradecanol	-	-	-	-	+	-
34 11-Hexadecen-1-ol	-	-	-	-	+	-
35 Eugenol	-	-	-	-	-	+
36 1-Hexadecanol	-	-	-	-	-	+
37 Phenyl ethyl alcohol	-	+	-	-	-	-
38 2.2-oxibis Ethanol	-	+	-	-	-	-
39 4-methyl phenol	+	-	+	-	-	-
40 1-Heptadecanol	+	-	-	-	-	-
41 14ctadecanol	+	-	-	-	-	-
42 p-Mentha-1 (7), 8 (10)-dien-9-ol	+	-	-	-	-	-
43 9,12,15-octadeca-trien-1-ol	+	-	-	-	-	-
44 Mentional	-	-	+	-	-	-
SubTotal	7	2	4	3	5	3
Nitrogen						
45 Indole	+	+	+	+	+	+
46 1-Phenyl-2-methylenaminopropane	-	-	-	-	+	-
47 2,5-dimethyl pyrazine	-	-	-	-	+	-
48 2 - (2-hydroxyethyl) piperidine	-	-	-	-	+	-

(continued on the next page)

Table 1 continued

Volatile compounds detected and identified by GC - MS	Terasi from six regions of northern coast of Central Java					
	Demak	Jepara	Pati	Pekalongan	Batang	Kendal
49 2-Phenyl-N-benzylidene ethylamine	-	-	-	-	+	-
50 3-Phenyl pyridine	-	-	-	-	-	+
51 N-(2-Feniletil), Asetamide	-	-	-	-	-	+
52 2,6-dimethyl-3-N-butilpyrazine	+	-	-	-	-	-
53 2,6-dimethyl pirazin	+	-	-	-	-	-
54 2,5-dimethyl-3-(3-methyl) Pirazin	-	-	+	-	-	-
55 4-Phenyl Piridine	+	-	-	-	-	-
56 4-methyl-2-phenyl pyrimidine	+	-	-	-	-	-
57 Benzeamine	-	-	+	-	-	-
58 1-Benzeneetamine, N-(Fenilmetil)	-	-	+	-	-	-
59 5-metiloksiindol	-	-	+	-	-	-
SubTotal	5	2	5	1	5	3
Fatty acid						
60 1,2 Benzene-dicarboxylic acid	-	+	-	-	+	-
61 Butanoic acid, 3-methyl, 2-phenyl	-	-	-	-	-	+
62 Pentanoic acid	-	+	-	-	-	-
63 Hexadecanoic acid	+	-	-	-	-	-
64 Tetradecanoic acid	-	-	+	-	-	-
SubTotal	1	2	1	-	1	1
Ester						
65 Hexadecanoic acid, Methyl ester	-	+	-	-	-	-
66 Ethyl linoleic	+	-	-	-	-	-
67 Ethyl laurate	-	-	-	-	+	-
SubTotal	1	1	0	0	1	0
Total	29	12	21	9	21	10




Figure 2 Chromatogram peak area (%) of each class of terasi volatile compounds in each sampling regions

Figure 2 shows that the largest total peak area of hexadecane was found in sample from Pekalongan. Hexadecane was formed during fermentation in which lipids were oxidized into fatty acids by microbial activities. Fatty acids were subsequently converted into other compounds one of which is a

hydrocarbon The highest total percentage area of carbonyl compounds was found in samples from Kendal (23.45 %). Carbonyl compounds were also derived from lipid oxidation by microbial activities which produced distinctive scent of fermented shrimp. Carbonyl compounds could also be

produced by non-enzymatic browning due to astaxanthin, the red pigment found in shrimp as the raw material. The pigments changed to brown color upon heating such as sun drying. Non-enzymatic browning enhanced the aroma and taste of a product. In samples from Jepara no carbonyl compounds were detected.

Sulphur compounds in shrimp paste was produced by enzymatic or heat protein degradation. The same sulfur compounds detected from samples of Demak, Jepara, Pati and Batang were dimethyl trisulphide, while none was detected in samples from Pekalongan and Kendal. In general, sulphur compounds of shrimp paste was very small, however they are quite strong and important. This was due to small threshold of sulphur compounds compared to the threshold of other volatile compounds [10]. The presence of sulfur compounds in shrimp paste gave the distinctive strong aroma of the product and therefore the strongest aroma of Terasi was found in samples from Pati.

Alcohols identified from Pekalongan terasi samples had the greatest percentage peak area compared to five other locations. Phenol were the same alcohols found in five locations. Phenol could be produced by lipids oxidation, sugar, and protein degradation. Protein degradation produced amino acids and peptides which were further broken down into other components such as hydrocarbons, esters and alcohols [10].

Indole as nitrogen compound identified in terasi samples from the six locations is an interesting volatile compound. It is interesting as it also found in jasmine and pork liver. Indoles in high concentration is well known as a typical off odor associated with putrefaction or fecal contamination, however no *E. coli* was found in all samples [11]. In contrast, when indole was combined in a very low concentration with other flowery chemical odor such as jasmine it would give depth to the existing aromas [10]. Nitrogen compounds was produced from the breakdown product of protein by the action of proteolytic bacteria during fermentation or the reaction of ammonia or amino acids with reducing sugars [9,11]. The highest total percentage area of nitrogen compounds was found from Batang and the smallest from Pekalongan. So samples from Batang would have stronger odor compared to samples from five other regions.

Fatty acid compounds were not found from Pekalongan samples. The largest total percentage area of fatty acid compounds were found from Jepara samples. Fatty acid compounds were produced from microbial enzymatic degradation of lipid [12]. The volatile fatty acids often lead to acidity in the aroma of terasi. Therefore terasi from Jepara has the most acidic odor compared with the other five regions.

Ester compounds in Terasi samples from Pati, Pekalongan, and Kendal were not found. The largest total percentage area of ester compounds in Terasi was found from Batang and the smallest one was

found from Demak. Ester compounds was produced by the reaction between alcohol and a carboxylic acid salt which was catalyzed by esterase released by microbial fermentation.

The processing steps in traditional terasi making was in principle the same. Shrimp was washed and sun dried for a day or more depending on the sun light intensity. The dry shrimps were pounded manually and salt could be added, mixed, and dried further. In the afternoon when the sun has already set, the dried shrimp was kept indoor. The next day, it was pounded again and sun dried. After alternate sun drying and pounding, the product was molded and wrapped in banana leaves or plastic and kept for several days or more or even up to 4 wk to let fermentation takes place. The final product was named terasi or known as shrimp paste. The product from each region showed variation on the number and peak area of volatile compounds. The three regions were located at the eastern part of Central Java and 3 others at western part (Figure 1). The locations have different ecological setting so that shrimps catch in those regions could have different characteristics. This combines with different processing namely length of sun drying, quantity of added salts, and length of fermentation time would give different fermentation process which consequently produces different microbial and enzymatic activities. These together would produce different degree of degradation products which contributed to different degree of volatile compounds typical of each region. We had record of processing procedure of terasi samples from three regions i.e Pekalongan, Batang, Kendal. The processing of sample from Pekalongan, Batang and Kendal were basically the same as described previously. However, terasi made in Pekalongan was added with salt and food color, Batang without added salt, and Kendal was also added with salt but no color. Together with the previous results which showed that terasi from Batang had higher number of total volatile compounds (21) compared to Pekalongan (9) and Kendal (10), this was more likely related to the processing without added salt. Shrimp paste processing (fermentation) with no salt produced a higher number of volatile compounds compared to processing with salt addition. This possibility was corroborated by previously developed study regarding proteolytic bacteria [10]. It was found that maximal growth of mesophilic proteolytic bacteria from terasi was found at zero (0) concentration of salt.

Terasi either fresh or briefly heated before or after mixing with other ingredients can release their volatiles which give an appetizing effect from the aromas. The sense of smell or termed olfactory sensation occurs when odorant molecules bind to specific scent receptors in our nasal cavity. The receptors can detect thousands of volatile chemicals. Volatile chemical odorants found in this study may all bind to the scent receptors simultaneously and send olfactory sensations which

produced appetite for the associated food intake. Food intake stimulates not only retronasal olfactory sensation but also the taste buds receptors found in mouth and gastro-intestine which ultimately send signals to brain and produce satiety [13]. For centuries, terasi has been used in various dishes and cooking, and most often mixed with chili, garlic, a pinch of salt, and palm sugar to yield a well known dish called Sambal. Sambal is a appetizing food which accompanying rice and other complementary food such as Tempe which therefore usually increase food intake and produce satiety.

4.0 CONCLUSION

Volatiles compounds of terasi from six regions of northern coast of Central Java belongs to hydrocarbons, carbonyls, sulphur, alcohols, nitrogen, fatty acids, and esters. The combined number, types, and concentration of the volatile compounds are different from each region and produced distinctive-signature smell of the product.

Acknowledgement

This study was financially supported by Centre of Ethno Food and Medicine, Research Institute Diponegoro University for financial support. Technical assistance from Anantasia Indriani is deeply appreciated.

References

- [1] Ramaekers, M. G., S. Boesveldt, G. T. Gort, C. M. M. Lakemond, M. A. J. S. van Boekel, and P. A. Luning. 2014. Sensory-Specific Appetite is Affected by Actively Smelled Food Odor and Remains Stable Over Time in Normal-Weight Women. *The Journal of Nutrition*. 144(8): 1314-1319.
- [2] Ramaekers, M. G., S. Boesveldt, C. M. M. Lakemond, M. A. J. S. van Boekeland, and P. A. Luning. 2013. Odor: Appetizing or Satiating? Development of Appetite During Odor Exposure Over Time. *International Journal of Obesity*. 38: 650-656.
- [3] Kumara, B. 1999. Identifikasi "Character Impact Compound" Flavor Buah Kawista - (Feronia limona) [Identification of Character Impact Fruit Flavor Kawista Compound (Feronia limona)]. Thesis. Bogor: IPB. [Bahasa Indonesia].
- [4] Larson, M. and L. Pool. 1990. Quick and Sample Extraction Method for Aroma Compound in Food Products. In Bessier, Y. and A. F. Thomas (eds). *Flavour Science and Technology*. Chicester: John Wiley and Sons.
- [5] Fardiaz, D. 1989. *Kromatografi Gas dalam Analisa Pangan* [Gas Chromatography in Food Analysis]. Bogor: IPB. [Bahasa Indonesia].
- [6] NIST. 2002. National Institute of Standards and Technology Standard Reference Database 1A NIST/EPA/NIH Mass Spectral Library and NIST Mass Spectral Search Program.
- [7] Cha, Y. J. and K. R. Cadwallader. 1995. Volatile Compounds in Salt-fermented Fish and Shrimp Pastes. *Journal of Food Science*. 60(1): 19-27.
- [8] Li, C., J. Wu, Y. Yan Li, and Z. Dai. 2013. Identification of the Aroma Compounds in Slinky Mandarin Fish (*Siniperca chuatsi*) and Comparison of Volatiles During Fermentation and Storage. *International Journal of Food Science & Technology*. 48(11): 2429-2437.
- [9] Pongsetkul, J., S. Benjakul, P. Sampavapol, K. Osako, and S. Faithong. 2015. Chemical Compositions, Sensory and Antioxidative Properties of Salted Shrimp Paste (Ka-pi) in Thailand. *International Food Research Journal*. 22(4): 1454-1465.
- [10] Choi, N. E. and J. H. Han. 2015. *How Flavor Works. The Science of Taste and Aroma*. UK: John Wiley and Sons, Ltd.
- [11] Murwani, R., Supriyadi, A. Subagio, Trianto, and Ambariyanto. 2015. Isolation and Identification of Thermophilic and Mesophylic Proteolytic Bacteria from Terasi. *AIP Proceeding 2nd International Conference on Chemical and Material Engineering*. 1699. (In Press).
- [12] Surono, I. S. and A. Hosono. 1994. Chemical and Aerobic Bacterial Composition of "Terasi", a Traditional Fermented Product from Indonesia. *Journal of the Food Hygienic Society of Japan*. 35(3): 299-304.
- [13] Moyer, M. 2015. What Makes Food Taste So Good? *Scientific American*. 24(2): 4-8.

Shrimp Paste "Terasi" Volatile Compounds From Northern Coast of Central Java

ORIGINALITY REPORT

11%	%	11%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

- 1** Scott C. Hutchings, Katy M. Horner, Victoria A. Dible, John M.V. Grigor, Dolores O'Riordan. "Modification of aftertaste with a menthol mouthwash reduces food wanting, liking, and ad libitum intake of potato crisps", *Appetite*, 2017
Publication 2%
- 2** I Samidjan, D Rachmawati. " Polyculture Engineering technology of red tilapia () and white shrimp () based for protease enzyme ", *IOP Conference Series: Earth and Environmental Science*, 2018
Publication 1%
- 3** Chaofan Ji, Jingbo Zhang, Xinping Lin, Jing Han, Xiuping Dong, Song Yang, Xiaoming Yan, Beiwei Zhu. "Metaproteomic analysis of microbiota in the fermented fish, *Siniperca chuatsi*", *LWT*, 2017
Publication 1%
- 4** Yanesti N Lestari, Retno Murwani, Tri W Agustini. " Flavor Enhancer From Catfish () 1%

Bekasam Powder and Angiotensin-I-Converting Enzyme (ACE) Inhibitory Activity in Various Dishes ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

5

"Microbiology of Fermented Foods", Springer Nature America, Inc, 1997

Publication

6

Raghavan, Sivakumar, and Hordur G. Kristinsson. "Influence of processing on lipids and lipid oxidation in aquatic foods", Antioxidants and Functional Components in Aquatic Foods, 2014.

Publication

7

David Arroyo, M. Cruz Ortiz, Luis A. Sarabia. "Optimization of the derivatization reaction and the solid-phase microextraction conditions using a D-optimal design and three-way calibration in the determination of non-steroidal anti-inflammatory drugs in bovine milk by gas chromatography–mass spectrometry", Journal of Chromatography A, 2011

Publication

8

Jaksuma Pongsetkul, Soottawat Benjakul, Kitiya Vongkamjan, Punnanee Sumpavapol, Kazufumi Osako, Nandhsha Faithong. "Changes in volatile compounds, ATP-related compounds

1%

1%

1%

1%

and antioxidative properties of Kapi , produced from *Acetes vulgaris* , during processing and fermentation", Food Bioscience, 2017

Publication

9

Henry, Theodore B. Menn, Fu-Min Fleming,. "Attributing effects of aqueous [c.sub.60] nano-aggregates to tetrahydrofuran decomposition products ", Environmental Health Perspectives, July 2007 Issue

1%

Publication

10

Zhang, Jun Simate, Geoffrey S. Lee, Sang. "Relating asphalt binder elastic recovery properties to HMA crack modeling and fatigue life predictio", Construction and Building Materials, May 15 2016 Issue

<1%

Publication

11

Agus Trianto, Agus Sabdono, Baskoro Rochaddi, Desy Wulan Triningsih, Dewi Seswita Zilda. "Identification Sponges-Associated Fungi From Karimunjawa National Park", IOP Conference Series: Earth and Environmental Science, 2018

<1%

Publication

12

Moyer, Michael. "The Food Issue", Scientific American, 2013.

<1%

Publication

13

Little, P.J.. "Tissue disposition of [³H]sarin and

its metabolites in mice", Toxicology and Applied Pharmacology, 198605

Publication

<1%

14

Omolo, M.O. Okinyo, D. Ndiege, I.O. Lwan. "Fumigant toxicity of the essential oils of some African plants against Anopheles gambiae sensu stric", Phytomedicine: International Journal of , March 2005 Issue

Publication

<1%

15

Cros, S.. "Desalination of mussel cooking juices by electrodialysis: effect on the aroma profile", Journal of Food Engineering, 200508

Publication

<1%

16

Wang, H.. "Persistent organic pollutants in water and surface sediments of Taihu Lake, China and risk assessment", Chemosphere, 200301

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off