

Total Sugar Content, Protein, Adhesiveness, and Hardness of Fermented

by Siti Susanti

Submission date: 11-Nov-2018 10:52AM (UTC+0700)

Submission ID: 1036655318

File name: ar_Content,_Protein,_Adhesiveness,_and_Hardness_of_Fermented.pdf (171.39K)

Word count: 1834

Character count: 10430



Total Sugar Content, Protein, Adhesiveness, and Hardness of Fermented Jackfruit Seeds Flour by *Kluyveromyces marxianus* and *Saccharomyces cerevisiae*

Fathiyatul Islamiyah¹, Siti Susanti¹, Ahmad Ni'matullah Al-Baarri^{1,2*}

¹Food Technology Department, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

²Laboratory of Food Technology, UPT Integrated Laboratory, Diponegoro University, Semarang, Indonesia

*Corresponding author (albari@undip.ac.id)

Abstract

This research aims to produce dough from fermented jackfruit seed using *Kluyveromyces marxianus* and *Saccharomyces cerevisiae*. Concentration of inoculum at 0, 4, 8 and 12% (v/v) was used to measure total sugar, protein content, adhesiveness and hardness in the dough. To produce dough, fermentation process for 1 hour at room temperature (27°C±1) was used. Total sugar was measured using refractometer and protein content was analyzed using Bradford method. Adhesiveness and hardness was analyzed using texture analyzer. Result shows no significant differences (P>0.05) on sugar content in obtained doughs using *K. marxianus* and *S. cerevisiae* resulting a value of 8.442±0.69 and 8.485±0.76°Brix, respectively. Protein content of the fermented dough was 3.51±1.16 and 4.4±2.09% for *K. marxianus* and *S. cerevisiae*, respectively, resulting the significant differences among treatments. The adhesiveness and hardness of fermented dough with *K. marxianus* was significantly higher than that of *S. cerevisiae*. This result may provide beneficial information to food industry that uses jackfruit seed as ingredient.

Article information:

Received 16 November 2016

Accepted 29 December 2016

Available online 31 December 2016

Keywords:

Kluyveromyces marxianus
Saccharomyces cerevisiae
jackfruit seeds flour
fermentation
dough

© 2016

Indonesian Food Technologists
All rights reserved

This is an open access article
under the CC BY-NC-ND license

doi: 10.17728/jaft.19

Introduction

Jackfruit (*Artocarpus heterophyllus* Lam.) consists of about 10-15% seed of the total weight of fruit. This seed contains a rich of carbohydrate and protein (Kumar et al., 1988). Seeds are normally discarded or steamed and eaten as a snack or used in some local dishes for example: "Gudeg"; one of Javanese dishes. In addition, the flour from this seed may provide potential application in bakery product since this seed is a rich source of protein, starch, and dietary fiber (Chowdhury et al., 2012) and potentially developed through fermentation process (Al-Baarri et al., 2013; Yamaoka et al., 2014). Therefore, this research was done to analyze total sugar, protein content, adhesiveness and hardness of dough made of jackfruit seed flour using fermentation process by inoculation of *K. marxianus* and *S. cerevisiae*. This research may provide beneficial information to food industry to open the ways for utilizing jackfruit seed as ingredient.

Materials and Methods

Materials

Jackfruit seeds was obtained from local market, *S. cerevisiae* (FNCC 3022) and *K. marxianus* (FNCC 3026) were obtained from Integrated Research and Testing Laboratory (LPPT) Gadjah Mada University, Yogyakarta, Indonesia. The chemicals used in this research were

analytical reagent grade.

Method

The study was conducted from August–October, 2016. The peeled and sliced jackfruit seeds (±3cm of thickness) was dried at 50°C for 6 h (Jading et al, 2011). Miller was used to convert the dried seed into flour. Hand-held refractometer was used to determine total sugar (Ihsan and Wahyudi, 2010), Bradford method was used to analyze protein content (Purwanto, 2014). Adhesiveness and hardness were analyzed using CT3 Texture Analyzer (Brookfield, 2014).

Fermentation of jackfruit seed flour was conducted using the following description. Fifty percent concentration of flour in the water was inoculated with 0, 4, 8, and 12% of *S. cerevisiae* and *K. marxianus*. Incubation was done for 1 hour at room temperature (27±1°C) in anaerobic condition.

The obtained data were statistically analyzed using one-way ANOVA and Duncan's Multiple Range Test using SPSS 16.0. The level significance was set at $\alpha = 0.05$.

Results and Discussion

Total sugar content in the dough that was composed from *K. marxianus* and *S. cerevisiae* was 8.442±0.69% and 8.485±0.76%, respectively (Figure 1a)

resulting in no significantly differences on b₂₄ value. Sugar content found in this research was 54% lower than that of reported by Fairus et al., (2010). This may be explained by the differences on the water activity in seeds. Furthermore, the enzyme was also plays important role in sugar degradation (Lubis, 2012)




Figure 1. Sugar content (a), total protein (b), adhesiveness (c), and hardness (d) of dough made of jackfruit seed using fermentation process by 0, 4, 8 and 12% of *K. marxianus* and *S. cerevisiae* for 1 hour at room temperature in anaerobic conditions. The superscript shows significantly difference using Duncan's Multiple Range Test.

It was understood that *K. marxianus* break the inulin into oligo-fructose (Yuliana et al., 2014). In addition, *S. cerevisiae* environmental dependently converted complex sugar into glucose and fructose (Awwalurrizki and Putra, 2009; Hasanah and Putra, 2010).

Protein content of dough made of *K. marxianus*

showed significant differences while *S. cerevisiae* showed no significant differences among concentrations of inoculum (Figure 1b). The highest protein content was in the fermented dough with 8% concentration of *S. cerevisiae* i.e. $5.37 \pm 3.22\%$. This value could be categorized as low since originally protein content in seed was low (Mukprasirt and Sajjanantakul, 2004). The reduction of protein content could be explained by proteolytic activity (Foukis et al., 2012) by protease enzyme, endopeptidases, and exopeptidases (Lopez et al., 2011). In order to increase the protein content, longer time of fermentation is needed (Kuswardani and Wijajaseputra, 1998), however, auto-biodegradation may reduce the protein content.

Adhesiveness of dough can be seen on Figure 1c. Fermented dough showed lowest value; i.e. 0.08 ± 0.01 N/mm and highest value; i.e. 0.18 ± 0.08 N/mm at 8% inoculation. Yusriyah and Agustini (2014) reported that the adhesiveness was inoculum concentration and fermentation rate dependent. This research shows the linear correlation between inoculum concentration and adhesiveness resulting the in line result with Rotcha et al. (2015). The low in adhesiveness of dough inoculated with *S. cerevisiae* may be assumed no strong relation between sugar and water in *S. cerevisiae* (Kumura et al., 2004)

Hardness of fermented dough by *K. marxianus* and *S. cerevisiae* can be seen on Figure 1d. The highest hardness was shown at 4% *S. cerevisiae* inoculation, i.e. 18.27 ± 2.41 g/cm². The value of hardness showed linear correlation with concentration of inoculum. This may be assumed by the content of gluten (Warechowska et al., 2016). Furthermore, the inoculum concentration affected the alcohol production resulting the change in hardness dough (Unika, 2015; Rocha et al., 2015).

Conclusion

K. marxianus and *S. cerevisiae* specifically affected protein content, adhesiveness, and hardness while the both of inoculum showed negligible differences on sugar content.

Acknowledgment

Authors would like to thank to Nurul Yaqin, STP for his support to finish the protein content analysis.

References

- Al-Baarri, A.N., Legowo, A.M., Fawaid, M.T. 2013. Alcohol profile from fermented whey using bagase. Journal Aplikasi Teknologi Pangan 2(1), 48-51. (In Bahasa Indonesia).
- Awwalurrizki, N., Putra, S.R., 2009. Sucrose hydrolysis using invertase enzyme to produce ethanol using *zymomonas mobilis*. Bachelor Thesis. Institut Teknologi Sepuluh Nopember. Surabaya. (In Bahasa Indonesia).
- Brookfield, 2014. Texture analyzer operating instruction. Brookfield Engineering Laboratories, USA.
- Chowdhury, A.R., Bhattacharyya, A.K., Chattopadhyay, P., 2012. Study on functional properties of raw and blended jackfruit seed flour (a non-conventional source) for food application. Indian Journal of

- Natural Product and Resources 3(3), 347–353
- Fairus, S., Haryono, Mirantheni, A., Aprianto, A., 2010. The Effect of HCl concentration and hydrolysis duration on the production of glucose from jackfruit seed. Proceeding of National Seminar Chemistry Engineering "Kejuangan", 1-6. (In Bahasa Indonesia).
- Foukis, A., Stergiou, P.Y., Theodorou, L.G., Papagianni, M., Papamichael, E.M., 2012. Purification, kinetic characterization and properties of a novel thermotolerant extracellular protease from *Kluyveromyces marxianus* IFO 0288 with potential biotechnological interest. Journal of Biosource Technology 123, 214-220. Doi: 10.1016/j.biortech.2012.06.090.
- Hasanah, E.N.I., Putra, S.R., 2010. Fiber characteristic of crude invertase enzyme derived from nalginat. Bachelor Thesis. Institut Teknologi Sepuluh Nopember. Surabaya.
- Ihsan, F., Wahyudi, A., 2010. Technical analysis to determine sucrose on papaya. Buletin Teknik Pertanian 15(1),10-12.
- Jading, A., Tethool, E., Payung P., Gultom, S., 2011. Sago starch physicochemical characteristics drying in fluidized by cross flow bed. Jurnal Reaktor 13(3),155-164. (In Bahasa Indonesia)
- Kumar, S., Singh, A.B., Abidi, A.B., Upadhyay, R.G., Singh, A., 1988. Proximate composition of jack fruit seeds. Journal Food Science Technology 25, 308-309.
- Kumura, H., Tanoue, Y., Tsukahara, M., Tanaka, T., Shimazaki, K., 2004. Screening of dairy yeast strains for probiotic applications. Journal of Dairy Science 87(12), 4050-4056. Doi: 10.3168/jds.S0022-0302(04)73546-8
- Kuswardani, I., Wijajaseputra, A.I., 1998. Production single cell protein *Phanerochaete chrysosporium* on waste product from tofu. Proceeding National Seminar "Teknologi Pangan dan Gizi", 604-613.
- Lopez, C.C., Paparella, A., Tofalo, R., Suzzi, G., 2011. Proteolytic activity of *Saccharomyces cerevisiae* strains associated with italian dry-fermented sausage in a model system. International Journal of Food Microbiology 150, 50-58. Doi: 10.1016/j.ijfoodmicro.2011.07.017.
- Lubis, M.R., 2012. Starch hydrolysis using H₂SO₄ as catalyst to produce glue. Jurnal Rekayasa Kimia dan Lingkungan 9(2)62-67.
- Mukprasirt, A., Sajjaanantakul, K., 2004. Physico-chemical properties of flour and starch from jackfruit seeds (*Artocarpus heterophyllus* Lamk.) compared with modified starches. International Journal of Food Science & Technology 39(3), 271-276. Doi: 10.1111/j.1365-2621.2004.00781.
- Purwanto, M.G.M., 2014. Comparison analysis of soluble protein using various methods by UV-visible spectrophotometry. Jurnal Ilmiah Sains dan Teknologi 7(2), 64-71 (In Bahasa Indonesia).
- Rotcha, K.A.P., Vera, N.G., Nicanor, A.B., Cruz, L.G., Uribe, J.P.H., Sanchez, A.T., 2015. Fortification of white bread with guava seed protein isolate. Pakistan Journal of Nutrition 14(11), 828-833.
- Unika, A. 2015. The effect of yeast quantity and fermentation duration on organoleptic characteristic fermented product from banana "pisang tanduk". E-journal Boga 4(1), 192-201.
- Warechowska, M., Markowska, A., Warechowski, J., Mis, A., Nawrocka, A., 2016. Effect of tempering moisture of wheat on grinding energy, middling and flour size distribution, and gluten and dough mixing properties. Journal of Cereal Science 69, 306-312. Doi: 10.1016/j.jcs.2016.04.007.
- Yamaoka, C., Kurita, O., Kubo, T., 2014. Improved ethanol tolerance of *Saccharomyces cerevisiae* in mixed cultures with *Kluyveromyces lactis* on high-sugar fermentation. Microbiological Research. 169, 907-914. Doi: 10.1016/j.micres.2014.04.007.
- Yuliana, R., Kusdiyantini, E., Izzati, M., 2014. *Aloe vera* flour to produce inulin to achieve final product of fructo-oligosaccharides (FOS) using *Kluyveromyces marxianus* DUCC-Y-003. Jurnal BIOMA 16(1), 39-49.
- Yusriyah, N.H., Agustini, R., 2014. Fermentation duration and concentration of kefir grain on the production of kefir from bovine milk. UNESA Journal of Chemistry 3(2), 53-57.

Total Sugar Content, Protein, Adhesiveness, and Hardness of Fermented

ORIGINALITY REPORT

31%

SIMILARITY INDEX

24%

INTERNET SOURCES

22%

PUBLICATIONS

16%

STUDENT PAPERS

PRIMARY SOURCES

1

appliedfoodtechnology.org

Internet Source

3%

2

www.locus.ufv.br

Internet Source

2%

3

Submitted to Universitas Diponegoro

Student Paper

2%

4

Submitted to Universiti Teknologi MARA

Student Paper

2%

5

Ghosh, Shiladitya, Ranjana Chowdhury, and Pinaki Bhattacharya. "Mixed consortia in bioprocesses: role of microbial interactions", *Applied Microbiology and Biotechnology*, 2016.

Publication

2%

6

Submitted to University of Stellenbosch, South Africa

Student Paper

2%

7

Submitted to Lovely Professional University

Student Paper

2%

8	www.redalyc.org Internet Source	2%
9	Vignolo, Graciela, Patricia Castellano, and Silvina Fadda. "Bioprotective Cultures", Handbook of Fermented Meat and Poultry, 2014. Publication	2%
10	tci-thaijo.org Internet Source	1%
11	www.innspub.net Internet Source	1%
12	eprints.undip.ac.id Internet Source	1%
13	alfonsototosaus.yolasite.com Internet Source	1%
14	jehse.biomedcentral.com Internet Source	1%
15	Rokhima Lusiantari, Miranti Dewi Pramaningtyas, Titis Nurmasitoh, Rachmi Hidayati Pattimura, Anggita Dewanti. "Shortening tends to increase aortic foam cell count and wall thickness in male Wistar rats", <i>Universa Medicina</i> , 2018 Publication	1%
16	"Yeast Diversity in Human Welfare", Springer	

Nature, 2017

Publication

1%

17

ejbiotechnology.info

Internet Source

1%

18

Amornrat Mukprasirt. "Physico-chemical properties of flour and starch from jackfruit seeds (*Artocarpus heterophyllus* Lam.) compared with modified starches", International Journal of Food Science and Technology, 3/2004

Publication

1%

19

jatp.ift.or.id

Internet Source

<1%

20

eprints.uny.ac.id

Internet Source

<1%

21

www.scilit.net

Internet Source

<1%

22

Clemencia Chaves-López, Rosanna Tofalo, Annalisa Serio, Antonello Paparella, Giampiero Sacchetti, Giovanna Suzzi. "Yeasts from Colombian Kumis as source of peptides with Angiotensin I converting enzyme (ACE) inhibitory activity in milk", International Journal of Food Microbiology, 2012

Publication

<1%

23 Clemencia Chaves-López, Antonello Paparella, Rosanna Tofalo, Giovanna Suzzi. "Proteolytic activity of *Saccharomyces cerevisiae* strains associated with Italian dry-fermented sausages in a model system", *International Journal of Food Microbiology*, 2011

Publication

24 article.sciencepublishinggroup.com

Internet Source

25 G. Oboh, M.M. Ekperigin, A.A. Akindahunsi. "Coagulants Modulate the Antioxidant Properties & Hypocholesterolemic Effect of Tofu (Curdled Soymilk)", *Nutrition and Health*, 2016

Publication

26 Cheok, Choon Yoong, Noranizan Mohd Adzahan, Russly Abdul Rahman, Nur Hanani Zainal Abedin, Norhayati Hussain, Rabiha Sulaiman, and Gun Hean Chong. "Current Trends of Tropical Fruit Waste Utilization", *Critical Reviews in Food Science and Nutrition*, 2016.

Publication

27 Swami, Shrikant Baslingappa, N. J. Thakor, P. M. Haldankar, and S. B. Kalse. "Jackfruit and Its Many Functional Components as Related to Human Health: A Review", *Comprehensive*

Reviews in Food Science and Food Safety, 2012.

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off