

Dietary Practical Ingredients With Emphasis on Protein Sources for Penaeid Shrimp

by Agung Sudaryono

Submission date: 16-Jul-2018 04:01PM (UTC+0700)

Submission ID: 982861309

File name: C7.pdf (88.13K)

Word count: 3333

Character count: 18156

DIETARY PRACTICAL INGREDIENTS WITH EMPHASIS ON PROTEIN SOURCES FOR PENAEID SHRIMP

Agung Sudaryono

Aquaculture Study Program Faculty of Fisheries and Marine Science, Diponegoro University
Jl. Hayam Wuruk 4-A, Semarang – Indonesia. E-mail: agungsud@indosat.net.id

Received August 10, 2001; accepted August 30, 2001

ABSTRACT

Studies on the utilization of dietary practical ingredients, in particular protein sources, by penaeid shrimp juveniles were reviewed in this paper. It has been proved by a number of researchers in shrimp nutrition that diets containing a mixture of two or more protein sources are better utilized by shrimp than those containing single protein sources. In general, evaluation of the nutritive value of a feed ingredient for growth is based on both the quantity and the quality of the protein in the ingredient. The amino acid composition and the relative proportions of other dietary nutrients are mostly relevant to the efficiency of protein assimilation by penaeids. Based on the essential amino acid index (EAAI) used to screen some potential protein sources for penaeid shrimp, there were found that fish meal, shrimp meal, squid meal and soybean meal were good dietary potential protein sources with EAAs between 0.87-0.98.

Key words: Penaeid shrimp, protein, dietary

INTRODUCTION

Experimental diets for penaeid shrimp are numerous. They include purified, semi-purified and practical diets and may be designed for larvae, postlarvae or juveniles (Akiyama, 1991). Purified diets refer to diets, which include a source of purified protein such as casein, gelatin and albumin alone or in combination. Practical diets are usually based on less refined protein sources, which are available for use in commercial diets. Commercial diets are diets processed in a factory and sold to farmers. In the previous section, some nutritional requirement studies of penaeids have been discussed. This section focuses on studies of the utilization of practical dietary ingredients, in particularly protein sources, by penaeid shrimp juveniles.

The Quality of Dietary Protein

The growth of shrimp is affected not only by the quantity of the dietary protein but also by the quality of the protein (Smith *et al.*, 1985). Diets containing a mixture of two or more protein sources are better utilized by shrimp than those containing a single protein source (Colvin, 1976; Alava and Lim, 1983; Rajyalakshmi *et al.*, 1986; Shiau *et al.*, 1991; Sudaryono *et al.*, 1995; Millamena and Trino, 1997). Sedgwick (1979) found that the optimal utilization of protein by *P. merguensis* was closely related to the energetic value of the diet and that carbohydrate and lipid can also increase growth efficiency at sub-optimal levels of protein. The efficiency of protein assimilation by penaeids is also likely to be affected by the relative proportions of

lipids and carbohydrates in the formulation as well as by the amino acid composition of whatever protein source is employed (Hanson and Goodwin, 1977).

The best protein sources for decapod crustaceans are those amino acid (AA) profile most closely mimic the decapod's protein AA profile (Deshimaru and Shigueno, 1972; Wilson and Poe, 1985; Aquacop and Cuzon, 1989; Kanazawa, 1990; Lim and Persyn, 1989; Sarac *et al.*, 1993). Since proteins constitute the largest chemical group within the decapod body, the nutritive value of a protein ingredient will be determined to a large extent by its amino acid composition, and in particular by the availability and profile of the essential amino acids (EAA) present. Deshimaru and Shigueno (1972) in their study on the artificial diets for *Penaeus japonicus* found that an inferior feed efficiency occurred when the diets were deficient in the essential amino acids, arginine, histidine, and lysine. Akiyama *et al.* (1992) stated that in making practical diet formulations for shrimp, one consideration is to provide enough of the dietary lysine, methionine, threonine and arginine, which are the most critical essential amino acids. In another trial, Deshimaru *et al.* (1985) reported that *P. monodon* fed diets with high percentages of all essential amino acids and high ratios of methionine and arginine to total essential amino acids, showed better growth, feed efficiency and survival. Lan and Pan (1993) added that the protein digestibility of diet increased when the amount of basic essential amino acids (arginine and lysine) and aromatic amino acid (phenylalanine) in the total essential amino acid composition of the diet increased. Akiyama *et al.* (1992) suggested that to maintain an optimum growth of shrimp, the dietary lysine: arginine ratio should be at 1:1.0 to 1:1.1. This is needed to avoid depressed growth due to a dietary lysine-arginine antagonism relationship.

Studies on Potential Protein Sources

Many workers incorporate squid meal as a dietary protein for shrimp. Squid meal is believed to be a good protein source in the diet and gives a significantly better growth for *P. indicus* (Bose, 1988), *P. japonicus* (Deshimaru and Shigueno, 1972), *P. monodon* (Lim *et al.*, 1979; Rajyalakshmi *et al.*, 1986; Cruz-Ricque *et al.*, 1987; Cruz-Suarez *et al.*, 1992), *P. setiferus* and *P. stylirostris* (Fenucci *et al.*, 1980) than other protein sources (concentrated soluble fish protein, fish meal, mysid meal, shrimp meal, soybean meal). Lim *et al.* (1979), using *P. monodon* post-larvae, also reported that squid meal and shrimp meal appeared to be good protein sources. Combinations of various dietary protein sources for *P. monodon* have also been tested. Pascual (1989) found a combination of 35% defatted soybean meal and 16% peruvian fish meal (40% protein) to give the best yield. Rajyalakshmi *et al.* (1986) reported that best performance was given by feeds with a combination of soybean and squid meals. Penaflores (1989) screened some potential protein sources for *P. monodon* feeds using the essential amino acid index (EAAI) calculated by comparing the ratio of EAAs in individual feedstuffs with that of whole *P. monodon* juvenile. The results showed that fishmeals, white and peruvian fishmeals, shrimp meal, squid meal, and soybean meal (EAAI of 0.92–0.95, 0.96, 0.94, 0.98, 0.96, and 0.87, respectively) were found to be good potential protein sources.

In growth experiments with various penaeid species, shrimp meal or shrimp waste meal also proved to be a good protein source (Chen *et al.*, 1986; Liang *et al.*, 1986; Godfred *et al.*, 1991; Hartnoll and Salama, 1992; Holland and Borski, 1993; Fox *et al.*, 1994). However, physical properties of shrimp meal with its high fibre content have resulted in a limitation of its use in shrimp diet. Increasing the dietary shrimp meal inclusion levels had a negative effect on

nutrient digestibility (Ali, 1992) and water stability (Akiyama *et al.*, 1992; Fox *et al.*, 1994) but a positive effect on palatability (Hartnoll and Salama, 1992; Fox *et al.*, 1994). Akiyama *et al.* (1992) suggested that dietary shrimp meal inclusion levels should not exceed 15%; normally shrimp meal is incorporated in shrimp diets with a range of 5–15%. Superiority of the mollusk meals, scallop meal as a protein source for the growth of juvenile *P. monodon* (Sudaryono *et al.*, 1995) and mussel extract (Murai *et al.*, 1985) have been reported. Hutabarat *et al.* (1995) found that the addition of a local material of kingworms (*Eisenia foetida*) in diets increased the growth of *P. monodon*.

Plant vs Animal Protein Sources

The use of soybean meal as a partial or complete alternative protein source for fish meal, shrimp meal, squid meal or a combination of two or more marine animal protein ingredients for various species of penaeid shrimp has been evaluated in several studies (Forster and Beard, 1973; Sick and Andrews, 1973; Colvin and Brand, 1977; Fenucci *et al.*, 1980; Lawrence *et al.*, 1986; Akiyama, 1989; Akiyama and FSGP, 1989; Lim and Dominy, 1990). Many of the reported results are contradictory. This may reflect differences in species and size of shrimp, diet composition, quality of the test ingredients, feeding management and culture conditions. Generally, however, high inclusion levels of dietary soybean meal or complete substitution of animal proteins resulted in poor growth and feed efficiency. Increasing the inclusion levels of dietary soybean meal over 42% was

Reported to have an adverse effect on pellet water stability and palatability (Lim and Dominy, 1990). Akiyama (1991) reviewed the utilization of soybean meal by marine shrimp and concluded that soybean meal was a highly digestible source of protein. Brunson *et al.* (1997) found that plant protein supplements

(soybean meal and cottonseed meal) were digested as efficiently as, or better than, proteins of animal origin (squid liver powder, menhaden fishmeal, shrimp meal and crab meal) by *P. setiferus*. Eusebio (1991) reported that *P. monodon* seemed to utilize the dehulled leguminous seeds (cowpea and rice bean) as protein sources more efficiently than the whole seeds.

GENERAL CONCLUSIONS

Although EAAI of the ingredients is a useful tool for predicting protein quality it does not mean that the individual amino acids present are biologically available and can be completely utilized by shrimp. To determine the biological availability of the amino acids, *in vitro* and/or *in vivo* digestibility measurements of ingredients are required (Akiyama *et al.*, 1989; Eid and Matty, 1989; Lan and Pan, 1993; Reigh *et al.*, 1990). The most common method used by researchers for estimating the nutrient digestibility of individual feedstuffs has been to evaluate a standard reference diet in which the test ingredient is incorporated at a single fixed level of 30% (Cho *et al.*, 1985; Akiyama *et al.*, 1989; Reigh *et al.*, 1990; Brunson *et al.*, 1997). Variations of nutrient digestibility value have been reported to attribute to differences in dietary protein source, ingredient dietary inclusion level, ingredient particle size, feed preparation method, and feeding regime (Forster and Gabbot, 1971; Akiyama and Dominy, 1991; Palaniswamy and Ali, 1991).

ACKNOWLEDGEMENTS

Thanks to Prof. L.H. Evans and Dr. E. Tsvetnenko from Aquatic Science Research Unit, Curtin University of Technology, Australia for their valuable review and suggestions of this manuscript.

Special thanks to anonymous reviewers for their revision on this manuscript.

REFERENCES

- Akiyama, D.M., 1991. Soybean meal utilisation by marine shrimp. In: D.M. Akiyama and R.K.H. Tan (Editors), Proceedings of the Aquaculture Feed Processing and Nutrition Workshop, Thailand and Indonesia, September 19-25. American Soybean Association, Singapore, pp. 207-225.
- Akiyama, D.M. and Dominy, W.G., 1991. Penaeid shrimp nutrition for the commercial feed industry. American Soybean Association and Oceanic Institute, Waimanalo, USA, 50 pp.
- Akiyama, D.M., Coelho, S.R., Lawrence, A.L. and Robinson, E.H., 1989. Apparent digestibility of feedstuffs by the marine shrimp *Penaeus vannamei* Boone. Bulletin of the Japanese Society of Scientific Fisheries, 55: 91-98.
- Akiyama, D.M., Dominy, W.G. and Lawrence, A.L., 1992. Penaeid shrimp nutrition. In: A.W. Fast and L.J. Lester (Eds.), Marine Shrimp Culture: Principles and Practices. Elsevier Science Publishers BV., pp. 535-568.
- Akiyama, D.M. and FSGP Aquaculture Research, 1989. The use of Soybean meal to replace white fishmeal in commercially processed *Penaeus monodon* Fabricius in Taiwan, R.O.C. In: Proc. Third Int. Symp. On Feeding and Nutr. in Fish, Toba, Japan, Aug. 28 - Sept. 1, 1989, pp. 289-299.
- Alava, V.R. and Lim, C., 1983. The quantitative dietary protein requirements of *Penaeus monodon* juveniles in a controlled environment. Aquaculture, 30: 53-61.
- Ali, S.A., 1992. Evaluation of some animal and plant protein sources in the diet of the shrimp *Penaeus indicus*. Asian Fisheries Society, 5: 277-289.
- Aquacop and Cuzon, G., 1989. Selected ingredients for shrimp feed. In: Advances in Tropical aquaculture, 20 Feb. - 4 March 1989, Tahiti. AQUACOP/IFREMER Actes de Colloque, 9: 405-412.
- Bose, S.V.C., 1988. Nursery rearing of seed of *Penaeus indicus* in nylon net cages. Presented at National Symposium on Research and Development in Marine Fisheries, Mandapam Camp (India), 16 Sep. 1987. CMFRI SPEC. PUBL., no. 40, pp. 55-56. Summary.
- Brunson, J.F., Romaine, R.P. and Reigh, R.C., 1997. Apparent digestibility of selected ingredients in diets for white shrimp *Penaeus setiferus* L. Aquaculture Nutrition, 3: 9-16.
- Chen, H.Y., Zein-Eldin, Z.P. and Aldrich, D.V., 1986. Combined effects of shrimp size and dietary protein source on the growth of *Penaeus setiferus* and *Penaeus vannamei*. Journal of the World Mariculture Society, 16: 288-296.
- Cho, C.Y., Cowey, C.B. and Watanabe, T., 1985. Fish nutrition in Asia. Methodological Approaches to Research and Development. Ottawa, Canada. International Development Research Center Publication No. IDRC-233e., 154 pp.

- Colvin, L.B. and Brand, C.W., 1977. The protein requirement of penaeid shrimp at various life cycle stages with compounded diets in controlled environment systems. Proceedings of The World Mariculture Society, 8: 821-840.
- Colvin, P.M., 1976. Nutritional studies on penaeid prawns: protein requirements in compounded diets for juvenile *Penaeus indicus* (Milne Edwards). Aquaculture, 7: 315-326.
- Cruz-Suarez, L.E., Ricque-Marie, D. and Aquacop, 1992. Effect of squid meal on growth of *Penaeus monodon* juveniles reared in pond pens and tanks. Aquaculture, 106: 293-299.
- Cruz-Suarez, L.E., Ricque-Marie, D., Martinez-Vega, J.A. and Wesche-Ebeling, P., 1993. Evaluation of two shrimp by-product meals as protein sources in diets for *Penaeus vannamei*. Aquaculture, 115: 53-62.
- Cruz-Ricque, L.E., Guillaume J. and Cuzon, G., 1987. Squid protein effect on growth of four penaeid shrimp. Journal of the World Aquaculture Society, 18: 209-217.
- Deshimaru, O. and Shigeno, K., 1972. Introduction to the artificial diet for prawn *Penaeus japonicus*. Aquaculture, 1: 115-133.
- Deshimaru, O., Kuroki, K., Mazid, M.A. and Kitamura, S., 1985. Nutritional quality of compounded diets for prawn *Penaeus monodon*. Bulletin of the Japanese Society of Scientific Fisheries, 51: 1037-1044.
- Eid, A.E. and Matty, A.J., 1989. Simple in vitro method for measuring protein digestibility. Aquaculture, 79: 111-119.
- Eusebio, P.S., 1991. Effect of dehulling on the nutritive value of some leguminous seeds as protein sources for tiger prawn, *Penaeus monodon*, juveniles. Aquaculture, 99: 297-308.
- Fenucci, J.L., Zein-Eldin, Z.P. and Lawrence, A.L., 1980. The nutritional response of two penaeid species to various levels of squid meal in a prepared feed. Proceedings of the World Mariculture Society, 11: 403-409.
- Forster, J.R.M. and Beard, T.W., 1973. Growth experiments with the prawn *Palaemon serratus* fed with fresh and compounded foods. Fisheries Investigation Series II, 27 (7), 16 pp.
- Fox, C.J., Blow, P., Brown, J.H. and Watson, I., 1994. The effect of various processing methods on the physical and biochemical properties of shrimp head meals and their utilization by juvenile *Penaeus monodon* Fabricius. Aquaculture, 122: 209-226.
- Forster, J.R.M. and Gabbott, P.A., 1971. The assimilation of nutrients from compounded diets by the prawns *Palaemon serratus* and *Pandalus platyceros*. Journal of Marine Biological Association United Kingdom, 51: 943-961.
- Godfred, J., Sriraman, K., Ramanathan, N. and Kannupandi, T., 1991. Feed formulation and its effect on the growth of four species of penaeid

- shrimp. *Aquaculture Europe*, 16: 26-28.
- Hanson, J.E. and Goodwin, H.L., 1977. Shrimp and Prawn Farming in the Western Hemisphere. Dowden, Hutchinson and Ross, Stroudsburg, Pa., 439 pp.
- Hartnoll, R.G. and Salama, A.J., 1992. The effect of protein source on the growth of the prawn *Palaemon elegans* Rathke, 1837 (Decapoda, Caridea). *Crustaceana*, 63: 81-90.
- Holland, K.N. and Borski, R.J., 1993. A palatability bioassay for determining ingestive stimuli in the marine shrimp *Penaeus vannamei*. *Aquaculture*, 109: 153-164.
- Hutabarat, J., Edhy, S. and Ma'ruf, W.F., 1995. Utilization of local raw materials to support shrimp feed industry in Central Java. Abstracts of the Fifth International Working Group on Crustacean Nutrition Symposium, April 22-24, 1995, Kagoshima University, Kagoshima, Japan, 62 pp.
- Kanazawa, A., 1990. Protein requirements of penaeid shrimp. In: *Advances in Tropical Aquaculture*, Tahiti, French Polynesia, Feb. 20 - March 4, 1989. IFREMER/SDP, Actes Colloq, 9: 261-271.
- Lan, C.C. and Pan, B.S., 1993. In-vitro digestibility simulating the proteolysis of feed protein in the midgut gland of grass shrimp (*Penaeus monodon*). *Aquaculture*, 109: 59-70.
- Liang, Yaquan, J. and Wenjuan, 1986. Protein requirement in formulated diets for penaeid shrimp (*Penaeus orientalis* Kishinouye) in different growth stage. *Marine Fisheries Research*, 7: 79-87.
- Lim, C. and Dominy, W.G., 1990. Evaluation of soybean meal as a replacement for marine animal protein in diets for shrimp (*Penaeus vannamei*). *Aquaculture*, 87: 53-63.
- Lim, C. and Persyn, A., 1989. Practical feeding - penaeid shrimps. In: T. Lovell (Editor), *Nutrition and Feeding of Fish*. Van Nostrand Reinhold, New York, pp. 205-222.
- Lim, C., Suraniranat, P and Platon, R.R., 1979. Evaluation of various protein sources for *Penaeus monodon* postlarvae. *Kalikasan, Philippines Journal of Biology*, 8(1): 29-36.
- Millamena, O.M. and Trino, A.T., 1997. Low-cost feed for *Penaeus monodon* reared in tanks and under semi-intensive and intensive conditions in brackishwater ponds. *Aquaculture*, 154: 69-78.
- Murai, T., Sumalangcay, A.Jr. and Pascual, F.P., 1985. Supplement of various attractants to a practical diet for juvenile *Penaeus monodon* Fabricius. *Fisheries Research Journal of Philippines*, 8: 61-67.
- Palaniswamy, R. and Ahmad Ali, S., 1991. The impact of particle size of ingredients in compound diets on pellet stability and performance in *Penaeus indicus* H. Milne Edwards. *Journal of Aquaculture in the Tropics*, 6: 119-127.
- Pascual, F.P., 1989. Nutritional feeding of *Penaeus monodon*. *Aquaculture*

- Extension Manual No. 3, June/89 (3rd Edition). Aquaculture Dept., SEAFDEC, Tigbauan, Iloilo, Philippines, 24 pp.
- Penafloida, V.D., 1989. An evaluation of indigenous protein sources as potential component in diet formulation for tiger prawn, *Penaeus monodon*, using essential amino acid index (EAAI). *Aquaculture*, 83: 319-330.
- Rajyalakshmi, T., Pillai, S.M., Roy, A.K. and Verghese, P.U., 1986. Feeds and nutrition of the prawn *Penaeus monodon*. *Environment and Ecology*, 4 (3): 415-422.
- Sarac, H.Z., Thaggard, H., Gravel, M., Saunders, J., Neill, A. and Cowan, R.T., 1993. Observations on the chemical composition of some commercial prawn feeds and associated growth responses in *Penaeus monodon*. *Aquaculture*, 115: 97-110.
- Sedgwick, R.W., 1979. Influence of dietary protein and energy on growth, food consumption and food conversion efficiency in *Penaeus merguensis* De Man. *Aquaculture*, 16: 7-30.
- Shiau, S.Y., Kwok, C.C. and Chou, B.S., 1991a. Optimal dietary protein level of *Penaeus monodon* reared in seawater and brackishwater. *Journal of the Japanese Society of Scientific Fisheries*, 57: 711-716.
- Sick, L.V. and Andrews, J.W., 1973. The effect of selected dietary lipids, carbohydrates and protein on the growth, survival and body composition of *Penaeus duorarum*. *Proceedings of the World Mariculture Society*, 4: 263-276.
- Smith, L.L., Lee, P.G. and Lawrence, A. L., 1985. Growth and digestibility by three sizes of *Penaeus vannamei* Boone : effect of dietary protein level and protein source. *Aquaculture*, 46: 85-96.
- Sudaryono, A., Hoxey, M.J., Kailis, S. G. and Evans, L. H., 1995. Investigation of alternative protein sources in practical diets for juvenile shrimp *Penaeus monodon*. *Aquaculture*, 134: 313-323.
- Wilson, R.P. and Poe, W.E., 1985. Relationship of whole body and egg essential amino acid pattern to amino acid requirement patterns in channel catfish, *Ictalurus punctatus*. *Comparative Biochemistry and Physiology*, 80B: 385-388.

Dietary Practical Ingredients With Emphasis on Protein Sources for Penaeid Shrimp

ORIGINALITY REPORT

38%

SIMILARITY INDEX

19%

INTERNET SOURCES

35%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Alava, V.R.. "The quantitative dietary protein requirements of *Penaeus monodon* juveniles in a controlled environment", *Aquaculture*, 198301 **7%**
Publication
 - 2** Lim, C.. "Evaluation of soybean meal as a replacement for marine animal protein in diets for shrimp (*Penaeus vannamei*)", *Aquaculture*, 19900515 **4%**
Publication
 - 3** A.G.J. TACON. "Nutritional studies in crustaceans and the problems of applying research findings to practical farming systems", *Aquaculture Nutrition*, 9/1996 **3%**
Publication
 - 4** Cuzon, G.. "Composition, preparation and utilization of feeds for Crustacea", *Aquaculture*, 199407 **2%**
Publication
-

5

www.aquacultureassociation.ca

Internet Source

2%

6

J. F. Brunson. "Apparent digestibility of selected ingredients in diets for white shrimp *Penaeus setiferus* L.", *Aquaculture Nutrition*, 3/1997

Publication

1%

7

Houng-Yung Chen. "Nutritional Requirements of the Black Tiger Shrimp: *Penaeus monodon*", *Reviews in Fisheries Science*, 3/1/1998

Publication

1%

8

Handbook on Ingredients for Aquaculture Feeds, 2000.

Publication

1%

9

Chhorn Lim. "Replacement of Marine Animal Protein with Peanut Meal in Diets for Juvenile White Shrimp, *Penaeus vannamei*", *Journal of Applied Aquaculture*, 05/30/1997

Publication

1%

10

www.j3.jstage.jst.go.jp

Internet Source

1%

11

Zhou, Y.-G., D.A. Davis, and A. Buentello. "Use of new soybean varieties in practical diets for the Pacific white shrimp, *Litopenaeus vannamei*", *Aquaculture Nutrition*, 2014.

Publication

1%

12	repository.seafdec.org.ph Internet Source	1%
13	docslide.us Internet Source	1%
14	Alam, Md.S.. "Supplemental effects of coated methionine and/or lysine to soy protein isolate diet for juvenile kuruma shrimp, <i>Marsupenaeus japonicus</i> ", <i>Aquaculture</i> , 20050729 Publication	1%
15	trjfas.org Internet Source	1%
16	researchonline.jcu.edu.au Internet Source	1%
17	www.researchgate.net Internet Source	1%
18	www.eolss.net Internet Source	1%
19	Sudaryono, A.. "Digestibility studies on fisheries by-product based diets for <i>Penaeus monodon</i> ", <i>Aquaculture</i> , 19960820 Publication	1%
20	A.E. Eid, A.J. Matty. "A simple in vitro method for measuring protein digestibility", <i>Aquaculture</i> , 1989 Publication	1%

21

jeb.biologists.org

Internet Source

1%

22

Louis D'Abramo. "Nutritional requirements, feed formulation, and feeding practices for intensive culture of the freshwater prawn *Macrobrachium rosenbergii*", *Reviews in Fisheries Science*, 1994

Publication

1%

23

www.2ndchance.info

Internet Source

1%

24

www.fao.org

Internet Source

<1%

25

Mauchamp, A.. "Submergence-induced damage of photosynthetic apparatus in *Phragmites australis*", *Environmental and Experimental Botany*, 200406

Publication

<1%

26

www.trjfas.org

Internet Source

<1%

27

Rebecca Lochmann. "Evaluation of Practical Feed Formulations and Dietary Supplements for Red Swamp Crayfish", *Journal of the World Aquaculture Society*, 9/1992

Publication

<1%

28

e.bangor.ac.uk

Internet Source

<1%

- 29 Modern Approaches to the Study of Crustacea, 2002. <1 %
Publication
-
- 30 www.ksuaquaculture.org <1 %
Internet Source
-
- 31 dyuthi.cusat.ac.in <1 %
Internet Source
-
- 32 Crisantema Hernández, Miguel A. Olvera-Novoa, David M. Smith, Ronald W. Hardy, Blanca Gonzalez-Rodriguez. "Enhancement of shrimp *Litopenaeus vannamei* diets based on terrestrial protein sources via the inclusion of tuna by-product protein hydrolysates", Aquaculture, 2011 <1 %
Publication
-
- 33 Perla S. Eusebio. "Effect of dehulling on the nutritive value of some leguminous seeds as protein sources for tiger prawn, *Penaeus monodon*, juveniles", Aquaculture, 1991 <1 %
Publication
-
- 34 Amin Oujifard. "Fish meal replacement with rice protein concentrate in a practical diet for the Pacific white shrimp, *Litopenaeus vannamei* Boone, 1931", Aquaculture International, 05/17/2011 <1 %
Publication
-

35

Stuart Biddle. "Towards an understanding of children's motivation for physical activity: Achievement goal orientations, beliefs about sport success, and sport emotion in zimbabwean children", *Psychology & Health*, 11/1/1996

Publication

<1%

36

Josafat Marina Ezquerro, Fernando L. García-Carreño, Roberto Civera, Norman F. Haard. "pH-stat method to predict protein digestibility in white shrimp (*Penaeus vannamei*)", *Aquaculture*, 1997

Publication

<1%

37

PATĪ, Siddhartha, SUMEDHA, Chinnari, CHATTERJĪ, Anil and DASH, Bisnu Prasad. "A Novel Food Formulation for the Juveniles of *Penaeus monodon* (Fabricius)", TÜBİTAK, 2016.

Publication

<1%

38

Penafiorida, V.D.. "An evaluation of indigenous protein sources as potential component in the diet formulation for tiger prawn, *Penaeus monodon*, using essential amino acid index (EAAI)", *Aquaculture*, 19891215

Publication

<1%

39

Lan, C.C.. "In-vitro digestibility simulating the proteolysis of feed protein in the midgut gland

<1%

of grass shrimp (*Penaeus monodon*"),
Aquaculture, 19930101

Publication

40

E.A.T. Floreto. "The effects of krill hydrolysate-supplemented soya-bean based diets on the growth, colouration, amino and fatty acid profiles of juvenile American lobster, *Homarus americanus*", Aquaculture Nutrition, 3/5/2001

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Dietary Practical Ingredients With Emphasis on Protein Sources for Penaeid Shrimp

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
