



PROCEEDINGS

VOLUME II

The 2nd International Conference
on Animal Nutrition and Environment
(ANI-NUE2017)

*“Towards the Betterment of Animal
Productivity, Conserving Resources
and Environment”*

November 1-4,
2017

Pullman Raja Orchid Hotel,
Khon Kaen, Thailand



EDITORS: Cherdthong, A., Foiklang, S., Mapato, C., Pilajun, R., Kang, SC. and Wanapat, M.

ISBN: 978-616-438-084-4

Organized by

Tropical Feed Resources Research and Development Center (TROFREC), Department of Animal Science,
Faculty of Agriculture, Khon Kaen University





In Recognition for

Professor Dr. Metha Wanapat
**“The Khon Kaen University Distinguished
Research Professor”**

for

**His 37 years of continuous serving in
teaching, research, education, technology
development, and transfer in Animal
Science nationally and internationally.**

**The 2nd International Conference on Animal Nutrition and
Environment (ANI-NUE2017) Organizing Committee,**

November 1, 2017

PROCEEDINGS
of
The 2nd International Conference on
Animal Nutrition and Environment
(ANI-NUE2017)
Volume II

Editors: Cherdthong, A., Foiklang, S., Mapato, C., Pilajun, R., Kang, SC. and Wanapat, M.

ISBN: 978-616-438-084-4

November 1-4, 2017

Pullman Raja Orchid Hotel, Khon Kaen, Thailand

Contents (Continued)

ANN-01-0071: “Effects of supplementation of Piper sarmentosum leaves powder in concentrates on feed efficiency, rumen fermentation and protozoa in Thai native beef cattle” <i>Anusorn Cherdthong and Benjamad Khonkhaeng-Thailand</i>	400
Session 9	
ANN-01-0061: “Determination of level protein intake to control fat and protein in carcass of fattened lambs” <i>Rizky Choirunnisa, Mukh Arifin, Robert Kusetyawan, Febrian Rhamadya, Ari Prima, Vita Restitrisnani, Nadlirotun Luthfi, Sutaryo and Agung Purnomoadi-Indonesia</i>	408
ANN-01-0044: “Effect of yeast-fermented cassava pulp levels on growth performance of growing goats fed Napier Pakchong 1 grass” <i>Saran Parisuthikul, Tichakorn Thumtong, Theerasant Phothee, Panuwat Teebklang, Wunchai Inthiseang, Wichan Kaewluan and Ruangyote Pilajun-Thailand</i>	415
ANN-01-0102: “Effect of mangosteen peel liquid protected soybean meal on methonogen and microbial population using in in vitro gas fermentation technique” <i>Kampanat Phesatcha, Burarat Phesatcha, Suban Foiklang, Metha Wanapat-Thailand</i>	422
ANN-01-0082: “Fermentation quality and in vitro digestibility of fermented total mix ration with difference roughage and fermentation period” <i>Pichad Khejornart, Teerayoot Jantanam and Metha Wanapat-Thailand</i>	429
ANN-01-0076: “Effect of replacement soybean meal by yeast waste on feed intake and rumen ecology in Thai native beef cattle” <i>Anusorn Cherdthong, Phussorn Sumadong, Suban Foiklang, and Nipa Milintawisamai-Thailand</i>	436
ANN-01-0077: “Chemical composition and In vitro gas production of the local Thai and India Moringa (<i>Moringa oleifera</i> Lam.) for ruminant” <i>Chaiyawan Wattanachant, Nattha Rattanacosol, Chaermpon Yuangklang, and Apichat Loopachr-Thailand</i>	441
Session 10	
ANN-01-0013: “Use of Body Measurements to Predict Intermuscular Fat in Thin-tailed Lambs” <i>Ulia Renfelia Baysi, Ari Prima, Farah Nabila, Pradhipta Hersandika, Endang Purbowati, Christina Maria Sri Lestari, and Agung Purnomoadi-Indonesia</i>	447
ANN-01-0099: “Effect of inclusion of enzyme from fermented tomato pomace with <i>Aspergillus niger</i> (FETPAN) in total mixed ration on feed intake and growth performance in beef cattle” <i>Chalermpon Yuangklang, Jiravan Khotsakdee, and Krisit Vasupen-Thailand</i>	452
ANN-01-0100: “Influence of Tropical Roughages Combined with Urea and Bamboo Grass (<i>Tiliacora Triandra</i>, Diels) Supplementation on Gas Production and In Vitro Degradability” <i>Chaichana Suriyapha, Metha Wanapat and Chinda Wann-Thailand</i>	457

Contents (Continued)

ANN-01-0087: “Effect of Bamboo grass (<i>Tiliacora Triandra</i>) pellet supplementation on feed intake, nutrient digestibility and rumen microbial population in Thai native beef” <i>Chinda Wann, Metha Wanapat, Choawarit Mapato and Chaichana Suriyapha-Thailand</i>	463
ANN-01-0105: “ <i>In Vitro</i> Fermentation and Methane Production Influenced by Leucaena Silage and Mangosteen Peel Powder” <i>Sungchhang Kang, Metha Wanapat, Ratana Khun, Laiheang Chhen, Piseth Try, Kimsan Sok, Ratha Long and Vannak Un-Cambodia</i>	469
ANN-01-0106: “Influence of Bamboo Leaf Meal Supplementation on <i>In Vitro</i> Gas Production and Digestibility” <i>Sungchhang Kang and Metha Wanapat-Cambodia</i>	475
Session 11	
ANN-01-0075: “Effect of Banana flower powder contained in high quality feed block and roughage to concentrate ratio on in vitro gas production kinetics, digestibility and fermentation” <i>Suban Foiklang, Metha Wanapat, Bounnaxay Viennasay, and Thitima Norrapoke-Thailand</i>	481
ANN-01-0047: “Feed intake and blood metabolite of goats fed urea-calcium hydroxide treated oil palm frond” <i>Pin Chanjula, Suradech phetarwut, and Anusorn Cherdthong-Thailand</i>	486
ANN-01-0070: “Blood chemistries of dairy cow during pre-calving, at calving and post-calving period” <i>Chamnanwit Promkot, Pitukpol Porn-aneke, and Teerawat Srinukool-Thailand</i>	491
ANN-01-0060: “The study on determination of feed digestibility using frequency of defecation on Thin Tailed lamb” <i>Febrianto Dwi Nugroho, Ari Prima, Edy Rianto, Agung Purnomoadi-Indonesia</i>	494
ANN-01-0062: “Effect of <i>Terminalia Chebula</i> RETZ. meal on nutrient intake, digestibility and microbial population of goats” <i>Nirawan Gunun, Pongsatorn Gunun, Anusorn Cherdthong, Sineenart Polyorach and Metha Wanapat-Thailand</i>	497
ANN-01-0090: “Effect of bamboo grass pellet (Bamboo-cass) levels on gas production kinetics and in vitro degradability” <i>Bounnaxay Viennasay and Metha Wanapat-Thailand</i>	500
ANN-01-0084: “Effect of fresh cassava root with feed block containing high sulfur on gas kinetics and rumen fermentation using <i>in vitro</i> gas production technique” <i>Gamonmas Dageaw and Anusorn Cherdthong-Thailand</i>	507
Session 12	
ANN-01-0072: “Inclusion of yeast waste as protein source to replace soybean meal in concentrate diet on ruminal fermentation and kinetics of gas using a gas production technique” <i>Anusorn Cherdthong, Rittikeard Prachumchai, Chanadol Supapong, Metha Wanapat, Suban Foiklang, and Nipa Milintawisamai-Thailand</i>	514

Contents (Continued)

ANN-01-0055: “Using of Urea and Molasses Fermented Cassava Pulp on Rumen Fermentation and Methane Production” <i>Thitima Norrapoke and Tanitpan Phongchongmit-Thailand</i>	519
ANN-01-0046: “The Meat Quality of Bali Beef fed with Supplement Blocks with Different Liquid Smoke Levels as Antioxidant and Binder” <i>Effendi Abustam, Muhammad Irfan Said, and Muhammad Yusuf-Indonesia</i>	526
ANN-01-0094: “Effect of Flemingia (<i>Flemingia macrophylla</i>) as a protein replacement of soybean meal on feed intake, digestibility of nutrients and microbial population in Thai native beef cattle” <i>Burarat Phesatcha and Metha Wanapat-Thailand</i>	532
ANN-01-0068: “Nutrient utilization and rumen ecology of Thai indigenous cattle given hay and sago palm pith with different levels of soybean meal” <i>Bunseelarp Wiyada and Ngampongsai Wanwisa-Thailand</i>	540
ANN-01-0085: “Comparison between Hay and Silage of <i>Pennisetum purpurium</i> cv. Mahasarakham feeding on feed intake, nutrient digestibility, and rumen fermentation in Thai native beef bulls” <i>Chaowarit Mapato and Metha Wanapat-Thailand</i>	545
ANN-01-0078: “Increasing Productive Performance of Native Chickens by Herbs in Rural Community” <i>Narumon Somkuna, Eakkasit Somkuna, Jarous Sawangtap and Phinithi Ratchwicha-Thailand</i>	549
Session 13	
ANN-01-0059: “Regression Models for Estimating Fat Carcass Percentage Using Chest Measurement in Thin Tailed Lambs” <i>Farah Nabila, Pradhipta Hersandika, Ari Prima, Vita Restitrisnani, Nadlirotun Luthfi, Endang Purbowati, Sutaryo and Agung Purnomoadi-Indonesia</i>	553
ANN-01-0065: “Development of Near Infrared Spectroscopy for nondestructive and rapid measurement of chemical compositions and somatic cell counts in raw milk” <i>Onanong Pongchompu, Metha Wanapat, Chaluntorn Vichasilp, Yaungyote Jindatajak and Pongsagorn Pongchompu-Thailand</i>	558
ANN-01-0014: “Effect of Hydroponic Maize Fodder Supplementation on Production Performance in Graded Murrah Buffaloes of Scarce Rainfall Zone” <i>Atturi Krishna Murthy, Dhanalakshmi Guduru, Y.G.Prasad, and Sarjan Reddy Kapa-India</i>	563
ANN-01-0081: “Effect of fermented total mixed ration with microbial culture on fermentation quality and <i>in vitro</i> digestibility” <i>Pichad Khejornsart and Metha Wanapat-Thailand</i>	567
ANN-01-0104- Effect of Addition of Siamese Neem Foliage on pH and Number of Lactic Acid Bacteria in Napier Grass Silage <i>Anan Petlum, Sukanya Kamphayae, Pramote Paengkoum, Walailuck Kaewwongsa, Smerjai Bureenok, Tanaporn Plong-uan, and Thanyalak Theppaw-Thailand</i>	574



Session 9-Orchid ballroom I

ANN-01-0061

Determination of Level Protein Intake to Control Fat and Protein in Carcass of Fattened Lambs

Rizky Choirunnisa, Mukh Arifin*, Robert Kusetyawan, Febrian Rhamadya, Ari Prima, Vita Restitrisnani, Nadlirotun Luthfi, Sutaryo, and Agung Purnomoadi

*Faculty of Animal and Agricultural Sciences, Diponegoro University,
Semarang, Indonesia, 50275*

**Corresponding email: mukh.arifin@live.undip.ac.id*

Abstract

Twenty four thin tailed lambs aged approximately 3 months old with an average body weight of 14.19 ± 2.41 kg (CV 16.98%) were used in this study to determine the level of dietary protein (percent protein intake per body weight; CPI in %BW) required to achieve a low fat content (less than 5%) in lamb meat preferred by consumer. The lamb were fed the diets containing 14, 16 and 18% crude protein (CP) and 60 and 70% total digestible nutrients (TDN). The lambs were slaughtered after 3 months raising under the feeding treatments which reached average slaughter weight 24.48 ± 3.45 kg. The carcass was weighed to obtain carcass weight, and was then separated into meat, fat and bone. The longissimus dorsi (LD) and Biceps femoris (BF) were used to determine the fat and protein content of carcass by proximate analysis. The data of level CPI (in %BW) was then correlated to fat and protein in carcass, and was analyzed to determine the level protein intake required to control carcass fat. The results showed that the level of CPI (in %BW) were significantly correlated ($P < 0.05$) with carcass fat, meat fat and carcass protein, being 0.11, 0.40 and 0.63 respectively. This study found that level of protein intake per body weight (CPI in %BW) for lamb should be less than 0.7%BW to obtain a consumer preferred low fat content (less than 5%) in lamb meat.

Keywords: lambs, percent protein intake per body weight, carcass fat, meat fat, meat protein



Introduction

In recent years, consumer realized that the high proportion of lipid consumption of meat give some effects on human health such as coronary heart disease, cancer and arthritis. Along with the increasing public awareness to healthy living, consumption of low meat fat increased (Scollan et al., 2006 ; World Cancer Research Fund/American Institute for Cancer Research, 2007 ; Bezerra et al., 2016) and the quality of meat can be identified through its flavor and tenderness (Jaworska et al., 2016 ; Malva et al., 2016 ; Zinder et al., 2017). Intramuscular fat (IMF) is being a factor to determine juiciness and flavour of meat (Lambe et al., 2017). Savel ad Cross (1998) claimed that a minimum level of IMF in meat lambs that can be accepted by consumer is 3%. Lambe et al. (2017) reported a preferred level of IMF of lamb meat is less than 5%.

In the postnatal period, adipose cells in the body of lamb is still carrying out to the hyperplasia process, adipose tissue develops its size of cell (hypertrophy) and increases the amount of cell (hyperplasia) (Hood and Allen, 1973). The increasing of adipose cells occurs as long as the live of lamb and fat accumulation in the meat of lambs will continue (Wangko and Wangko, 2010). Protein is one of nutrient contents that can affect fat proportion in meat. The excess of amino acids from protein would be utilized for fat synthesis in the carcass of lambs and being low quality carcass (Ponnampalam et al., 2003). Therefore, it is necessary to determine the level protein intake to get best quality of meat lambs.

Materials and Methods

Twenty four thin tailed lambs were used with an average body weight of 14.19 kg \pm 2.41 (CV 16.98%) aged \pm 3 months. The diets contained 14, 16 and 18% of crude protein (CP) and 60 and 70% of total digestible nutrients (TDN). The feed ingredients consisted of rice brand, cassava peel, sugarane top, cassava flour, soybean meal, fish meal, molasses and minerals. The feed and water were given ad libitum. The lambs were reared for 3 months and slaughtered. Lambs were fasted 6 hour prior to be slaughtered. The carcass of lambs was aging at 18°C for 10 hours, then was weight to determine carcass weight. Then carcass was separated between meat, fat and bone to obtained the percentage of carcass fat. The longissimus dorsi (LD) and Biceps femoris (BF) were used to determine the proximate composition (fat and protein).



The data was obtained and analyzed using correlation analysis. The relationship between the two variables was evaluated using the magnitude of the correlation value described by Steel and Torrie (1960).

Results and Discussion

Percent protein intake per body weight, carcass fat (%), fat of meat (%) and protein of meat (%) are presented in the Table 1. The dietary protein intake was 0.46 – 0.99 % per body weight and 0.67% as an average. The carcass fat (%) was around 1.41 - 33.96 % and 16.51% as an average. Fat of meat (%) was around 2.41 – 8.88 % and 4.22 % as an average. Protein of meat (%) was around 12.25 – 20.02% and 16.864% as an average).

Table 1 Protein intake per body weight (%), carcass fat (%), meat fat (%) and meat protein (%) of lambs.

Parameter	Range	Average	Standar Deviation
Protein intake / BW (%)	0.462 – 0.992	0.677	0.137
Carcass Fat (%)	1.413 – 33.964	16.516	6.597
Meat Fat (%)	2.410 – 8.880	4.225	1.436
Meat Protein (%)	12.250 – 20.020	16.864	2.682

Francisco et al. (2015) found that carcass fat of lamb was 20,9% with protein intake 0,7% per BW. Based on fat class and conformation (Speedy, 1980), lambs in this study is included as class 2 of fat. It was due to its carcass weight was around 14.18 kg (fat 16.71%, muscle 62.21% and bone 18.51%). The meat fat of lambs in this study was higher than those of previous study that taken by Francisco et al. (2015), the meat fat of lambs were 2%. According to Purbowati et al. (2010) the protein of lamb meat were 18,32% which is higher than this study. The growth of animals starts from nerves growth, bone growth, muscle growth and fat growth (Owens et al., 1993). Savel and Cross (1998) claimed that a minimum level of IMF in meat lambs that can be accepted by consumer is 3%. According to Lambe et al (2017), a preferred level of IMF of meat lamb is less than 5%. Protein content of meat is about 16 – 22% and the largest component was 75 – 80% of dry matter (Lawrie, 1995).




Figure 1 Correlation between % CP intake per BW to carcass fat (CF), meat fat (MF) and meat protein (MP).; %meat fat (●), %carcass fat (○), %meat protein (◆)

The result showed that protein intake per body weight (%) was linearly correlated with carcass fat (%), meat fat (%) and meat protein (%) ($P < 0.05$). Protein intake per body weight (%) has weak correlation with carcass fat (%) ($r = 0.1118$) and has medium correlation with meat fat (%) ($r = 0.3967$). However, protein intake per body weight (%) has strong correlation with meat protein (%) ($r = 0.6261$). Figure 1 showed that every increasing of protein intake per body weight (%) increases the carcass fat (%) and meat fat (%) but decreases the meat protein (%). It is due to the muscle has maximum capacity to receive protein intake. Therefore, the excessive protein intake would be used to grow fat in form of adipose cells by hyperplasia process. Muscle tissue in lambs grows as cell enlargement process (hypertrophy) but adipose cells in the body of lambs is still carrying out to be hyperplasia process where is adipose tissue develops size of cell (hypertrophy) and increases of the amount of cell (hyperplasia) during growth period. (Hood and Allen, 1973). The excess of amino acids from protein would be utilized for fat synthesis in the carcass of lambs and being low quality carcass (Ponnampalam et al., 2003).



If lamb with average body weight ± 25 kg consumed 3.5% dry matter intake/BW it can be calculated that lamb consumed 0.875 kg DMI. From this research it is known that the best protein intake to get low fat of meat is 0.7%/BW. Then, protein consumed by lamb with the average body weight ± 25 kg is 175 g. Therefore, protein content on diet that should be given to the lamb to get low fat meat is less than 20% .

Conclusions

Based on this study, it can be concluded that protein intake per body weight of lamb to achieve low fat meat (<5%) which is preferred by consumer as healthy meat is 0.7%. Then, it can be suggested that protein content on diet for lamb should be given less than 20%.

Acknowledgements

The authors would like to thank you to the Domba Cemara Team of Faculty of Animal and Agricultural Sciences, Diponegoro University, for their support and help in this study.

References

- Bezerra, L.S., A.M. Barbosa, G.G.P Carvalho, J.I. Simionato, J.E. Freitas, M.L.G.L. Araujo, L. Pereira, R.R Silvia, E.C.Q Lacerda and B.M.A. Carvalho. 2016. Meat quality of lambs fed diets with peanut cake. *Meat Sci.* 121: 88-95.
- Francisco, A., M.T. Detinho, S.P. Alves, P.V. Portugal, F. Fernandes, S. Sengo, E. Jeronimo, M.A. Oliveira, P. Costa, A. Sequeira, R.J.B. Bessa and J.S. Silva. 2015. Growth performance, carcass and meat quality of lambs supplemented with increasing levels of a tanniferous bush (*Cistus ladanifer* L.) and vegetable oils. *Meat Sci.* 100: 275-282.
- Hood, R. L. and C. E. Allen. 1973. Cellularity of bovine adipose tissue. *J. Lipid Research.* 14: 605-610.
- Jaworska, D., M. Czauderna, W. Przybylski, and A.J.R. Rozbicka-Wieczorek. 2016. Sensor quality and chemical composition of meat from lambs fed diets enriched with fish and rapessed oils, carnosic acid. *Meat Sci.* 119: 185-192.



- Lambe, N.R., K.A. McLean, J. Gordon, D. Evans, N. Clelland, and L. Bunger. 2017. Prediction of intramuscular fat content using CT Scanning of package lamb cuts and relationship with meat ageing quality. *Meat Sci.* 123: 112-119.
- Lawrie, R.A. 1995. *Meat Science*. Woodhead Publishing Limited, CRC Press, New York, DC.
- Malva, A.D, M. Albenzio, G. Annicchiarico, M. Caroprese, A. Muscio, and A. Santillo. 2016. Relationship between slaughtering age, nutritional and organoleptic properties of Altamura lamb meat. *Small Rum. Res.* 135: 39-45.
- Owens, F.N., P. Dubeski and C.F. Hanson. 1993. Factors that alter the growth and development of ruminants. *J. Anim. Sci.* 71: 3138-3150.
- Ponnampalam, E.N., B.J. Hosking, and A.R. Egan. 2003. Rate of carcass components gain, carcass characteristics and muscle longissimus tenderness in lambs fed dietary protein sources with a low quality roughage diet. *Meat Sci.* 63: 143-149.
- Purbowati, E., U. Hasanah, R. Adiwinarti, C. I. Sutrisno, E. Baliarti, S. P. S. Budi and W. Lestariana. Komposisi kimia daging Domba Lokal akibat pemberian pakan komplit dari berbagai limbah pertanian dan agroindustri. *Prosiding Seminar Nasional Teknologi Peternakan dan Veteriner*. Bogor, 3-4 Agustus 2010. Puslitbang Peternakan, Bogor. 573 – 543.
- Scollan, N.D., J.F. Hocquette, K. Nuernberg, D. Dannenberger, R.I. Richardson, and A. Maloney. 2006. Innovations in beef production systems that enhance the nutritional and health value of beef lipids and their relationship with meat quality. *Meat Sci.* 74: 17–33.
- Steel, R.G.D. and J.H. Torrie,. 1960. *Principles and Procedures of Statistics*. McGraw-Hill Book Co., New York.
- Wangko, W.S and S. Wangko. 2010. Adipogenesis tumbuh kembang adiposit. *Biomedik* 2 (3): 153-161.
- World Cancer Research Fund/American Institute for Cancer Research. 2007. *Food, Nutrition, Physical Activity and the Prevention of Cancer: A Global Perspective*. American Institute for Cancer Research, Washington, DC.
- Zinder, M.C., A. Orlov, O. Trofimiyuk, R. Agmon, R. Kabiya, E.S. Shimoni, E.K Wagner, K. Hussey, H. Leibovich, J. Miron, and A. Shabtay. 2017. Dietary