

JURNAL

by Tri Retnaningsih Soeprbowati

Submission date: 13-Feb-2019 03:36PM (UTC+0700)

Submission ID: 1077517691

File name: d_with_human_activities_in_Lake_Warna_Dieng_Plateau_Indonesi.pdf (2.22M)

Word count: 6652

Character count: 35300



Diatom assemblage in the 24 cm upper sediment associated with human activities in Lake Warna Dieng Plateau Indonesia



Tri Retnaningsih Soeprbowati^{a,b,*}, Sri Widodo Agung Suedy^b, Hadiyanto^{a,c},
Ali Arman Lubis^d, Peter Gell^e

^a School of Postgraduate Studies, Diponegoro University, Semarang, Indonesia

^b Department of Biology, Faculty of Science and Mathematics, Diponegoro University, Semarang, Indonesia

^c Department of Chemical Engineering, Faculty of Engineering, Diponegoro University, Semarang, Indonesia

^d Center for Isotopes and Radiation Application, National Nuclear Energy Agency, Indonesia

^e Water Research Network, Federation University Australia, Ballarat, Australia

19

ARTICLE INFO

Article history:

Received 30 July 2017

Received in revised form 21 March 2018

Accepted 31 March 2018

Available online 11 April 2018

Keywords:

Diatoms
Human activities
Lake Warna Dieng
pH
Eutrophic
Paleolimnology

ABSTRACT

Lake Warna is a small shallow crater lake on the Dieng Plateau, Central Java, the second highest plateau in the world after Nepal. A 24 cm sediment core was extracted from Lake Warna to reconstruct environmental changes in the Lake and its catchment from present diatom assemblages. Diatoms are microalgae in the Bacillariophyte that have siliceous cell walls that can be preserved in sediments. As diatom species are sensitive to water quality changes in the assemblages upcore reflect changes in lake condition. Sediment cores were collected from two sites, sliced at 1 cm intervals for diatom analysis and bulked across 3 cm for ²¹⁰Pb radiometric dating. Examination of diatoms in a 24 cm sediment core from Lake Warna reveals clear correlation with human activities in the catchment area over the past 124 years. The record is divided into 2 zones based on sustained changes in the diatom assemblages. The lowest zone Zone I (21–15 cm, estimated 1935–1954) was dominated by *Frustulia crassinervia* (Brebisson ex W. Smith) Ross, *Gomphonema parvulum* (Kutzing) Kutzing, *Pinnularia valdetolerans* Mayama & H. Kobayasi, *P. viridis* (Nitzsch) Ehrenberg, and *Aulacoseira distans* (Ehrenberg) Simonson. Zone II (15–0 cm, estimated 1980–2013), the uppermost zone was dominated by *P. viridiformis* Krammer, *P. latevittata* Cleve, *E. monodon* var. *tropica* (Hustedt) Hustedt, *S. seminulum* Grunow, *P. a crossphaeria* W. Smith, *E. zygodon* Ehrenberg, *P. gibba* Ehrenberg, and *P. viridiformis* Krammer. The 15 cm core from TW3 commenced sediment accumulation 114 years ago and is divided into 2 zones as well. Zone I (15–7 cm, estimated 1901–1981) was dominated by *Brachysira brebissonii* R. Ross, *G. parvulum* (Kutzing) Kutzing, *E. monodon* var. *tropica* (Hustedt) Hustedt, *E. zygodon* Ehrenberg and *S. seminulum* Grunow. Zone II (7–0 cm, estimated 1981–2006) was dominated by *P. viridiformis* Krammer, *P. gibba* Ehrenberg, *Pinnularia viridis* (Nitzsch) Ehrenberg, *Sellaphora bacillum* (Ehrenberg) Mann, *S. seminulum* Grunow, and *Stauroneis* sp. These assemblages reflect ongoing acid, clear water conditions for the time represented by the cores. The recent rise in *S. seminulum* Grunow reflects recent lake eutrophication likely owing to the accelerated agricultural and urban development in the lake's catchment in recent decades.

© 2018 Elsevier B.V. All rights reserved.

5

* Corresponding author at: School of Postgraduate Studies, Diponegoro University, Semarang, Indonesia.

E-mail address: trsoeprbowati@live.undip.ac.id (T.R. Soeprbowati).

20

<https://doi.org/10.1016/j.eti.2018.03.007>

2352-1864/© 2018 Elsevier B.V. All rights reserved.

1. Introduction

The Dieng Volcanic Complex, called the Dieng Plateau, hosts the highest village on Java, located in the Wonosobo and Banjarnegara Regencies, on the slopes of the active Sindoro and Sumbing Volcanoes. The Dieng Plateau, the second highest plateau in the world after Nepal, has many surface manifestations of hydrothermal activity, including hot lakes, fumaroles/solfatara, mud pools, altered rocks, and hot springs. Dieng Plateau is also rich in geothermal resources and is subject to lethal outbursts of gas. The many temples on the Plateau attest to the long establishment of ancient Indian Hindu culture that established a firm footing in Java from about AD 400 (van Bergen et al., 2000). The Dieng Plateau is 14 km long and 6 km wide and geologically consists of late Quaternary to Recent volcanic cones and explosion craters. There are 3 major volcanic activities resulting in volcanic rocks comprising Old Dieng, Adult Dieng, and Young Dieng (Sukhyar, 1994). During the second episode, a number of stratovolcanoes emerged within the depression, producing basalts, basaltic andesites and pyroxene-andesites. Pyroclastic fall deposits, believed to have been erupted from all of these volcanoes, blanket the Dieng and Batur depressions. They are collectively referred to as the 'Dieng tephra' for which dating yielded an age of 16,770 years. A parasitic eruption centered on the southern slope contains an 800 m wide and > 150 m deep crater lake (Telaga Menjer), which is used for hydroelectric power and irrigation. Panganan and Merdada are two stratocones east of Nagasari. The latter has a crater lake which is used for drinking water by local villagers. This crater, which has been partly filled by a biotite-andesite lava from the third episode, contains a colored lake, Lake Warna (van Bergen et al., 2000).

The high sulfur content, and possibly other elements, causes this lake's colorful state which ranges from blue to yellowish-green, and sometimes brownish, especially when exposed to sunlight. The color is a reflection of sunlight by sediment/rock at the bottom of the lake or refraction from fine sediments within the water. Red and yellow colors are a reflection of the sulfur deposits, and the white color comes from the deposition of limestone rocks and quartz. Now however, the only color visible is mostly blue on account of the sediment surface (Soeprbowati et al., 2017). In terms of chemical composition, Lake Warna is the most interesting crater lake in the Dieng area. The original shape of the crater has been modified by a lava flow and now the water occupies <1 km². Gas bubbles with a sulfurous odor can be seen rising to the lake surface. The water has a pH of about 3 (van Bergen et al., 2000) to 5 (Soeprbowati et al., 2017) which may fluctuate depending on seasonal conditions. The lakes chemistry is unusual with high concentrations of sulfate and chloride. Strong emissions of CO₂-rich gas are variable but have been intense enough to occasionally kill animals, warranting the closure of a path on the north side of the lake (van Bergen et al., 2000).

Dieng plantations over the years only used a monoculture system with potatoes as the main commodity. The slope of the land in Dieng area ranges from 35% to more than 45%, therefore, the farmers developed tillage systems terracing on contours as far as the tops of the hills. The erosion rate has reached 161 tons/hectare/year. About 7758 hectares of land has become critically unstable and has high erosion rates. In 2007, the Dieng area was 63.22% dominated by non-forest areas and this increased to 66.1% in 2010 (Mulyana, 2008). Lack of vegetation cover and the use of monoculture systems had induced sedimentation and damaged the ecosystems within this important watershed, such as Tulis and Serayu (Fig. 1, Pradana et al., 2015).

Lake Pengilon (Pengilon = mirror) located next to and connected to Lake Warna, is also notable. The two lakes host specific ecosystems that differ from other lakes. However, due to land use changes in the surrounding area both lakes are now facing environmental degradation. The land use in the surrounding area is dedicated to intensive agriculture with the main crops being vegetables, especially potatoes. Agricultural practices for plantation potato production use water from Lake Pengilon to irrigate the plants by pumping and irrigation. Agricultural practices expose soils to erosion, which contributes sediment to the lake via surface runoff. The volume of the lake is gradually decreasing due to water abstraction. Further, the overuse of fertilizer for agricultural production contributes nutrients to the lake leading to eutrophication (Sudarmadji and Purastuti, 2015).

Diatoms are microalgae that can be found in most aquatic ecosystems. Their silicified cell walls are often well preserved in sediments so diatoms are particularly useful for reconstructing environmental history. Different species have distinct optima to given environmental variables, and they are taxonomically distinct allowing identification to species and sub species level (Smol, 2008). They are highly sensitive to variations in the aquatic environment so they can be used as bioindicators of lake conditions. The lake's sediment reflects the history of changes in the catchment area (Battarbee et al., 2011; Di et al., 2013; Tolkkinen et al., 2014; Gell and Reid, 2014). This history can be reconstructed by reference to the nutrient requirements or typical habitats of each fossil taxon, or by consideration of their relative species abundance.

Palaeolimnological studies offer an opportunity to understand past and present lake condition and so, from the inferred trajectory, to predict future conditions. Preliminary research was conducted on the water quality of Lake Warna over the period of 2014–2015. Diatoms are particularly sensitive to pH and the acidification of surface waters is known to negatively impact on aquatic organisms, reduce diversity, and induce shifts in community structure. Diatoms can act as an early detection mechanism for decreasing pH and so can flag the need for the implementation of mitigation measures.

Hustedt (1937–1939) found *E. septentrionalis* lived in acid waters (pH ca.3) of the sulfur-springs of Sumatra. Negoro reported these varieties occurring as epiphytes from many highly acidic environments in Japan (DeNicole, 2000). The presence of *E. septentrionalis* in Lake Warna may be on account of the influence of the sulfur springs as was recorded in Sumatra (Soeprbowati et al., 2017).

Studies on diatoms in Indonesian lake sediments have in general been very limited in the analysis to determine indicator species and the changes of relative abundance through time (Soeprbowati et al., 2018). Notable among these are Dama



Fig. 2. Sites for coring sediment in Telaga Warna Dieng (Soeprbowati et al., 2017). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

disks for 10 min while stirring. ^{210}Po was assumed to be in radioactive equilibrium with ^{210}Pb . The measurement of activity was conducted using an alpha spectrometer equipped with a PIPS detector. For calibration of the detector, a point source of alpha emitters (^{238}U , ^{234}U , ^{239}Pu and ^{241}Am) was used while reference material IAEA-301 was used for controlling the method. Supported ^{210}Pb was calculated from the mean constant of concentration of ^{210}Pb at the bottom of the sediment core (Cossa et al., 2014). The excess ^{210}Pb was equal to the total ^{210}Pb minus the supported fraction. The depth-age profile was determined using the constant rate of supply (CRS) model that assumes a constant influx of unsupported, atmospheric ^{210}Pb to the site (Appleby, 2001).

Diatoms were chosen as a primary palaeolimnological tool. Diatom extraction was conducted following the procedure of Battarbee et al. (2001), whereby the subsamples are digested in 10% HCl and subsequently 10% H_2O_2 with multiple washes in distilled water between stages. Samples were mounted onto slides and a minimum of 300 diatom valves were identified under 1,000x magnification with oil immersion using a compound microscope (Soeprbowati et al., 2016a). The diatoms were identified to species level where possible, whereas the others were identified to genus level following standard texts (Sonneman et al., 2000; Krammer and Lange-Bertalot, 2004a, 2004b, 2004c; Krammer and Lange-Bertalot, 2010; Taylor et al., 2007; Karthick et al., 2013; Bahls, 2017). Guiry and Guiry (2011) was used for consulting taxonomy and nomenclature. The counts of diatom valves were expressed as relative abundance. Species with a relative abundance less than 1% were not included in the analysis and the remaining assemblage was graphed using the software package C2 version 1.7.7 (Juggins, 2016). The diatom assemblages were positioned into statistically significant clusters by Bray Curtis Clustering performed using PAST (Paleontological statistics software package for education and data analysis) version 2.17c (Hammer et al., 2001; Hammer, 2017).

3. Results

Seventeen sediment samples were used to establish the chronology of the 72 cm long TW2 core, and 16 sediment samples were used to establish the chronology of the 65 cm long TW3 core. The unsupported ^{210}Pb activities in the TW2 core exhibited an overall decreasing profile with depth (Fig. 3). The unsupported ^{210}Pb activities below 30 cm (TW2) and 18 cm (TW3) are close to being constant, showing that the activity had reached background and that the sediments below these levels were deposited > 100 years ago. This also indicates that the sediments in the upper 24 cm were deposited in the last 124.9 years (deposited since the year 1891) for TW2. Meanwhile, the sediment in the upper 12–15 cm of TW3 were deposited in the last 114 years (deposited since the year 1901) (Tables 1a and 1b).

Diatoms were well preserved in the lake sediments with most valves being entire or many valves were fragmented or dissolved. A total of 62 diatom species were identified from TW2 and 47 diatom species were identified from TW3. The identifications were made by comparing the species descriptions in published floras and comparing the morphometrics of valves measured down the microscope (length, width, striae density etc.) to the published ranges. Where any valves observed from the Warna sediments did not clearly match the features of published species they were assigned affinity status.

Based on the cluster analysis, basically the two cores of TW2 and TW3 can be divided into 2 zones, upper (Zone II) and bottom (Zone I), that are separated with a blue line in Fig. 2. Within the upper zones there were sub zones with red color (Figs. 3 and 4). For TW2, the zonations are: **Zone I (21–15 cm, estimated 1935–1954)**, the lowest zone was dominated by *Frustulia crassinervia* (Brebisson ex W. Smith) Ross, *Gomphonema parvulum* (Kutzing) Kutzing, *Pinnularia valdetolerans* Mayama & H. Kobayasi, *P. viridis* (Nitzsch) Ehrenberg, and *Aulacoseira distans* (Ehrenberg) Simonson. **Zone II (15–0 cm, estimated 1980–2013)** can be divided into 2 sub zones IIb (9–0 cm, estimated 2003–2013), the uppermost zone was




Fig. 3. Total ^{210}Pb versus depth of the TW2 core (left) and TW3 core (right), decay trend.

Table 1a

Age estimates determined from ^{210}Pb analysis in TW2.

Depth (cm)	Estimated date (year)	Calendar age	Sediment accumulation rates ($\text{gr}/\text{cm}^2 \text{ y}$)
0–3	2.4 ± 0.1	2013	0.31 ± 0.013
3–6	6.9 ± 0.2	2008	0.18 ± 0.005
6–9	12.4 ± 0.2	2003	0.13 ± 0.002
9–12	18.3 ± 0.9	1997	0.13 ± 0.007
12–15	34.7 ± 1.3	1980	0.04 ± 0.002
15–18	60.9 ± 2.6	1954	0.03 ± 0.001
18–21	81.9 ± 6.8	1933	0.03 ± 0.003
21–24	124.2 ± 16.0	1891	0.02 ± 0.002

Table 1b

Age estimates determined from ^{210}Pb analysis in TW3.

Depth (cm)	Estimated date (year)	Calendar age	Sediment accumulation rates ($\text{gr}/\text{cm}^2 \text{ y}$)
0–3	10.8 ± 0.6	2004	0.10 ± 0.005
3–6	21.1 ± 1.3	1994	0.05 ± 0.003
6–9	33.7 ± 2.0	1981	0.031 ± 0.002
9–12	62.8 ± 3.8	1952	0.015 ± 0.001
12–15	113.7 ± 6.6	1901	0.008 ± 0.001

dominated by *P. viridiformis* Krammer, *P. latevittata* Cleve, *E. monodon* var. *tropica* (Hustedt) Hustedt, *S. seminulum* Grunow, and *P. acrosphaeria* W. Smith. Sub zone IIa (15–9 cm, estimated 1997–1980) was dominated by *E. zygodon* Ehrenberg, *S. seminulum* Grunow, *P. gibba* Ehrenberg, and *P. viridiformis* Krammer. Diatoms assemblages in 1891 differed to that in other layers owing to the dominance of *F. saxonica* Rabernhorst in these layers.

For TW3, the 2 zones are: Zone I (15–7 cm, estimated 1901–1981), above Zone I of TW2 that was dominated by *Brachysira brebissonii* R. Ross, *G. parvulum* (Kutzing) Kutzing, *E. monodon* var. *tropica* (Hustedt) Hustedt, *E. zygodon* Ehr 36 erg and *S. seminulum* Grunow. Zone II (7–0 cm, estimated 1981–2006), just below the Zone II of TW2, was dominated by *P. viridiformis* Krammer, *P. gibba* Ehrenberg, *P. viridis* (Nitzsch) Ehrenberg, *Sellaphora bacillum* (Ehrenberg) Mann, *S. seminulum* Grunow, and *Stauroneis* sp.

The interesting note from the diatom stratigraphy from TW 2 and TW3 is that the population of *P. gibba* tended to increase in the uppermost sediment in contrast with *P. viridiformis* Krammer and *G. parvulum* (Kutzing) Kutzing that tended to decrease in the uppermost sediment (see Fig. 5).

displaced towards the secondary side of the sternum; internally they are liform and straight. The proximal (central) part of each raphe branch curves gently towards the primary side externally and the central raphe endings are slightly drop-like. Internally, the raphe branches remain slit-like as they approach the center and are bent towards the primary side. Beyond the helictoglossa, a larger foramen-like (pit) can be seen (Figs. 295, 300, 301) at the poles. Distal raphe endings strongly curved onto the mantle, towards the secondary side (Figs. 295–298).

Over the last 100 years, Lake Warna has been known for its acidic clear water. *Sphagnum* is dominant in the bed of the lake, and, owing to the water clarity, benthic diatoms were also dominant compared to planktonic diatoms. Several planktonic species, such as *A. distans* (Ehrenberg) Simonson, and *M. varians* C. Agardh were found sporadically in several layers but with low numbers. Some species of *Eunotia* and *Pinnularia* that were found are acidophilus, so Lake Warna remained acidic throughout. One of the most widely distributed taxa in acid lakes is *P. acoricola*. Hustedt described it from sulfate rich waters in Java in 1935, and fossil specimens of *P. acoricola* var. *lanceolata* Hustedt was found in Lake Toba, Sumatra (Hustedt, 1935). A trend of increasing abundance and diversity of *Pinnularia* spp. from Zone II indicates that the pH was acid to neutral (Sonneman et al., 2000; Hobbs et al., 2009). This is supported by the results of the modern water quality which recorded pH values in the range of 2.2 to 5.35 from 2014–2015.

Lake Warna is a geothermal lake that releases air bubbles to the surface of the lake. This indicates that the region still experiences magmatism activity, that is a hydrothermal process that emits vapors containing sulfur. This ensures that Lake Warna is an acid lake with pH below 5. Lakes with pH < 4 are typically dominated by acidophilus diatom species, such as *Eunotia paludosa* and *E. veneris* (Sonneman et al., 2000), many species in the genera *Eunotia*, *Pinnularia*, *Nitzschia*, and *Frustulia* as well as *Achnanthis minutissima* (DeNicole, 2000). In Lake Rawapening, *E. bilunaris*, *E. minor*, *E. monodon* (Hustedt) Hustedt, and *E. pectinalis* var. *pectinalis* found in abundance indicates a pH of less than 6 or neutral (Soeprbowati et al., 2012). In the case of Lake Rawapening, temperature, phosphate and calcium were the modern parameters that were found to influence the diatom assemblage (Soeprbowati et al., 2016b).

E. monodon var. *tropica* (Hustedt) Hustedt and *E. zygodon* Ehrenberg numbers sharply increased after 1980 and tended to decline in the upper layers of Lake Warna. Those *Eunotia* seem to be specific to acidic lakes, and, assuming their pH preference remains stable over time, these changes can be used to infer pH changes in Lake Warna. As their abundance fluctuated and shifted it is clear that recent inputs have influenced lake acidity. *E. monodon* var. *tropica* (Hustedt) Hustedt was the dominant species from 1980–1997 while *E. zygodon* Ehrenberg was dominant from 1997–2003. Their decline after 2003 may suggest that incoming nutrients have advantaged other species in recent times.

E. exigua (Brébisson ex Kützing) Rabenhorst is a cosmopolitan species that is found from tropical to arctic regions in lakes, ponds, bogs and hot springs (Joynt and Wolfe, 2001). The dominance of the acidobiontic taxon *E. exigua* (Brébisson ex Kützing) Rabenhorst in Lake Warna from 1991–2007 indicates a pH of 2–4 through this time (Hobbs et al., 2009). The acidic condition was also indicated by the dominance of *A. minutissimum* (Kützing) in uppermost centimeter of sediment (Soeprbowati et al., 2017).

F. saxonica Rabenhorst was dominant and only found in 1891, and *F. crassineervia* dominant in the following layers and tend to decrease from 1980. *Frustulia* Rabenhorst is a genus 36 phytobenthic diatoms, that are abundant in acidic habitats, such as *Sphagnum* peat bogs and many ephemeral habitats (Round et al., 2000; Wehr and Sheath, 2002; Siver and Baskette, 2004). The dominance of *F. crassineervia* indicated that, from 1891–1954, Lake Warna was oligotrophic. The two species 22 similar morphologically with small differences were correlated with environmental condition. Variation in the shape and size of valves has shown strong relationships with different types of peat land habitats (Kulichova and Fialova, 2016).

G. parvulum (Kützing) Kützing, *E. exigua* (Brébisson ex Kützing) Rabenhorst, and *E. minor* are only found in Zone 1 (1954–1933). *G. parvulum* (Kützing) Kützing is a cosmopolitan species (Abarca et al., 2014), mostly found in circumneutral and eutrophic waters (Kovacs et al., 2006; Karthick et al., 2013) but Zelazna-Wieczorek and Mamainska (2006) found *G. parvulum* (Kützing) Kützing in a pH of 3. *G. parvulum* (Kützing) Kützing is also reported as a common species in clean or metal contaminated systems (Ivorra et al., 2002).

The sediment core was separated into the relatively natural part (1891–1954 yr, lower layer: 24–15 cm) and the human disturbed part (1980–2013 yr, upper layer: 15–0 cm). The diatom abundance changes are abrupt in 1997. The use of fertilizer for intensive agricultural improvement since 1970s is likely to increase the nutrient loads in the Lake Warna that induced a eutrophic lake, indicated by an increase in *S. seminulum* Grunow since 1980 (Fig. 4). The deforestation and illegal logging combined to enable extensive monocultures of potatoes. As a result, there was high erosion and a reduced water supply (Sudarmadji and Pudjiastuti, 2018), and decreased land fertility that has impacted on potato productivity since 1997 (Pradana et al., 2015). This low land fertility is shown on the shift at 9 cm of diatom abundance from *S. seminulum* Grunow to *P. viridiformis* Krammer and *E. zygodon* Ehrenberg. This might suggest reduced nutrient inputs, or the onset of a secondary impact such as increased acidity or sediment input. The causes of these detected changes can be related to the changes in land use in the catchment area through reference to the chronostratigraphy and 15 interpretation of the historical record. The diatom record reveals continuous change through time and has enabled the impacts of changing land use practices on the lake's water quality to be identified. This trajectory indicates that the services provided by the lakes are at risk if effective management is not implemented.

Water quality monitoring of Telaga Warna commenced in 2014–2016. Lake Warna is an acidic eutrophic lake, with pH in the range of 2.2–5.35. In the dry season, the pH is about 2, and in the wet season, the pH is between 4 and 5. Today Lake Warna is eutrophic to hypereutrophic with the recorded total nitrogen concentration above 1.9 mg/L and total phosphorus




Plate 1. 1. *Eunotia zygodon* Ehrenberg. 2. *Eunotia minor* Ehrenberg. 3. *Frustulia saxonica* Rabenhorst. 4. *Frustulia crassinervia* (Brébisson) Lange-³⁵lot and Krammer. 5. *Gomphonema parvulum* Kützing. 6. *Sellaphora seminulum* (Grunow) D.G. Mann. 7. *Aulacoseira distans* (Ehrenberg) Simonsen. 8. *Eunotia exigua* (Brébisson ex Kützing) Rabenhorst. 9. *Eunotia minor* Kützing. 10. *Pinnularia acrosphaeria* W. Smith. 11. *Pinnularia gibba* Ehrenberg. 12. *Pinnularia viridiformis* Krammer. 13. *Neidium affine* (Ehrenberg) Pfitzer. 14. *Pinnularia viridis* (Nitzsch) Ehrenberg.

concentration above 0.1 mg/L. The recorded cadmium and lead content in Lake Warna was higher than the Indonesian water quality standard for agriculture, that was 0.01 and 0.03 mg/L, respectively

There is only little evidence of the changing conditions from these data owing to the short duration of the record (24 cm). But, as this is the only historical record available, the documentation of changing diatoms in the lake, as recorded in the sediment sequence, represents the only means by which the history of this lake can be reconstructed. This provides evidence of the trajectory of change as well as the variability of condition over time. It attests to the modern Lake Warna being different in condition to that evident in 1891. This then provides evidence that it has been impacted by human activity and flags the need to manage human impact to restore the condition of the lake. Paleolimnological studies in Indonesia are still limited (Soeprbowati et al., 2018). This study was initiated to provide evidence that sound lake management can be developed when the past condition is well understood. Other works, such as pollen analysis to determine crop pollen, XRF to determine erosion input, or stable isotope analysis – e.g. ^{15}N , would be useful to strengthen understanding to geode management options.

5. Conclusion

Based on diatom record from a 24 cm sequence of sediment from the upper section of a core taken from Telaga Warna has shown that human activities (deforestation and potatoes farming) in the catchment area have changed an acid oligotrophic lake into an acid eutrophic one. This is indicated by changes in the diatom assemblage from *Frustulia* spp. to *S. seminulum* Grunow. This initial result represents a baseline for further research for the development of appropriate management plans for the sustainable use of the lake (see Plate 1).

Acknowledgments

This article was produced as a part of project supported by Diponegoro University Indonesia through Research Grant for International Publication (RPI), PNBP DIPA Undip Number 1052-57/UN7.5.1/PG/2016. The heartfelt thanks to Kenanga Sari, M. Hadi El Amin, Siti Mudakiroh, Alam, and Geyga for helping in the field work and lab work.

References

- Abarca, N., Jahn, R., Zimmermann, J., Enke, N., 2014. Does the cosmopolitan diatom *Gomphonema parvulum* (Kützting) Kützting have a biogeography? *PLoS One* 9 (1), e86885.
- Appleby, P.G., 2001. Chronostratigraphic techniques in recent sediments. In: Last, W.M., Smol, J.P. (Eds.), *Tracking Environmental Change using Lake Sediments: Basin Analysis, Coring, and Chronological Techniques*. Kluwer Academic Press, Dordrecht, pp. 171–204.
- Bahls, L.L., 2017. *Diatoms from Western North America 1. Some New and Notable Biraphid Species*. Montana, USA.
- Battarbee, R.W., Bennion, H., Gell, P., Rose, N., 2011. Human impacts on lacustrine ecosystems. In: *The Sage Handbook of Environmental Change*. Sage, pp. 583–606.
- Battarbee, R., Jones, V.J., Flower, R.J., Cameron, N.G., Bennion, H., Carvalho, L., Juggins, S., 2001. Diatoms. In: Smol, J.P., Birks, H.J.B., Last, W.M. (Eds.), *Tracking Environmental Change using Lake Sediments. Volume 3: Terrestrial, Algal and Silicious Indicators*. Kluwer Academic Publishers, pp. 155–202.
- Carroll, J., Williamson, M., Lerche, I., Karabanov, E., Williams, D.F., 1999. Geochronology of lake baikal from ^{210}Pb and ^{137}Cs radioisotopes. *Appl. Rad. Isotop.* 50, 1105–1119.
- Cossa, D., Buscail, R., Puig, P., Chiffolleau, J.F., Radakovitch, O., Jeanty, G., Heussner, S., 2014. Origin and accumulation of trace elements in sediments of the northwestern Mediterranean margin. *Chem. Geol.* 380, 61–73.
- Dama, R.A.C., Fluin, J., Suparan, P., van der Kaars, S., 2001. Palaeoenvironmental development in the lake Tondano area (N Sulawesi Indonesia), since 33,000 yr BP. *Palaeogeog. Palaeoclim. Palaeoecol.* 171, 147–183.
- DeNicole, D., 2000. A review of diatoms found in highly acidic environments. *Hydrobiologia* 433 (1), 111–122.
- Di, B., Liu, D., Wang, Y., Dong, Z., Li, X., Shi, Y., 2013. Diatom and silicoflagellate assemblages in modern surface sediments associated with human activity: A case study in Sishili Bay, China. *Ecol. Indicators* 24, 23–30.
- Gell, P., Reid, M., 2014. Assessing change in floodplain wetland condition in the Murray Darling Basin, Australia. *Anthropocene* 057, e1–e7.
- Guiry, M.D., Guiry, G.M., 2018. *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. [13.02.18]. <http://www.algaebase.org>.
- Hammer, O., 2017. PAST: Paleontological statistics software package for education and data analysis version 2.17c.
- Hammer, O., Harper, D.A.T., Ryan, P.D., 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontol. Electron.* 4 (1), 9. http://palaeo-electronica.org/2001_1/past/issue1_01.htm.
- Hobbs, W.O., Wolfe, A.P., Inskeep, W.P., Amskold, L., Konhauser, K.O., 2009. Epipellic Diatoms from an Extreme Acid Environment: Beowulf Spring, Vol. 135. *Yellowstone National Park, U.S.A. Nova Hedwigia, Beiheft*, pp. 71–83.
- Hustedt, F., 1935. Die fossile Diatomeenflora in den Ablagerungen des Tobasees auf Sumatra.–Tropische Binnengewässer, Band VI.–Archiv für Hydrobiologie. Supplement 14, 143–192.
- Ivorra, N., Barranguet, C., Jonker, M., Kraak, M.H.S., Admiraal, W., 2002. Metal-induced tolerance in the freshwater microbenthic diatom *Gomphonema parvulum*. *Environ. Pollut.* 116, 147–157.
- Joynt II, E.H., Wolfe, A.P., 2001. Palaeoenvironmental inference models from sediment diatom assemblages in Baffin Island lakes (Nunavut, Canada) and reconstruction of summer water temperature. *Can. J. Fish. Aquat. Sci.* 58, 1222–1243.
- Juggins, S., 2016. *The European Diatom Database User Guide Version 1.7.7*. University of New Castle, New Castle, UK.
- Karthick, B., Hamilton, P.B., Kociolek, P., 2013. *An Illustrated Guide to Common Diatoms of Peninsular India*. Gubbi Labs LLP.
- Kovacs, C., Kahlert, M., Padisak, J., 2006. Benthic diatom communities along pH and TP Gradients in Hungarian and Swedish streams. *J. App. Phycol.* 18 (2), 105–117.
- Kramer, K., Lange-Bertalot, Ha, 2004a. *Susswasserflora Von Mitteleuropa, Bd. 02/2: Bacillariophyceae: Teil 2: Bacillariophyceae, Epithemiaeaceae, Surirellaceae*. Spektrum, Berlin.
- Kramer, K., Lange-Bertalot, Ha, 2004b. *Susswasserflora Von Mitteleuropa, Bd. 02/3: Bacillariophyceae: Teil3: Centrales, Fragillariaceae, Eunotiaceae*. Spektrum, Berlin.
- Kramer, K., Lange-Bertalot, Hc, 2004c. *Susswasserflora Von Mitteleuropa, Bd. 02/4: Bacillariophyceae: Teil 4: Achnanthes S.L., Navicula Sstr.* Spektrum, Berlin.
- Krammer, K., Lange-Bertalot, H., 2010. *Susswasserflora Von Mitteleuropa, Bd. 2/1: Bacillariophyceae, Teil 1: Naviculaceae*, first ed. Spektrum Akademischer Verlag, p. 882 ISBN-10:3827426154.
- Kulichova, J., Fialova, M., 2016. Correspondence between morphology and ecology: Morphological variation of the *Frustulia crassinervia-saxonica* Species Complex (Bacillariophyta) Reflects the Ombro-Minerotrophic Gradient. *Cryptogam. Algal.* 37 (1), 15–28.
- Lubis, A.A., 2006. Constant of rate supply (CRS) model for determining the sediment accumulation rates in the coastal area using ^{210}Pb . *J. Coast. Dev.* 10, 9–18.
- Mann, D.G., Thomas, S.J., Evans, K.M., 2008. Revision of the diatom genus *Sellaphora*: a first account of the larger species in the British Isles. *Fottea, Olomouc* 8 (1), 15–78.
- Mulyana, A.R., 2008. Evaluation of Land Utilization and Mud Flow Controlling in Dieng Plateau. Pusat Litbang Sumberdaya Air, Yogyakarta, Indonesia, pp. 5–7.
- Pradana, A., Sara, P.H., Wahdaningrum, W., 2015. The analysis of environmental degradation and carica agroforestry system as an attempt of environmental restoration in dieng plateau. *Int. J. Environ. Sci. Dev.* 6 (11), 861–867.
- Round, F.E., Crawford, R.M., Mann, D.G., 2000. *The Diatoms, Biology and Morphology of the Gera*. 3th Reprint. Cambridge University Press, Cambridge.
- Siver, P.A., Baskette, G., 2004. A morphological examination of *Frustulia* (Bacillariophyceae) from Ocala National Forest, Florida USA. *Ca. J. Bo.* 82, 629–644.
- Smol, J.P., 2008. *Pollution of Lakes and Rivers a Palaeoenvironmental Perspective*, second ed., Blackwell Publishing, USA.

- Soeprbowati, T.R., Hadisusanto, S., Gell, P., Zawadski, A., 2012. The diatom stratigraphy of Rawapening Lake, implying eutrophication history. *Am. J. Environ. Sci.* (3), 334–344.
- Soeprbowati, T.R., Suedy, S.W.A., Hadiyanto, , 2017. Diatoms and water quality of Telaga Warna Dieng, Java. *IOP. Conf. Ser. Earth. Environ. Sci.* 55, 012051.
- Soeprbowati, T.R., Suedy, S.W.A., Hadiyanto, , 2018. Find the future from the past: Paleolimnology in Indonesia. *E3S Web Conf.* 31, 08002.
- Soeprbowati, T.R., Tandjung, S.D., Sutikno, , Hadisusanto, S., Gell, P., 2016a. The minimum number of valves for diatom identification in Rawapening Lakee, Central Java. *Biotropia* 23 (2), 96–104.
- Soeprbowati, T.R., Tandjung, S.D., Sutikno, , Hadisusanto, S., Gell, P., Hadiyanto, , Suedy, S.W.A., 2016b. The water quality parameters controlling diatoms assemblage in Rawapening Lake, Indonesia. *Biodiversitas* 17 (2), 657–664.
- Sonneman, J.A., Sincock, A., Fluin, J., Reid, M., Newall, P., Tibby, J., Gell, P., 2000. *An Illustrated Guide to Common Stream Diatom Species from Temperate Australia*. Cooperative Research Centre for freshwater Ecology, Albury.
- Sudarmadji, , Pudjiastuti, H., 2015. Impact of agricultural practices on the sustainability of Telaga Warna Lake, Dieng Plateau, Central Java. In: *Proceeding Intngernational Symposium Integrated Actions for Global Water and Environmental Sustainability*, Medan 21-22 October 2015. Commemoration of the 70th Anniversary of UNESCO, pp. 46–49.
- Sudarmadji, , Pudjiastuti, H., 2018. Impacts of agricultural practices and tourism activities on the sustainability of Telaga Warna and Telaga Pengilon lakes, dieng plateau, central java. *E3S Web Conf.* 31, 08030.
- Sukhyar, R., 1994. Dieng Plateau, Central Java. Guidebook of the 1994. In: *Fifth Field Workshop on Volcanic Gases, Volcanological Survey of Indonesia*, Bandung.
- Taylor, J.C., Harding, W.R., Archibald, C.G.M., 2007. *An Illustrated Guide to Some Common Diatom Species from South Africa*, WRC Report TT 282/07.
- Theng, T.L., Ahmad, Z., Mohamed, C.A.R., 2003. Estimation of sedimentation rates using ²¹⁰Pb and ²¹⁰Po at the coastal water of Sabah, Malaysia. *J. Rad. Nuclear. Chem.* 256 (1), 115–120.
- Tolkkinen, M., Marttila, H., Saukkoriipi, J., Martinmki, K., Tammela, S., Tertsunen, J., Heikkinen, K., Tuohino, J., Ihme, R., Klove, B., 2014. pH-levels in intensively drained and peatland-dominated river basin: Palaeolimnological approach to detect impacts of past land use. *Ecol. Eng.* 64, 367–376.
- van Bergen, M.J., Bernard, A., Sumarti, S., Sriwana, T., Sitorus, K., 2000. *Crater lakes of Java: Dieng, Kelud and Ijen*. Excursion Guidbook. IAVCEI General Assembly, Bali.
- Wehr, J.D., Sheath, R.G., 2002. Freshwater habitats of algae. In: Wehr, J.D., Sheath, R.G. (Eds.), *Freshwater Algae from North America; Ecology and Classification*. Elsevier Science, San Diego, California USA, pp. 11–57.
- Wetzel, C.E., Ector, L., Van De Vijner, B., Compere, P., Mann, D.G., 2015. Morphology, typi cation and critical analysis of some ecologically important small naviculoid species (Bacillariophyta). *Fottea, Olomouc* 15 (2), 203–234.
- Zelazna-Wieczorek, J., Mamainska, M., 2006. Algoflora and vascular flora of a limestone spring in the Warta River Valey. *Acta Soc. Bot.* 75 (2), 131–143.

JURNAL

ORIGINALITY REPORT

17%

SIMILARITY INDEX

14%

INTERNET SOURCES

10%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | link.springer.com
Internet Source | 2% |
| 2 | STOCKNER, JOHN G., and WOODRUFF W. BENSON. "The succession of diatom assemblages in the recent sediments of Lake Washington", <i>Limnology and Oceanography</i> , 1967.
Publication | 1% |
| 3 | Sudarmadji, Hermin Pudjiastuti. "Impacts of Agricultural Practices and Tourism Activities on the Sustainability of Telaga Warna and Telaga Pengilon Lakes, Dieng Plateau, Central Java", <i>E3S Web of Conferences</i> , 2018
Publication | 1% |
| 4 | volcano.si.edu
Internet Source | 1% |
| 5 | www.ilec.or.jp
Internet Source | 1% |
| 6 | ainse.edu.au
Internet Source | 1% |

7	biodiversitas.mipa.uns.ac.id Internet Source	1%
8	federation.edu.au Internet Source	1%
9	esp.cr.usgs.gov Internet Source	<1%
10	comaniciu.net Internet Source	<1%
11	tel.archives-ouvertes.fr Internet Source	<1%
12	aos.wisc.edu Internet Source	<1%
13	www.omicsonline.com Internet Source	<1%
14	Muhammad Nur, Aribat Solichin, Endang Kusdiyantini, Tri A. Winarni et al. "Ozone production by Dielectric Barrier Discharge Plasma for microbial inactivation in rice", 2013 3rd International Conference on Instrumentation, Communications, Information Technology and Biomedical Engineering (ICICI-BME), 2013 Publication	<1%
15	Kathryn H. Taffs. "A diatom-based Holocene record of human impact from a coastal	<1%

environment: Tuckean Swamp, eastern
Australia", Journal of Paleolimnology, 01/2008

Publication

16

pythagoras.federation.edu.au:8443

Internet Source

<1%

17

espace.library.uq.edu.au

Internet Source

<1%

18

Portevin, Damien, Sudarkodi Sukumar, Mireia Coscolla, Guanghou Shui, Bowen Li, Xue Li Guan, Anne K. Bendt, Douglas Young, Sebastien Gagneux, and Markus R. Wenk. "Lipidomics and genomics of Mycobacterium tuberculosis reveal lineage-specific trends in mycolic acid biosynthesis", MicrobiologyOpen, 2014.

Publication

<1%

19

www.rslakes.com

Internet Source

<1%

20

www.mysmu.edu

Internet Source

<1%

21

uad.portalgaruda.org

Internet Source

<1%

22

www.cryptogamie.com

Internet Source

<1%

23

eprints.nottingham.ac.uk

Internet Source

<1%

24	doaj.org Internet Source	<1%
25	users.monash.edu.au Internet Source	<1%
26	www.er-c.org Internet Source	<1%
27	ddd.uab.cat Internet Source	<1%
28	qspace.library.queensu.ca Internet Source	<1%
29	www.phycologia.org Internet Source	<1%
30	bomax.botany.pl Internet Source	<1%
31	espace.curtin.edu.au Internet Source	<1%
32	Mackay, A.W.. "The paleoclimatology of Lake Baikal: A diatom synthesis and prospectus", Earth Science Reviews, 200706 Publication	<1%
33	repository.up.ac.za Internet Source	<1%
34	"Applications of Paleoenvironmental	<1%

Techniques in Estuarine Studies", Springer
Nature, 2017

Publication

35

www.algology.ru

Internet Source

<1%

36

jlimnol.it

Internet Source

<1%

37

Van de Vijver, Bart, Luc Ector, Myriam de Haan, and Ralitsa Zidarova. "THE GENUS *MICROCOSTATUS* IN THE ANTARCTIC REGION", Diatom Research, 2010.

Publication

<1%

38

Teresa Noga, Łukasz Peszek, Jadwiga Stanek-Tarkowska, Anita Pajączek. "The Pinnularia genus in south-eastern Poland with consideration of rare and new taxa to Poland", Oceanological and Hydrobiological Studies, 2014

Publication

<1%

39

jalgalbiomass.com

Internet Source

<1%

40

ir.canterbury.ac.nz

Internet Source

<1%

41

Kittel, Piotr, Błażej Muzolf, Mateusz Płóciennik, Scott Elias, Stephen J. Brooks, Monika Lutyńska, Dominik Pawłowski, Renata

<1%

Stachowicz-Rybka, Agnieszka Wacnik, Daniel Okupny, Zbigniew Głąb, and Aldona Mueller-Bieniek. "A multi-proxy reconstruction from Lutomiersk–Koziówki, Central Poland, in the context of early modern hemp and flax processing", *Journal of Archaeological Science*, 2014.

Publication

42

www.biogeosciences-discuss.net

Internet Source

<1%

43

Coles, Anna E., Carlos E. Wetzel, Núria Martínez-Carreras, Luc Ector, Jeffrey J. McDonnell, Jason J. Frentress, Julian Klaus, Lucien Hoffmann, and Laurent Pfister. "Diatoms as a tracer of hydrological connectivity: are they supply limited? : DIATOMS AS A TRACER OF HYDROLOGICAL CONNECTIVITY", *Ecohydrology*, 2015.

Publication

44

"Palaeolimnological Proxies as Tools of Environmental Reconstruction in Fresh Water", Springer Nature America, Inc, 2009

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

