

South_African_Journal_of_Animal_Science_-2017.pdf

by T Yudiarti

Submission date: 15-Aug-2018 07:02PM (UTC+0700)

Submission ID: 990136299

File name: South_African_Journal_of_Animal_Science_-2017.pdf (502.84K)

Word count: 6305

Character count: 31738

1
Effect of feeding duration of *Spirulina platensis* on growth performance, haematological parameters, intestinal microbial population and carcass traits of broiler chicks

11 **S. Sugiharto[#], T. Yudiarti, I. Isroli & E. Widiastuti**
Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Central Java, Indonesia (50275)

35
(Received 2 October 2017; Accepted 7 November 2017; First published online 16 November 2017)

2
Copyright resides with the authors in terms of the Creative Commons Attribution 4.0 South African Licence.
See: <http://creativecommons.org/licenses/by/4.0/za>
Condition of use: The user may copy, distribute, transmit and adapt the work, but must recognise the authors and the South African Journal of Animal Science.

Abstract

Spirulina platensis is a good candidate as an in-feed antibiotics substitute for broilers. However, its use seems impractical owing to its high price, especially when being administered throughout the whole rearing period. This study aimed to investigate the effects of feeding duration of *S. platensis* on growth, haematological parameters, intestinal microbial population, and carcass traits of broiler chicks. A total of 288 one-day-old broiler chicks were randomly allotted to one of four groups, including control (basal diet with 0.04% zinc bacitracin) (CONT) and birds receiving basal diet supplement with 1% of *S. platensis* for the first seven days (SP-7), for 21 days (SP-21), and for 35 days (SP-35). In this study, treatments had no significant effect on the growth performance of broilers. The caecum relative weight was significantly higher in SP-35 than in CONT and SP-21 birds. The values of haemoglobin, erythrocytes, and haematocrit were significantly lower in SP-35 than in other birds. Compared with CONT, SP-35 birds had significantly lower numbers of leukocytes, lymphocytes, and a lower number of eosinophils. The numbers of coliform were significantly lower in the ileum of SP-21 than in CONT and SP-7 birds. In the caecum, coliform tended to be lower in SP-21 than in other birds. There was no significant difference in the carcass traits of broilers across the groups. In conclusion, the administration of *S. platensis* for the first 21 days of broilers' life resulted in similar or even better responses than administration of *S. platensis* or in-feed antibiotics throughout the rearing period.

Keywords: Blood profile, bodyweight, broilers, carcass quality, feeding period, green alga, in-feed antibiotics
[#] Corresponding author: sgh_undip@yahoo.co.id

Introduction

To reach the maximum genetic potential for growth in broiler chickens, several nutritional strategies have been conducted, one of which is the use of antibiotics as feed additive (in-feed antibiotics). Sugiharto (2016) highlighted that in-feed antibiotics may be associated with improved growth and feed efficiency, as well as reduced morbidity and mortality in broiler production. However, owing to the development of antibiotic-resistant bacteria in humans and animals, the use of such additives was banned in most countries. Indeed, the withdrawal of synthetic antibiotics from broiler feeds led to performance and health problems in broiler chickens (Pourhossein *et al.*, 2015; Sugiharto, 2016). For food safety reasons and sustainable broiler production, using alternatives to replace in-feed antibiotics for broilers was therefore crucial. Microalgae, for instance *Chlorella vulgaris* and *Spirulina platensis*, have recently attracted considerable interest among poultry nutritionists owing to their high nutritional and functional properties, which may be beneficial for broiler chickens (Jamil *et al.*, 2012; Sugiharto & Lauridsen, 2016). With regard to *S. platensis* in particular, this microalga has high contents of protein, essential amino acids, vitamins, minerals, essential fatty acids and pigments (Beheshtipour *et al.*, 2013; Holman & Malau-Adu, 2013). *Spirulina platensis* is also rich in polysaccharides, which may function as prebiotics (Beheshtipour *et al.*, 2013; de Jesus Raposo *et al.*, 2016).

Several studies have demonstrated the growth-promoting effects of *S. platensis* on broiler chicks. Jamil *et al.* (2015) showed that feeding 2, 4, or 8 g of *S. platensis*/kg feed increased and decreased the body weight and feed conversion ratio (FCR) of broiler chickens, respectively. Similar results were reported by Shanmugapriya *et al.* (2015a) when feeding 1% of *S. platensis* to broiler chicks. These authors revealed that such treatment resulted in increased villi height, and hence improved absorption capacity of broiler intestines. With regard to the health of chicks, *S. platensis* administration has been demonstrated to

decrease the numbers of *Escherichia coli* and increase lactic acid bacteria (LAB) in the intestine of broilers (Shanmugapriya *et al.*, 2015b). Concomitantly, Yusuf *et al.* (2016) reported an increased *Lactobacilli* population in the gut of Japanese quails with feeding *S. platensis*. In terms of immunity, earlier studies by Qureshi *et al.* (1996) and Raju *et al.* (2004) reported that feeding *S. platensis* enhanced the humoral and cellular immune responses and lymphoid organ development of chicks. Recently, Lokapirnasari *et al.* (2016) showed that treatment with *S. platensis* increased the number of leukocytes and decreased the mortality rate of broiler chicks. In their review, Farag *et al.* (2016) pointed out that the antimicrobial and immunomodulatory (and anti-inflammatory) capacities as well as antioxidant potential seemed to be responsible for the health-promoting effect of *S. platensis* on poultry. As well as growth and health performances, treatment with *S. platensis* was reported to increase carcass percentage and ready-to-cook yields of broiler chicks in the studies of Raju *et al.* (2004), Kaoud (2012) and Holman & Malau-Aduli (2013). Moreover, Bonos *et al.* (2016) showed that Spirulina supplement (5 g/kg) was capable of improving the meat quality of broiler, that is, increasing the contents of eicosapentaenoic acid (EPA), docosapentaenoic acid (DPA) and docosahexaenoic acid (DHA) in the thigh muscles of broiler chicks. Taken together, *S. platensis* seems to have potential to replace the role of antibiotics in broiler diets.

As an alternative to in-feed antibiotics, *S. platensis* has commonly been administered to broiler diets from day of hatch to slaughter age (Qureshi *et al.*, 1996; Jamil *et al.*, 2015; Bonos *et al.*, 2016). Compared with antibiotics and other broiler feed ingredients, *S. platensis* is more expensive (Holman & Malau-Aduli, 2013). Apart from its benefits, the application of Spirulina as the replacement for in-feed antibiotics in broiler production therefore seems impractical, especially when used throughout the rearing period. For this reason, poultry nutritionists should reconsider the feeding duration of *S. platensis* to reduce the amount and cost of the microalga used in broiler production. With regard to probiotics, Pietras (2001) found no effect of feeding duration (that is, from days 1 to 21, from days 22 to 49 and from days 1 to 49 old) of probiotic *Lactobacillus acidophilus* and *Streptococcus faecium* on the production variables of broilers. In the current study, *S. platensis* was administered in broiler feeds for various durations within the rearing period. The objective of this study was therefore to investigate the effects of feeding duration of *S. platensis* on growth performance, haematological parameters, intestinal microbial population and carcass traits of broiler chicks.

Materials and methods

Two hundred and eighty-eight Lohmann (MB-202) one-day-old broiler chicks (bodyweight 42.0 ± 0.22 g) (means \pm standard deviation) were used in the current study. The chicks were randomly distributed to one of four groups of 72 chicks each (6 replicates of 12 chicks). Throughout the study period, the chicks were raised in an open-sided broiler house with rice husk-littered floor pens. The treatment groups included control birds (birds receiving basal diet with 0.04% zinc bacitracin) (CONT) and birds receiving basal diets supplemented with 1% of *S. platensis* for the first seven days (SP-7), for 21 days (SP-21), and for 35 days (SP-35). The proportion of *S. platensis* supplemented to basal diets was based on Shanmugapriya *et al.* (2015a). Feeds (as mash form) and water were provided *ad libitum* throughout the study period. A coccidiostat was not included in the feeds. The chemical compositions of *S. platensis* powder and basal diets (formulated to meet the Indonesian National Standards for Broiler Feed (SNI, 2006)) provided to broiler chicks are presented in Tables 1 and 2, respectively. Zinc bacitracin or *S. platensis* was added at the expense of the feeds. *S. platensis* was obtained from PT. Neoalga Indonesia Makmur (Sukoharjo, Central Java, Indonesia). The microalgae were grown in fresh water. The experiment was conducted according to the standard procedures of rearing and treating of farm animals as stated in law of the Republic of Indonesia, number 18, 2009, concerning animal husbandry and health.

Table 1 Chemical composition of *Spirulina platensis* (as-dry basis)

Items	Composition (%)
Moisture	91.8
Crude ash	11.9
Crude fat	0.63
Crude protein	52.4
Crude fibre	34.2

Table 2 Ingredients and nutrient composition (as-dry basis) of basal diet used in the study

Items	Composition (% , unless otherwise noted)
Maize	45.5
Soybean meal (CP 46%)	17.0
Wheat flour	10.0
Bread flour	5.00
Rice bran	4.45
Crude palm oil	3.50
Corn gluten meal (CP 62%)	3.60
Distiller dried grains (CP 27%)	3.00
Meat bone meal (CP 49%)	2.80
Chicken feather meal (CP 79%)	2.00
Bone meal (CP 22%)	1.50
Lysine	0.55
Methionine	0.37
L-threonine	0.08
Salt	0.15
Premix ¹	0.50
Analysed composition:	
Metabolizable energy (kcal/kg) ²	3,300
Dry matter	89.6
Crude protein	21.9
Crude fat	6.40
Crude fiber	5.62
Ash	6.39

¹Mineral-vitamin premix per kg of diet: C 250 g, P 0.625 g, Fe 3.570 mg, Cu 0.640 mg, Mn 5.285 mg, Zn 0.003 mg, Co 0.001 mg, Se 0.013 mg, I 0.016 mg, vit A 375 IU, vit D 150 IU and vit E 0.027 mg

²Metabolizable energy was calculated according to formula (Bolton, 1967): $40.81 \{0.87 (\text{crude protein} + 2.25 \text{ crude fat} + \text{nitrogen-free extract}) + 2.5\}$

CP: crude protein

The chicks were vaccinated with commercial Newcastle disease virus (NDV) vaccine through eye drops and drinking water at days 4 and 18 of the experiment, respectively. Bodyweight and feed intake were determined weekly. The FCR was determined as the feed intake per weight gain. To determine haematological profile, blood was obtained from the wing veins and collected in vacutainers containing ethylenediaminetetraacetic acid (EDTA) at day 32. The rest of the blood was collected in the vacutainers with no anticoagulant, let to clot at room temperature, and centrifuged at 2000 rpm for 15 minutes to produce serum. The serum was frozen until analyses of antibody titres and serum biochemistry (Sugiharto *et al.*, 2017a). At day 35, a total of 24 chicks were slaughtered, de-feathered and eviscerated. The internal organs were immediately taken out and weighed. Digesta were collected from the ileum and caecum of broilers to determine pH and for microbiological analyses.

Complete blood counts were determined using a hematology analyser (Prima Fully Auto Hematology Analyser, PT. Prima Alkesindo Nusantara, Jakarta, Indonesia). The NDV antibody titers of serum were determined based on haemagglutination inhibition (HI) assay (Villegas, 1987). The titers were presented as geometric mean titers (\log_2). Total triglyceride, total cholesterol, high-density lipoprotein (HDL) and low-density lipoprotein (LDL) cholesterol and uric acid serum were measured with the enzymatic colorimetric/colour method. Total protein, albumin, alanine aminotransferase (ALT), and aspartate aminotransferase (AST) in serum were determined with spectrophotometric/photometric tests. Data of globulin were obtained from the difference between the values of total protein and albumin in serum. The counts of certain bacteria in the intestinal digesta of broilers were determined according to Sugiharto *et al.*

(2017b) with few modifications. For enumeration of coliform bacteria and lactose-negative enterobacteria, samples were cultured on MacConkey agar. After aerobic incubation at 38 °C for 24 h, coliform and lactose-negative enterobacteria were counted as red and colourless colonies, respectively. The sum of coliform bacteria and lactose-negative enterobacteria was defined as enterobacteria. Lactic acid bacteria (LAB) were determined on de Man, Rogosa and Sharpe (MRS) agar after anaerobic incubation at 38 °C for 48 h.

Data were analysed based on a completely randomized design by ANOVA using the General Linear Models Procedure in SAS (SAS Inst. Inc., Cary, NC, USA). Pen was treated as the experimental unit. Significant differences among treatment groups were further analysed using Duncan's multiple-range test. A significant level of $P < 0.05$ was implemented.

Results

Data on the performances of broilers are presented in Table 3. There was a tendency for CONT birds to have higher ($P = 0.05$) weight gain for the first week of rearing compared with other birds. On days 22 to 35, FCR was lower ($P < 0.05$) in SP-21 birds than in CONT and SP-7 birds, but not different when compared with that in SP-35 birds. For the entire rearing period (days 0 to 35), no difference ($P > 0.05$) was observed with regard to weight gain, feed intake and FCR of broiler chicks.

Table 3 Effect of feeding duration of *Spirulina platensis* on performances of broiler chicks (means and SE)

Items (g)	Treatments				SE	P value
	CONT	SP-7	SP-21	SP-35		
Weight gain						
Days 0–7	135	118	112	122	5.54	0.05
Days 8–21	635	636	642	632	11.3	0.93
Days 22–35	1,000	984	1,048	1,018	26.5	0.37
Days 0–35	1,770	1,737	1,803	1,771	35.2	0.64
Feed intake						
Days 0–7	151	148	145	151	2.26	0.20
Days 8–21	970	968	976	970	9.31	0.95
Days 22–35	1,712	1,688	1,665	1,689	32.6	0.79
Days 0–35	2,833	2,804	2,786	2,811	33.9	0.80
FCR						
Days 0–7	1.14	1.26	1.29	1.25	0.04	0.09
Days 8–21	1.53	1.52	1.52	1.54	0.02	0.90
Days 22–35	1.72 ^a	1.72 ^a	1.60 ^b	1.66 ^{ab}	0.03	0.04
Days 0–35	1.60	1.62	1.55	1.59	0.02	0.19

^{a,b} Means with different superscripts in each row are significantly different

CONT: birds receiving basal diet with 0.04% of zinc bacitracin, SP-7: birds receiving 1% of *S. platensis* for the first seven days, SP-21: birds receiving 1% of *S. platensis* for the first 21 days, SP-35: birds receiving 1% of *S. platensis* for 35 days, FCR: feed conversion ratio, SE: standard error

Table 4 shows the data on internal organs of broiler chicks. The relative weight of the caecum was higher ($P < 0.05$) in SP-35 than CONT and SP-21 birds, but the difference was not substantial when compared with SP-7 birds. There are no significant differences in other internal organs of birds.

Data on haematological parameters of broiler chicks are presented in Table 5. The values of haemoglobin, erythrocytes and haematocrit were lower ($P < 0.05$) in SP-35 than in other birds. The numbers of leukocytes and lymphocytes were lower ($P < 0.05$) in SP-35 than in CONT birds, but the difference was not pronounced when compared with that in SP-7 and SP-21 birds. Concomitantly, the number of eosinophils tended ($P = 0.07$) to be lower in SP-35 than in CONT birds. The serum biochemical parameters and antibody titer against NDV were not different ($P > 0.05$) across the treatment groups.

Table 4 Effect of feeding duration of *Spirulina platensis* on internal organs of broiler chicks (means and SE)

Items (% live body weight)	Treatments				SE	P value
	CONT	SP-7	SP-21	SP-35		
Heart	0.51	0.51	0.46	0.45	0.03	0.28
Liver	2.41	2.61	2.43	2.40	0.16	0.77
Proventriculus	0.46	0.48	0.47	0.46	0.02	0.92
Gizzard	1.31	1.16	1.13	1.29	0.06	0.11
Pancreas	0.28	0.29	0.25	0.26	0.02	0.36
Duodenum	0.51	0.52	0.47	0.48	0.03	0.68
Jejunum	0.98	1.01	1.09	1.04	0.08	0.79
Ileum	0.68	0.79	0.74	0.76	0.09	0.84
Caecum	0.26 ^b	0.27 ^{ab}	0.24 ^b	0.32 ^a	0.02	0.03
Spleen	0.10	0.12	0.13	0.13	0.03	0.82
Thymus	0.38	0.42	0.30	0.34	0.05	0.28
Bursa of Fabricius	0.14	0.16	0.12	0.17	0.02	0.34

^{a,b} Means with different superscripts in each row are significantly different

CONT: birds receiving basal diet with 0.04% of zinc bacitracin, SP-7: birds receiving 1% of *S. platensis* for the first seven days; SP-21: birds receiving 1% of *S. platensis* for the first 21 days; SP-35: birds receiving 1% of *S. platensis* for 35 days; SE: standard error

The numbers of coliform bacteria were lower ($P < 0.05$) in the ileal digesta of SP-21 than in CONT and SP-7 birds, but the difference was not significant when compared with that in SP-35 birds (Table 6). In accordance, there was a tendency ($P = 0.08$) for the lower number of coliform bacteria in caecal digesta of SP-21 than in other birds. There were no differences ($P > 0.05$) in the numbers of lactose negative-enterobacteria, enterobacteria and LAB both in the ileum and caecum of broiler chicks.

Data on carcass traits of broiler chicks are presented in Table 7. In general, there was no difference ($P > 0.05$) in the carcass traits of broilers.

Discussion

A number of studies have revealed consistent benefits of *S. platensis* on the growth and health performances of broiler chickens (Qureshi *et al.*, 1996; Raju *et al.*, 2004; Kaoud, 2012; Jamil *et al.*, 2015; Shanmugapriya *et al.*, 2015a; Lokapinasari *et al.*, 2016; Yusuf *et al.*, 2016). Based on these published data, the authors inferred that *S. platensis* could be an alternative to in-feed antibiotics for broiler chickens. With the focus on replacing synthetic antibiotics in feed, the periods during which *S. platensis* was administered in broiler feeds were compared with solely antibiotics-supplemented feed. Hence, the authors did not provide an unsupplemented (neither with antibiotics nor *S. platensis*) feed group (as negative control) in the present study. The present results showed that the period during which *S. platensis* was supplemented in broiler feed did not affect growth performance of broilers. This finding was similar to that of Pietras (2001), which showed no effect of feeding duration of probiotic *Lactobacillus acidophilus* and *Streptococcus faecium* on production parameters of broiler chickens. Irrespective of feeding duration, dietary supplementation with 1% of *S. platensis* resulted in a corresponding effect on the growth performance when compared with feeding zinc bacitracin as a growth promoter to broilers. Indeed, inclusion of 1% of *S. platensis* in the diets for the first 21 days improved FC of broilers during days 22 to 35. Taken together, the current data may confirm the efficacy of *S. platensis* as an alternative to antibiotics in broiler diets.

In this study, feeding *S. platensis* for the entire rearing period enhanced the relative weight of caecum of broilers. The heavier caecum may be associated with the improved fermentation function of broiler chicks (Clench & Mathias, 1995). The definite explanation for the enhanced weight of caecum in broilers fed *S. platensis* for 35 days was not known, but it was possible that long-term feeding of *S. platensis* led to greater fibre intake, and thus resulted in heavier caeca (Moen *et al.*, 2016). Note that *S. platensis* in this study contained 34.2% of fibre.

Table 5 Effect of feeding duration of *Spirulina platensis* on haematological parameters of broiler chicks (means and SE)

Items	Treatments				SE	P value
	CONT	SP-7	SP-21	SP-35		
Complete blood counts						
Haemoglobin (g/dL)	9.78 ^a	10.2 ^a	10.3 ^a	7.78 ^b	0.57	0.02
Erythrocytes (10 ⁶ /μL)	2.30 ^a	2.30 ^a	2.39 ^a	1.82 ^b	0.11	0.01
Hematocrit (%)	30.2 ^a	31.0 ^a	31.9 ^a	24.8 ^b	1.59	0.02
MCV (fl)	131	135	134	138	1.51	0.11
MCH (pg)	42.6	44.2	42.8	41.9	1.28	0.63
MCHC (g/dL)	32.5	32.8	32.0	30.7	0.93	0.39
Leukocytes (10 ³ /μL)	22.2 ^a	17.1 ^{ab}	16.8 ^{ab}	13.4 ^b	2.04	0.04
Heterophils (10 ³ /μL)	0.63	0.55	0.47	0.42	0.12	0.58
Eosinophils (10 ³ /μL)	0.92	0.65	0.73	0.57	0.09	0.07
Lymphocytes (10 ³ /μL)	20.7 ^a	15.9 ^{ab}	15.6 ^{ab}	12.5 ^b	1.89	0.04
Biochemical parameters						
Total cholesterol (mg/dL)	128	146	150	139	7.65	0.23
HDL (mg/dL)	64.2	54.5	57.2	64.5	11.3	0.88
LDL (mg/dL)	46.5	67.2	75.1	61.6	11.7	0.39
Total triglyceride (mg/dL)	73.2	148	95.8	127	37.6	0.48
AST (U/L)	204	236	247	254	25.2	0.52
ALT (U/L)	7.63	6.52	3.00	7.57	1.82	0.25
Total protein (g/dL)	2.76	2.90	2.90	3.13	0.13	0.28
Albumin (g/dL)	1.25	1.22	1.32	1.25	0.05	0.60
Globulin (g/dL)	1.56	1.68	1.58	1.88	0.13	0.29
A/G ratio	0.81	0.73	0.86	0.70	0.06	0.22
Uric acid (mg/dL)	5.73	5.99	8.05	7.29	1.62	0.68
Antibody titer against NDV (Log ₂ GMT)	3.50	3.00	3.00	4.67	0.57	0.16

^{a,b} Means with different superscripts in each row are significantly different

CONT: birds receiving basal diet with 0.04% of zinc bacitracin; SP-7: birds receiving 1% of *S. platensis* for the first seven days; SP-21: birds receiving 1% of *S. platensis* for the first 21 days; SP-35: birds receiving 1% of *S. platensis* for 35 days; MCV: mean corpuscular volume; MCH: mean corpuscular hemoglobin; MCHC: mean corpuscular hemoglobin concentration; HDL: high-density lipoprotein; LDL: low-density lipoprotein; AST: aspartate aminotransferase; ALT: alanine aminotransferase; A/G ratio: albumin to globulin ratio; NDV: Newcastle disease virus; GMT: geometric mean titer; SE: standard error

Dietary supplementation of *S. platensis* has been reported to increase the number of erythrocytes and haemoglobin in broiler chicks (Jamil *et al.*, 2015). In contrast, feeding such algae during the whole rearing period resulted in lower values of erythrocytes, haemoglobin and haematocrit of broilers in the current study. To date, the cause of this decrease is not known, but long-term exposure of broilers to *S. platensis* may have a negative effect on the liver, where erythrocytes and haemoglobin are partly produced. Indeed, Iwasa *et al.* (2002) showed the presence of hepatotoxin (microcystin) in *Spirulina*, which may cause liver disruptions. Concomitantly, Roy-Lachapelle *et al.* (2017) found cyanotoxins in *Spirulina* that may induce liver problems. However, the authors' inference should be interpreted with caution as there was no significant impact of various feeding durations of *S. platensis* on the indicators of liver disruptions (e.g. AST, ALT and the relative liver weight). In parallel with the erythrocyte profile, the numbers of leukocytes, lymphocytes and eosinophils were lower in birds supplemented with *S. platensis* during the rearing period. These findings were different from those reported by Jamil *et al.* (2015) and Lokapinasari *et al.* (2016), which showed increased leukocytes with feeding *S. platensis* in broiler chicks. Again, the liver disruptions owing to long-term exposure to toxins in *S. platensis* seemed to contribute to the lower leukocytes and the differential leukocytes of

broilers in the present study. Interesting data were seen in the study of Shanmugapriya *et al.* (2015a), at which feeding 1.5% of *Spirulina* resulted in lower final bodyweight when compared with feeding 0.5 or 1% of *S. platensis*. Perhaps, excessive intake of *S. platensis* resulted in metabolic disturbances and affected the liver function leading to retarded growth rate in broilers. Considering the contradictory results between our study and others negatively, the differences in nutritional and functional properties of *S. platensis* used in the studies may to some extent be responsible. Note that *S. platensis* used in the current study was grown in fresh water, not in sea water as is the common commercial *S. platensis* available in the market.

Table 6 Effect of feeding duration of *Spirulina platensis* on intestinal microbial populations of broiler chicks (means and SE)

Items (log cfu/g)	Treatments				SE	P value
	CONT	SP-7	SP-21	SP-35		
Ileum						
Coliform	6.80 ^a	6.74 ^a	6.09 ^b	6.51 ^{ab}	0.16	0.03
Lactose negative-enterobacteria	1.56	1.64	2.17	1.67	1.02	0.97
Enterobacteria	8.36	8.37	8.26	8.18	1.03	0.99
LAB	8.77	8.87	8.75	8.73	0.19	0.96
Caecum						
Coliform	6.32	6.21	5.79	6.44	0.19	0.08
Lactose negative-enterobacteria	4.59	4.20	4.33	4.83	0.64	0.90
Enterobacteria	10.9	10.4	10.1	11.3	0.61	0.56
LAB	8.73	8.78	8.66	8.84	0.13	0.80

^{a,b} Means with different superscripts in each row are significantly different

CONT: birds receiving basal diet with 0.04% of zinc bacitracin; SP-7: birds receiving 1% of *S. platensis* for the first seven days; SP-21: birds receiving 1% of *S. platensis* for the first 21 days; SP-35: birds receiving 1% of *S. platensis* for 35 days; LAB: lactic acid bacteria; SE: standard error

Table 7 Effect of feeding duration of *Spirulina platensis* on carcass traits of broiler chicks (means and SE)

Items	Treatments				SE	P value
	CONT	SP-7	SP-21	SP-35		
(% live weight)						
Giblet ¹	4.23	4.29	4.02	4.14	0.19	0.76
Eviscerated carcass	67.9	69.4	69.3	68.7	0.62	0.28
(% eviscerated carcass)						
Breast	36.3	34.3	35.7	35.6	0.81	0.40
Thigh	15.9	15.8	16.3	15.5	0.52	0.77
Drumstick	13.5	13.1	13.4	13.8	0.45	0.78
Wing	10.7	10.9	10.6	11.0	0.41	0.88
Abdominal fat	2.43	2.22	2.46	1.99	0.26	0.56

¹Giblet: heart, gizzard and liver

CONT: birds receiving basal diet with 0.04% of zinc bacitracin; SP-7: birds receiving 1% of *S. platensis* for the first seven days; SP-21: birds receiving 1% of *S. platensis* for the first 21 days; SP-35: birds receiving 1% of *S. platensis* for 35 days; SE: standard error

In the present study, feeding 1% of *S. platensis*, especially during the first 21 days, decreased the numbers of coliform in the ileal and caecal digesta of broiler chicks. This finding was in accordance with that

reported by Shanmugapriya *et al.* (2015b), which showed the decreased *E. coli* population in the ileal and caecal digesta of broilers when feeding 1% of *S. platensis*. In the study of Yusuf *et al.* (2016), it was apparent that feeding 2% of *S. platensis* decreased the numbers of coliform bacteria in the ileocecal contents of Japanese quails fed a vegetarian protein diet. Concomitantly, Nuhu (2013) showed the bacterial clearance capacities of *S. platensis* in chicks injected with *E. coli* or *Staphylococcus aureus*. In such cases, the antimicrobial activities (against pathogenic bacteria) of *S. platensis* seemed to be responsible. In their *in vitro* study, Mala *et al.* (2009) showed that extract of *S. platensis* was able to inhibit the growth of *Klebsiella pneumoniae*, *Shigella shigae*, *E. coli*, *S. aureus*, *Proteus vulgaris*, *Pseudomonas aeruginosa* and *Salmonella typhi*. Similar to this, Kaushik & Chauhan (2008) showed the antibacterial activities of *S. platensis* against *S. aureus* and *E. coli*, while El-Baz *et al.* (2013) showed the antibacterial activities of *S. platensis* extract against *E. coli*, *S. aureus*, *S. typhi*, and *Enterococcus faecalis*. Indeed, several compounds, including γ -linolenic acid, active fatty acid lauric and palmitoleic acid, have been attributed to the antimicrobial activities of *S. platensis* (El-Sheekh *et al.*, 2014). With its prebiotic properties (Beheshtipour *et al.*, 2013; de Jesus Raposo *et al.*, 2016), *S. platensis* is known to possess a stimulating effect on the growth of LAB (*Lactobacillus acidophilus*, *Lactobacillus casei* and *Streptococcus thermophilus*) (Bhowmik *et al.*, 2009). In broiler chickens, *S. platensis* administration has been associated with the increased intestinal population of LAB (Shanmugapriya *et al.*, 2015b), while in Japanese quails, feeding such algae tended to increase the count of intestinal lactobacilli (Yusuf *et al.*, 2016). Unlike these earlier studies, treatment with *S. platensis* in the present study did not affect the ileal and caecal populations of LAB. To date, the explanation for these divergent data remains unclear, but the different levels and the nutritional qualities of *S. platensis* as well as the trial conditions may be responsible.

Several studies reported that feeding *S. platensis* increased carcass percentage of broiler chicks (Raju *et al.*, 2004; Kaoud, 2012; Marley *et al.*, 2014) and Japanese quails (Jamil *et al.*, 2015). In contrast, the present findings did not show any effect of *S. platensis* (regardless of feeding duration) on carcass traits of broilers. The data of the current study were concomitant with those reported by Cheong *et al.* (2016), which showed no significant effect of feeding up to 8% of *S. platensis* on the carcass, breast and legs percentage of Japanese quails. No substantial effect on the carcass weight was seen in rabbit, as reported by Mahmoud *et al.* (2017) when including *S. platensis* to replace soybean in the diets of rabbits (with percentage of 20, 40 and 60%). The precise reason for these different results is not known, but the relatively similar final bodyweight may result in the similar carcass percentage of broilers among the treatment groups in the present study. This inference was supported by Marley *et al.* (2014), who suggested that carcass weight was in parallel with the live body weight of broilers. That is, the increased carcass weight and total edible part were attributed to the increased final live bodyweight of broilers, and vice versa.

Conclusion

From the present study, it can be concluded that the administration of *S. platensis* for the first 21 days of broiler's lives resulted responses that were similar to or better than administration of *S. platensis* or antibiotics throughout the entire rearing period. It is therefore not necessary to administer *S. platensis* throughout the rearing period to obtain antibiotic-like benefits for broilers.

64

Acknowledgements

The study was funded by the Faculty of Animal and Agricultural Sciences, Diponegoro University through the Faculty Research Grant No: 84/UN7.5.5/PP/2017, 18 May 2017

Authors' Contributions

SS designed the study, carried out the animal experiment, data analysis and manuscript drafting. II and EW carried out the animal experiment. TY conducted the laboratory analysis.

20

Conflict of Interest Declaration

The authors declare that they have no competing interests.

References

- Beheshtipour, H., Mortazavian, A.M., Mohammadi, R., Sohrabvandi, S. & Khosravi-Darani, K., 2013. Supplementation of *Spirulina platensis* and *Chlorella vulgaris* algae into probiotic fermented milks. *Compr. Rev. Food Sci. Food Saf.* 12, 144-154.
- Bhowmik, D., Dubey, J. & Mehra, S., 2009. Probiotic efficiency of *Spirulina platensis*-stimulating growth of lactic acid bacteria. *American-Eurasian J. Agric. & Environ. Sci.* 6, 546-549.
- Bolton, W., 1967. Poultry nutrition. MAFF Bulletin No.174. HMSO, London.
- Bonos, E., Kasapidou, E., Kargopoulos, A., Karampampas, A., Christaki, E., Florou-Paneri, P. & Nikolakakis, I., 2016. *Spirulina* as a functional ingredient in broiler chicken diets. *S. Afr. J. Anim. Sci.* 46, 94-102.

- Cheong, D.S.W., Kasim, A., Sazili, A.Q., Omar, H. & Teoh, J.Y., 2016. Effect of supplementing *Spirulina* on live performance, carcass composition and meat quality of Japanese quail. *Walailak J. Sci. Tech.* 13, 77-84.
- Clench, M.H. & Mathias, J.R., 1995. The avian cecum: a review. *Wilson Bull.* 107, 93-121.
- de Jesus Raposo, M.F., de Morais, A.M.M.B. & de Morais, R.M.S.C., 2016. Emergent sources of prebiotics: seaweeds and microalgae. *Mar. Drugs*. 14, 27; doi:10.3390/md14020027
- El-Baz, F.K., El-Senousy, W.M., El-Sayed, A.B. & Kamel, M.M., 2013. In vitro antiviral and antimicrobial activities of *Spirulina platensis* extract. *J. Appl. Pharm. Sci.* 3, 52-56.
- El-Sheekh, M.M., Daboor, S.M., Swelim, M.A. & Mohamed, S., 2014. Production and characterization of antimicrobial active substance from *Spirulina platensis*. *Iran. J. Microbiol.* 6, 112-119.
- Farag, M.R., Alagawany, M., El-Hack, M.E.A. & Dhama, K., 2016. Nutritional and healthical aspects of *Spirulina* (*Arthrospira*) for poultry, animals and human. *Int. J. Pharmacol.* 12, 36-51.
- Holman B.W.B. & Malau-Aduli, A.E.O., 2013. *Spirulina* as a livestock supplement and animal feed. *J. Anim. Physiol. Anim. Nutr.* 97, 615-623.
- Iwasa, M., Yamamoto, M., Tanaka, Y., Kaito, M. & Adachi, Y., 2002. *Spirulina* associated hepatotoxicity. *Am. J. Gastroenterol.* 97, 3212-3213.
- Jamil, A.B.M.R., Akanda, R., Rahman, M., Hossain, A. & Islam, S., 2015. Prebiotic competence of spirulina on the production performance of broiler chickens. *J. Adv. Vet. Anim. Res.* 2, 304-309.
- Kaoud, H.A., 2012. Effect of *Spirulina platensis* as a dietary supplement on broiler performance in comparison with prebiotics. *Sci. J. App. Res.* 1, 44-48.
- Kaushik, P. & Chauhan, A., 2008. In vitro antibacterial activity of laboratory grown culture of *Spirulina platensis*. *Indian J. Microbiol.* 48, 348-352.
- Lokapimasari, W.P., Yulianto, A.B., Legowo, D. & Agustono, 2016. The effect of *Spirulina* as feed additive to myocardial necrosis and leukocyte of chicken with avian influenza (H5N1) virus infection. *Proc. Chem.* 18, 213-217.
- Mahmoud, A.E., Naguib, M.M., Higazy, A.M., Sultan, Y.Y. & Marrez, D.A., 2017. Effect of substitution soybean by blue green alga *Spirulina platensis* on performance and meat quality of growing rabbits. *Am. J. Food Technol.* 12, 51-59.
- Mala, R., Sarojini, M., Saravanababu, S. & Umadevi, G., 2009. Screening for antimicrobial activity of crude extracts of *Spirulina platensis*. *J. Cell Tissue Res.* 9, 1951-1955.
- Mariey, Y.A., Samak, H.R., Abou-Khashba, H.A., Sayed, M.A.M. & Abou-Zeid, A.E., 2014. Effect of using *Spirulina platensis* algae as a feed additives for poultry diets: 2–Productive performance of broiler. *Egypt. Poult. Sci.* 34, 245-258.
- Moen, B., Henjum, K., Måge, I., Knutsen, S.H., Rud, I., Hetland, R.B. & Paulsen, J.E., 2016. Effect of dietary fibers on cecal microbiota and intestinal tumorigenesis in azoxymethane treated A/J Min/+ mice. *PLoS ONE.* 11, e0155402. <https://doi.org/10.1371/journal.pone.0155402>
- Nuhu, A.A., 2013. *Spirulina* (*Arthrospira*): An important source of nutritional and medicinal compounds. *J. Marine Biol.* <http://dx.doi.org/10.1155/2013/325636>
- Pietras, M., 2001. The effect of probiotics on selected blood and meat parameters of broiler chickens. *J. Anim. Feed Sci.* 10, 297-302.
- Pourhossein, Z., Qotbi, A.A.A., Seidavi, A., Laudadio, V., Centoducati, G. & Tufarelli, V., 2015. Effect of different levels of dietary sweet orange (*Citrus sinensis*) peel extract on humoral immune system responses in broiler chickens. *Anim. Sci. J.* 86, 105-110.
- Qureshi, M.A., Garlich, J.D. & Kidd, M.T., 1996. Dietary *Spirulina platensis* enhances humoral and cell-mediated immune functions in chickens. *Immunopharmacol. Immunotoxicol.* 18, 465-476.
- Raju, M.V.L.N., Rama Rao, S.V., Radhika, K. & Chawak, M.M., 2004. Effects of *Spirulina platensis* or furazolidone on the performance and immune response of broiler chickens fed with aflatoxin contaminated diet. *Indian J. Anim. Nutr.* 21, 40-44.
- Roy-Lachapelle, A., Sollicec, M., Bouchard, M.F. & Sauvé, S., 2017. Detection of cyanotoxins in algae dietary supplements. *Toxins.* 9, 76; doi:10.3390/toxins9030076
- SAS Institute, Inc., 1985. SAS user's guide: Statistics, version 5. Cary, NC, USA.
- Shanmugapriya, B., Babu, S.S., Hariharan, T., Sivaneswaran, S., Anusha, M.B. & Raja, P.U., 2015a. Synergistic effect of *Spirulina platensis* on performance and gut microbial load of broiler chicks. *Indo-Asian J. Multidisc. Res.* 1, 149-155.
- Shanmugapriya, B., Babu, S.S., Hariharan, T., Sivaneswaran, S. & Anusha, M.B., 2015b. Dietary administration of *Spirulina platensis* as probiotics on health and histopathology in broiler chicks. *Int. J. Recent Sci. Res.* 6, 2650-2653.
- SNI (Indonesian National Standard), 2006. Standard for broiler feed (SNI 01-3930-2006). National Standardization Agency of Indonesia, Jakarta, Indonesia (article in Bahasa).
- Sugiharto, S., 2016. Role of nutraceuticals in gut health and growth performance of poultry. *J. Saudi Soc. Agric. Sci.* 15, 99-111.
- Sugiharto, S. & Lauridsen, C., 2016. Dietary *Chlorella* supplementation effect on immune responses and growth performances of broiler chickens exposed to post hatch holding time. *Livest. Res. Rur. Dev.* 28; <http://www.lrrd.org/lrrd28/7/sugi28119.html>
- Sugiharto, S., Yudiarti, T., Isroli, I., Widiastuti, E. & Putra, F.D., 2017a. Intestinal microbial ecology and hematological parameters of broiler fed cassava waste pulp fermented with *Acremonium charticola*. *Vet. World.* 10, 324-330.

- Sugiharto, S., Yudiarti, T., Isroli, I., Widiastuti, E. & Putra, F.D., 2017b. Effect of dietary supplementation with *Rhizopus oryzae* or *Chrysonilia crassa* on growth performance, blood profile, intestinal microbial population, and carcass traits in broilers exposed to heat stress. *Arch. Anim. Breed.*, 60, 347–356.
- Villegas, P., 1987. Avian virus diseases laboratory manual. College of Veterinary Medicine. University of Georgia, Athens, Georgia, USA.
- Yusuf, M.S., Hassan, M.A., Abdel-Daim, M.M., El Nabtiti, A.S., Ahmed, A.M., Moawed, S.A., El-Sayed, A.K. & Cui, H., 2016. Value added by *Spirulina platensis* in two different diets on growth performance, gut microbiota, and meat quality of Japanese quails. *Vet. World.* 9, 1287-1293.

ORIGINALITY REPORT

18%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|----------|---|-----------|
| 1 | Submitted to Universidad San Francisco de Quito
Student Paper | 1% |
| <hr/> | | |
| 2 | Brand, TS, and L Brundyn. "Effect of supplementary feeding to ewes and suckling lambs on ewe and lamb live weights while grazing wheat stubble", South African Journal of Animal Science, 2015.
Publication | 1% |
| <hr/> | | |
| 3 | L. L. Mikkelsen. "Effects of Physical Properties of Feed on Microbial Ecology and Survival of Salmonella enterica Serovar Typhimurium in the Pig Gastrointestinal Tract", Applied and Environmental Microbiology, 06/01/2004
Publication | 1% |
| <hr/> | | |
| 4 | Absalan, M, A Afzalzade, M Mirzaee, SD Sharifi, M Khorvash, and M Kazemi-Benchenari. "Feeding of whole cottonseed on performance, carcass characteristics and intestinal morphology of Zandi lambs", South African Journal of Animal Science, 2011. | 1% |

5 www.i-scholar.in 1%

Internet Source

6 V Jazi, A D Foroozandeh, M Toghyani, B Dastar, R Rezaie Koochaksaraie, M Toghyani. 1%

"Effects of *Pediococcus acidilactici*, mannan-oligosaccharide, butyric acid and their combination on growth performance and intestinal health in young broiler chickens challenged with *Salmonella Typhimurium*", *Poultry Science*, 2018

Publication

7 Niu, ZY, YN Min, HY Wang, J Zhang, WC Li, L Li, and FZ Liu. "Effects of dietary dihydropyridine 1%

on laying performance and lipid metabolism of broiler breeder hens", *South African Journal of Animal Science*, 2011.

Publication

8 scialert.net 1%

Internet Source

9 pt.scribd.com <1%

Internet Source

10 scienceandnature.org <1%

Internet Source

11 www.cambridge.org <1%

Internet Source

- | | | |
|----|--|-----|
| 12 | Mohamed M. Abdel-Daim. "Pharmacodynamic interaction of Spirulina platensis with erythromycin in Egyptian Baladi bucks (Capra hircus)", Small Ruminant Research, 2014
Publication | <1% |
| 13 | Daneshmand, A., GH. Sadeghi, A. Karimi, A. Vaziry, and S.A. Ibrahim. "Evaluating complementary effects of ethanol extract of propolis with the probiotic on growth performance, immune response and serum metabolites in male broiler chickens", Livestock Science, 2015.
Publication | <1% |
| 14 | academic.oup.com
Internet Source | <1% |
| 15 | T. K. Lim. "Edible Medicinal And Non-Medicinal Plants", Springer Nature, 2013
Publication | <1% |
| 16 | www.frontiersin.org
Internet Source | <1% |
| 17 | translational-medicine.biomedcentral.com
Internet Source | <1% |
| 18 | www.dfrc.ars.usda.gov
Internet Source | <1% |
| 19 | banglajol.info
Internet Source | <1% |

- | | | |
|----|---|-----|
| 20 | Yoshihisa HIRAKAWA. "Evaluation of gender differences in caregiver burden in home care: Nagoya Longitudinal Study of the Frail Elderly (NLS-FE)", Psychogeriatrics, 9/2006
Publication | <1% |
| 21 | www.emisphere.com
Internet Source | <1% |
| 22 | A. Daneshmand, GH. Sadeghi, A. Karimi, A. Vaziry. "Effect of oyster mushroom (<i>Pleurotus ostreatus</i>) with and without probiotic on growth performance and some blood parameters of male broilers", Animal Feed Science and Technology, 2011
Publication | <1% |
| 23 | world-rabbit-science.com
Internet Source | <1% |
| 24 | Submitted to Universitas Diponegoro
Student Paper | <1% |
| 25 | repositorio.ufpe.br
Internet Source | <1% |
| 26 | jas.fass.org
Internet Source | <1% |
| 27 | www.animalproduction.net
Internet Source | <1% |
| 28 | Mburu, M.. "PROXIMATE COMPOSITION AND | |

NUTRITIONAL CHARACTERIZATION OF CHIA ENRICHED YOGHURT.(Report)", African Journal of Food, Agriculture, Nutrition and Development

Publication

<1%

29

spandidos-publications.com

Internet Source

<1%

30

Submitted to University of Stellenbosch, South Africa

Student Paper

<1%

31

Sugiharto Sugiharto, Turrini Yudiarti, Isroli Isroli, Endang Widiastuti, Endang Kusumanti. "Dietary supplementation of probiotics in poultry exposed to heat stress – a review", Annals of Animal Science, 2017

Publication

<1%

32

e-sciencecentral.org

Internet Source

<1%

33

ediss.uni-goettingen.de

Internet Source

<1%

34

www.wellnessresources.com

Internet Source

<1%

35

www.mdpi.com

Internet Source

<1%

36

Raffaella Campana, Veronica Martinelli, Stefano

<1%

Scoglio, Evelin Colombo, Serena Benedetti, Wally Baffone. "Influence of Aphanizomenon flos-aquae and two of its extracts on growth ability and antimicrobial properties of Lactobacillus acidophilus DDS-1", LWT - Food Science and Technology, 2017

Publication

37

Alzawqari, M. H., A. A. Al-Baddany, H. H. Al-Baadani, I. A. Alhidary, Rifat Ullah Khan, G. M. Aqil, and A. Abdurab. "Effect of feeding dried sweet orange (Citrus sinensis) peel and lemon grass (Cymbopogon citratus) leaves on growth performance, carcass traits, serum metabolites and antioxidant status in broiler during the finisher phase", Environmental Science and Pollution Research, 2016.

Publication

38

Kumar, V., A.V. Elangovan, A.B. Mandal, P.K. Tyagi, S.K. Bhanja, and B.B. Dash. "Effects of feeding raw or reconstituted high tannin red sorghum on nutrient utilisation and certain welfare parameters of broiler chickens", British Poultry Science, 2007.

Publication

39

A. de la Jara, C. Ruano-Rodriguez, M. Polifrone, P. Assunção, Y. Brito-Casillas, A. M. Wägner, L. Serra-Majem. "Impact of dietary Arthrospira (Spirulina) biomass consumption on human

<1%

<1%

<1%

health: main health targets and systematic review", Journal of Applied Phycology, 2018

Publication

40

ijpsr.com

Internet Source

<1%

41

M. AZIZI, A.R. SEIDAVI, M. RAGNI, V. LAUDADIO, V. TUFARELLI. "Practical applications of agricultural wastes in poultry feeding in Mediterranean and Middle East regions. Part 1: citrus, grape, pomegranate and apple wastes", World's Poultry Science Journal, 2018

Publication

<1%

42

preview-cardiab.biomedcentral.com

Internet Source

<1%

43

EIShenway, Abeer, and Mosaas Soltan. "Effect of Dietary Probiotic and/or Prebiotic Supplementation on Growth Performance, Carcass Traits and Some Serum Biochemical Alterations in Broiler Chicken", Journal of Animal Science Advances, 2015.

Publication

<1%

44

G.F. Castillo-Mitre, A. Olmedo-Juárez, R. Rojo-Rubio, M. González-Cortázar et al. "Caffeoyl and coumaroyl derivatives from Acacia cochliacantha exhibit ovicidal activity against Haemonchus contortus", Journal of

<1%

Ethnopharmacology, 2017

Publication

45	www.aasv.org Internet Source	<1%
46	unair.ac.id Internet Source	<1%
47	www.intaquares.com Internet Source	<1%
48	ijmbonline.com Internet Source	<1%
49	www.omicsonline.org Internet Source	<1%
50	espace.library.uq.edu.au Internet Source	<1%
51	"EFFECTS OF INCLUSION OF AERIEL DRIED PARTS OF SOME HERBS IN BROILER DIETS. (Report)", Journal of Animal and Plant Sciences, Sept 30 2011 Issue Publication	<1%
52	www.odermatol.com Internet Source	<1%
53	www.jarvm.com Internet Source	<1%
54	ijas.pagepress.org Internet Source	<1%

55 Lisa Karr-Lilienthal, N. R. Merchen, Christine Grieshop, Marianne Smeets-Peeters, G. C. Fahey. "Selected Gelling Agents in Canned Dog Food Affect Nutrient Digestibilities and Fecal Characteristics of Ileal Cannulated Dogs", *Archiv für Tierernaehrung*, 2002

Publication

<1%

56 www.tdx.cat

Internet Source

<1%

57 S. Khan, R. U. Khan, W. Alam, A. Sultan. "Evaluating the nutritive profile of three insect meals and their effects to replace soya bean in broiler diet", *Journal of Animal Physiology and Animal Nutrition*, 2018

Publication

<1%

58 Yasmina M. Abd El-Hakim, Wafaa A. Mohamed, Abeer E. El-Metwally. "Spirulina platensis attenuates furan reprotoxicity by regulating oxidative stress, inflammation, and apoptosis in testis of rats", *Ecotoxicology and Environmental Safety*, 2018

Publication

<1%

59 Christaki, Efterpi, Eleftherios Bonos, Ilias Giannenas, and Panagiota Florou-Paneri. "Aromatic Plants as a Source of Bioactive Compounds", *Agriculture*, 2012.

Publication

<1%

60

www.tandfonline.com

Internet Source

<1%

61

www.wildlifedisease.org

Internet Source

<1%

62

Engberg, R. M., M. S. Hedemann, T. D. Leser, and B. B. Jensen. "Effect of zinc bacitracin and salinomycin on intestinal microflora and performance of broilers", Poultry Science, 2000.

Publication

<1%

63

Engberg, R. M., M. S. Hedemann, S. Steinfeldt, and B. B. Jensen. "Influence of Whole Wheat and Xylanase on Broiler Performance and Microbial Composition and Activity in the Digestive Tract", Poultry Science, 2004.

Publication

<1%

64

Sugiharto, Sugiharto, Turrini Yudiarti, and Isroli Isroli. "Functional Properties of Filamentous Fungi Isolated from the Indonesian Fermented Dried Cassava, with Particular Application on Poultry", Mycobiology, 2015.

Publication

<1%

65

Naghi Shokri, Ali, Hossein A. Ghasemi, and Kamran Taherpour. "Evaluation of Aloe vera and synbiotic as antibiotic growth promoter substitutions on performance, gut morphology, immune responses and blood constituents of

<1%

broiler chickens : Aloe Vera and Synbiotic as Antibiotic Alternatives", Animal Science Journal, 2016.

Publication

66

Grinstead, G.S.. "Effects of Spirulina platensis on growth performance of weanling pigs", Animal Feed Science and Technology, 20000306

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On