

Nanoparticle of silver nitrate (Ag₂NO₃) and Carbamate (CO₂NH₃ derivatives) for vector control of *Aedes aegypti*

by Mursid Rahardjo

Submission date: 05-Dec-2019 02:38PM (UTC+0700)

Submission ID: 1227575150

File name: artikel.pdf (1.44M)

Word count: 3087

Character count: 16136

9

Nanoparticle of Silver Nitrate (Ag₂NO₃) and Carbamate (CO₂NH₃ derivatives) For Vector Control of *Aedes aegypti*

Mursid Raharjo^{1,*}, Agus Subagyo², and Sulistyani³

5

¹Magister of Environmental Health, Faculty of Public Health, Diponegoro University, Semarang - Indonesia

²Faculty of Sains and Mathematics, Diponegoro University, Semarang – Indonesia

³Magister of Environmental Health, Faculty of Public Health, Diponegoro University, Semarang - Indonesia

Abstract. The climate change has an impact on various lives, including in the vector breeding. The *Aedes aegypti* resistance as an infectious agent of dengue fever is harmful for life, especially to non-target organisms and the environment. Nano-silver (Ag₂NO₃), is a new form of engineering in vector control. The aim of this research was the toxicity test of nano-silver (Ag₂NO₃) and carbamate (CO₂NH₃ derivatives) on *Aedes aegypti* larvae. This was a laboratory research (true experiment), used for 3-4 instar stage larvae toxicity test. The number of samples was calculated using an experimental approach, with a completely randomized design. As simply, can formulated as follows, $(t-1)(r-1) > 15$. The amount of this samples were 14. The toxicity test with carbamate compounds, at concentration 0.01 mg / l as many as 80% larvae were death, whereas on concentrations 0.5 mg / l as many as 92% of the test animals were death. Toxicity test with nano-silver on concentration 0.3 mg / l, all of the test animals (100%) experienced were death. The results of the observations also showed that the instar more active in the low concentration of Ag₂NO₃ (0.01 mg/l - 0.1 mg/l). The conclusion are, nano-silver is more effective than carbamate compounds, and there is a potential for resistance.

Keywords : Nanosilver; Carbamate; *Aedes aegypti*.

1 Background

The climate change has an impact to various lives including the vector breeding [1]. The changes effecting to vectors especially in bionocyclic vectors, including *Aedes aegypti* as vector of infectious dengue fever [2]. Climate change also has the potential to caused the appearance of rare species [3].

The occurrence of *Aedes aegypti* resistance as a carrier of dengue fever is dangerous for life [4]. Resistance demanding the dose increase of insecticide use. The dose increase threaten the non target organism and harmful for the vector's environment [5].

Carbamate (CO₂NH₃ derivat) commonly use for vector control [6]. The result of another research show that 100% resistance on 0,03 mg/l concentration [7]. WHO classified the active materials of carbamate types carbofuran, butocarbocoxim dan butoxycarbocoxim to class lb (Highly hazardous) [8].

Nanotechnology with its growth has been expanding to some utilization [9]. In the health sector, there was some materials tried in another country with Nano-TiO₂, Sillica and herbs material [10]. Nanosilver which come from Ag₂NO₃ easy to obtain and can be used in the health sector. The biomecanism principle of carbamate and organofosfat which inhibit Acetylcholinesterase (AChE),

expected has a change when synthesized with nano-silver. The nano-silver particles, beside on theory able to penetrate Ache enzyme, so it more effective and could use in lower dose [4].

The problem on this research is how to diversify of Ag₂NO₃ nanoparticles for vector control compared to carbamate compounds. The aim of this research is to do toxicity test of nanoparticles (Ag₂NO₃) and carbamate (CO₂NH₃ derivat) to the *Aedes aegypti* larvae

2 Methods

This was laboratory research (true experiment) by controlling all of variables. The research started with nano-silver making which come from Ag₂NO₃. Ag₂NO₃ materials obtained from chemical supplier. Carbamate obtained from the markets product and standarized to determine the concentration. The *Aedes aegypti* vector which used for toxicity test was larvae instar 3-4. Toxicity test for larvae used the liquid-immersion method. Amount of the samples measured by experimental approach with complete randomized design. As simply, can be formulated as $(t-1)(r-1) > 15$. [11-12]. The amount of the samples was 14. The solutions used as test solutions were carbamate (Furadan 3gr) and Ag₂NO₃ nano-particle (0.2%). Steps of making the mother liquor (Furadan 3gr) to obtain two concentrations are 0,1 ppm and 1 ppm. The

4

* Corresponding author: arsikazk@gmail.com

Ag₂NO₃ nano-particle solution designed with 2 ppm concentration of mother liquor.

Toxicity test use the *Aedes aegypti* species instar 3-4 with carbamate and Ag₂NO₃ solutions. The steps of this test are 1) Transparent glasses are prepared about 14 pieces used for 2 types of solutions test with 6 concentrations and 1 control in each solution. 2) Observations were did in the first 2 hours then the observations were recorded. 3) Observations carried out again in the next 24 hours then recorded the results of these observations.

3 Results

3.1 Carbamate (CO₂NH₃ derivatives)

Carbamate toxicity test was carried out on 25 instars of *Aedes aegypti*, using controls. Observations are made in the first 2 hours and after 24 hours. In the first 2 hours, at concentration 0.01 mg / l as many as 56% larvae of *Aedes aegypti* was died. At the highest concentration at the test 0.5 mg/l, 84% of the larvae were died. There was percentage increase from 56% (0.01 mg / l) to 84% (0.5 mg / l).

After 24 hours observation of the larvae test, it was obtained as follows, at concentration 0.01 mg / l, 80% of the test animals experienced were death. The death increased by 92% at concentration 0,5 mg / l. The results

of the carbamate toxicity test on instar 3-4 of *Aedes aegypti* are presented in Table 1 and Figure 1.

3.2 Ag₂NO₃ Nanoparticles (Silver Nitrate) Test

Nano Silver nitrate particles (Ag₂NO₃) toxicity test were carried out on *Aedes aegypti* larvae instar 3-4. The total of 25 larvae were tested with various nano concentrations of silver nitrate particles. The observations are obtained as follows, in the first 2 hours observation, at 0.01 mg/l, no test animals were died. Likewise the 0.05 mg/l and 0.1 mg/l. The 0.3 mg/l concentration, there were 4% of animal's death, while in 0.4 mg / l and 0.5 mg / l concentration, all test animals (100%) experienced were death. At 0.01 mg/l, 0.05 mg/l and 0.1 mg / l concentration shows the test animals move more actively.

At 24 hours observation, obtained at concentrations 0.01 mg / l there were no (0%) test animals were death. The new death of test animals were seen at concentrations of 0.05 mg / l (4%) and 0.1 mg/l (4%). At concentrations 0.3 mg / l to 0.5 mg / l all of test animals (100%) experienced were death. The results of the observations also still show the occurrence of instar activity in the low concentration of Ag₂NO₃ which is 0.01 mg / l - 0.1 mg / l. The results of the toxicity test of Ag₂NO₃ nanoparticles on instar 3-4 of *Aedes aegypti* are presented in Table 1 and Figure 2.

Table 1. The result of Ag₂NO₃ and carbamate toxicity test on instar 3-4 *Aedes aegypti*

Concentration of Karbamat (mg/l)	2 hours	24 hours	Concentration of Ag ₂ NO ₃ (mg/l)	2 hours	24 hours
0,01	56%	80%	0,01	0%	0%
0,05	68%	80%	0,05	0%	4%
0,1	76%	88%	0,1	0%	4%
0,3	84%	88%	0,3	4%	100%
0,4	84%	92%	0,4	100%	100%
0,5	84%	92%	0,5	100%	100%




Fig. 1. The death percentage's development of instar 3-4 *Aedes aegypti* with carbamate (CO₂NH₃ derivatives)




Fig. 2. The death percentage's development of instar 3-4 *Aedes aegypti* with nano materials of Ag₂NO₃

3 Discussion

Chemical control by utilizing various synthetic material, processed, along with organophosphate and carbamate derivatives are still widely used. Control is carried out to the vector at the larvae-pupa stage until the mature stage. Oxygen and mineral that contains in the water are utilized by Instar stage of *Aedes aegypti* to live in the water. Oxygen absorption through the surface of the skin and minerals by mouth are the biochemical mechanism in the life cycle of larvae. Biochemical mechanisms of larvae's body are used to choose the type of intervention so that the larvae are easily destroyed and not given the chance to become a mosquito.

The results of the study of silver nanoparticle toxicity tests (Ag₂NO₃) shows, that at low concentrations (0,01 – 0,1 mg/lit) the higher activity of *Aedes aegypti* larvae occurs. This high activity shows the suitability of nitrate minerals as essential material for growth and development to boost larvae activity.

Acetylcholinesterase Enzym (AChE) can grow optimally. According to Liebig's theory that the growth of organisms is controlled by a limiting factor [5].

At a concentration of 0.3 mg / lt, all test animals (100%) were died. This concentration is smaller than carbamate. At the same concentration, carbamate can only kill 88% of test animals. Heeding to the characteristics of nanoparticles, it has size 1-100 nm, and has a high penetrating power to media around it.

At low concentration, the mortality rate of toxicity test with carbamate was higher than nano particle because of the active materials of carbamates is poisonous although in the low concentration, while the nitrate is not poisonous in the low concentration and can be the essential material for larva's growth. In the high concentration, the nitrate can be poisonous for the organism life, so thats why it can kill all of animal test and be more effective with its smaller size than carbamate, so it easily inhibit the larvae enzym. Nano-silver (Ag₂NO₃) material has smaller particle size than larvae's pores in the inter-abdominal segment. Silver nitrate compounds will react directly with the ACHe hormone system, so that it will directly affect the larvae's

metabolism. The inhibition of the enzyme's performance causes the death of larvae. Carbamate compounds with a larger diameter only possible to enter the larvae through the digestive tract which caused only a part of the compound reacts with Ache enzym (Acethyl Cholinesterase), so it can be understood why Nano-silver concentrations are lower than Carbamat (CO_2NH_3 derivatives) concentration.

The larvae breathe through the funnel on their tail, and enter the food through the gastrointestinal tract and through the abdominal pores [13]. Larval body pores are estimated [2] have a larger size than Ag_2NO_3 nano particles. Overall, nano refers to a size scale between 1 nanometer (nm) and 100 nm. In comparison, the wavelength of visible light is between 400 nm and 700 nm. A leukocyte has a size of 10,000 nm, a bacterium 1000-10000 nm, a virus 75-100 nm, a protein of 5-50 nm, deoxyribonucleic acid (DNA) ~ 2 nm (width), and atoms of ~ 0.1 nm. In this scale, the physical, biological and chemical characteristics of materials have fundamentally difference between one and another and often unexpected actions are showed from them [14].

Ag_2NO_3 nanoparticles are 1-100 nm in size. These particles have penetrating power on Aedes larvae. Nanoparticles are dispersions of particles or solid colloid structures that the diameter is start from 1 to 1000 nm. They consist of synthetic, semi-synthetic and natural polymer active therapeutic molecules that have the ability to be trapped, encapsulated, dissolved, absorbed, or chemically attached [15]. Due to biodegradability, biocompatibility, and flexibility in utilization [16,17].

The potential for the entry of nanoparticles through pores and food tract, causes increasing of particles's rate to interact with the enzyme system in the body of the larvae. So that in low concentrations, it has accumulated in the body of the larvae which is able to inhibit the work system of the Acetylcholinesterase Enzym (AChE). It is understand that the reason why Nano Silver concentration (Ag_2NO_3) is lower than carbamate.

This study strengthens the present study. There is a better expediency, especially nano particles Ag_2NO_3 against other organisms. Compared to the use of other insecticides, Ag_2NO_3 Nano is more environmentally friendly. Artificial nanoparticle particles have been known to affect non-target aquatic organisms [18]. There is information about the occurrence of acute toxicity to non-target aquatic species. Silver that are synthesized by *Plumeria rubra*- and *Pergularia daemia* nanoparticles did not show toxicity effects on *P. reticulata* fish after 48 hours of exposure to LC_{50} [19]. Silver nanoparticles were synthesized using extracts of *Solanum nigrum* berry which were non-toxic to two mosquito predators, *Toxorhynchites* larvae and *Diplonychus annulatum* [20]. Nano Silver can be applied to various health care products [10]. The most interesting thing is the potential use of silver nanoparticles (AgNP) in mosquito control. More than several groups have reported AgNP synthesizing which can limit the growth of larvae and/or pupae from *Andopheles* [21-22].

Carbamate (CO_2NH_3 derivatives)

Carbamate is the most widely used compound for insecticides. Pesticides that are categorized as

carbamates: propoxur, furadan and bendiocarb [6, 23]. Insecticides made from active carbamate (methonyl) in *Ae. aegypti*, the concentration needed for LC_{50} is 0.802 mg / l and LC_{95} is 18.832 mg / l. Effective concentration of *Ae. aegypti* requires a smaller amount than carbamate, for LC_{50} is equal to 0.64 mg / l [7].

The results of the study for carbamate toxicity showed that concentration of 0.5 mg / lt, the mortality rate was 92%. At a lower concentration of 0.01 mg / lt, the mortality of test animals was 80%. A concentration greater than 0.5 mg / lt is needed to kill all test animals. These concentration figures resistance of *Aedes aegypti* larvae.

This research also occurs in other countries. Variations in mortality and Aedes's resistance to insecticides in various countries were influenced by the different bio-ecological characteristics of each mosquito. These characters are controlled by the metabolism intensity of Monoxygenases enzym, Esters enzym, Acetylcholinesterase enzym (AChE) and Glutathione-S-transferase (GST) enzymes in the process of detoxifying insecticides [4].

That performance of the enzyme as a determinant of insecticide absorption against the target mosquito. The specific target site is usually an enzyme or protein that acts as an effector of insecticides [10]. Other studies show the high metabolic activity of enzymes Esters in *Aedes aegypti*. In Sri Lanka it causes Potential Resistance of Malathion Low activity of the enzyme Glutathione-S-transferase (GSTs) in *Aedes aegypti* and also it causes mosquito has a high susceptibility status to λ - Cyhalothrin [24-26].

Carbamates act to inhibit the acetylcholinesterase enzyme (AChE) which caused the accumulation of acetylcholine (ACh) Acetylcholine which accumulated in the Central Nervous System (CNS) will induce tremors, coordination, convulsions and others [7]. Acetylcholinesterase (AChE) is a proven target for controlling dengue mosquitoes (*Aedes aegypti*). A single amino acid mutation (G119S) in *Aedes aegypti* forms AChE-I (AgAChE), which provides resistance AChE inhibitors. G119S resistance, AgAChE mutants, and high toxicity to *An. gambiae* has the effect of properties changes. Inhibitors can be raised from 'core' shape of carbamates like aldicarb and pirazol-4-yl methylcarbamate 4a - e although none of these compounds show selectivity which is useful for inhibiting AgAChE to AChE [6].

The high concentration of carbamate to kill *Aedes aegypti* larvae, caused by the entry of that material through only the gastrointestinal tract. High concentration is needed to achieve the occurrence of enzyme system disorders in *Aedes aegypti* larvae. Acetylcholinesterase Enzym (AChE), as a controlling metabolism of *Aedes aegypti* larvae, has not been completed with 0.5 mg / l concentration of carbamate.

The synthesis of nano-silver particles (Ag_2NO_3) with carbamate is the next step to find the effective concentration in the use of carbamate, or nano silver independently. Synthesis is possible for the combination of chemical entry systems into the larva's body. The nano-silver able to penetrate through by the pores between segments in the abdomen, while carbamates can enter the

body of the larvae through the gastrointestinal system. Synthesis also can produce the new compounds with specific properties of Ag_2NO_3 and CO_2NH_3 , so it is possible will be a substitution reaction which resulting a new compound of Ag_2NH_3 , with the cutting of NH_3 chain from CO_2NH_3 .

4 Conclusion

The concentration of 0.5 mg/lit carbamate compounds (CO_2NH_3) can kill 92% of test animals while the concentration of 0.3 mg/l nano-silver (Ag_2NO_3) can kill 100% of them. The result shows that nanosilver (Ag_2NO_3) is more effective to kill animals test (*Aedes aegypti* larvae).

References

1. M. Rahardjo, IJSBAR **22**, 1 (2015)
2. M. Rahardjo, Y.H. Darundiati, Praba G., *Proceeding of International Meeting Of Public Health (IMOPH)*, Kne Publisher, (2018)
3. M. Rahardjo, Y.H. Darundiati, Praba G., IJPHRD **9**, (2018)
4. Dono D, et al, J. Entomol. Indon. **7**, 1, 9-27, (2010)
5. Beroya AM, *Mengenal Lingkungan Hidup* (Yakoma, 2000)
6. Wong MD et al, Plos One **7**, 10, (2012)
7. Pujiastuti E, Aspirator **2**, 2, 77–83, (2010)
8. World Health Organization (WHO). *The WHO Recommended Classification of Pesticides by Hazard and Guidelines to Classification*, (2009)
9. Subagyo A, IOP Conf. Series: Journal of Physics, (2018)
10. Amerasan D, et al, J Pest Sci, **89**, 249–256, (2016)
11. Haldar KM, et al, Parasitol Res, doi: 10.1007/s00436-013-3288-4, (2013).
12. Santhoshkumar T, et al, Parasitol Res, **108**, 693-702, (2013)
13. Sutiningsih D, Martono E, Satoto TBT, AJPJR, **11**, 10, (2018)
14. Pam DD et al, IJHAS, **16**, 2, (2017)
15. IOPScience, *Nanotechnology encompasses the Understanding of Fundamental physic, Chemistry*, **15**, (2019)
16. Denis et al, Ther, **3**, 4, (2015)
17. Sailaja AP, Int J Pharm, **3**, 45-50, (2016)
18. Benelli G, JKUS, 424-435, (2017)
19. Salunkhe RB, Parasitol Journal, **109**, 823-831, (2015)
20. Santhoshkumar T et al, Parasitol Res, **108**, 693-702, (2011)
21. Birla S, Lett Appl Microbiol **48**, 173–179, (2016)
22. Murugan K, Parasitology Journal, **153**, 129–138, (2015)
23. Kenneth D, *Organophosphate Toxicity, eMedicine, Drug and Disease*, (2018)
24. N. Endo, E.A.B. Eltahir, Malaria Journal, **48**, (2018)
25. Nwane P, Etang J, Chouaïbou M, Toto JC, Koffi A, Mimpfoundi R, et al. Parasit Vectors, **6**, 41, (2013)
26. Oduola AO, Idowu ET, Oyebola MK, Adeogun AO, Olojede JB, Otubanjo OA, et al. Parasit vectors, **11**, 5, 116, (2012)

Nanoparticle of silver nitrate (Ag_2NO_3) and Carbamate (CO_2NH_3 derivatives) for vector control of *Aedes aegypti*

ORIGINALITY REPORT

10%

SIMILARITY INDEX

6%

INTERNET SOURCES

7%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Universitas Diponegoro 2%
Student Paper
- 2** textroad.com 2%
Internet Source
- 3** Wong, Dawn M., Jianyong Li, Qiao-Hong Chen, Qian Han, James M. Mutunga, Ania Wysinski, Troy D. Anderson, Haizhen Ding, Tiffany L. Carpenetti, Astha Verma, Rafique Islam, Sally L. Paulson, Polo C.-H. Lam, Maxim Totrov, Jeffrey R. Bloomquist, and Paul R. Carlier. "Select Small Core Structure Carbamates Exhibit High Contact Toxicity to "Carbamate-Resistant" Strain Malaria Mosquitoes, *Anopheles gambiae* (Akron)", PLoS ONE, 2012. 1%
Publication
- 4** www.shs-conferences.org 1%
Internet Source
- 5** www.dovepress.com 1%
Internet Source

6	Submitted to Florida International Baccalaureate Consortium Student Paper	1%
7	Niken Subekti, Rizka Saputri. "The application of Cinnamomum aromaticum nanoparticle and chlorpyrifos for controlling Tribolium castaneum", AIP Publishing, 2019 Publication	1%
8	Submitted to University of Sheffield Student Paper	1%
9	eprints.undip.ac.id Internet Source	1%

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On