Jurnal International Bereputasi
Scopus Q4

Ahmad Ni'matullah Al-Baarri
Sebagai Corresponding Author
(Bukti terlampir)

International Journal of Dairy Science 9 (2): 56-62, 2014
ISSN 1811-9743 / DOI: 10.3923/ijds.2014.56.62
© 2014 Academic Journals Inc.

Resistance of Immobilized Lactoperoxidase Activity from Bovine Whey Against Storage Solutions

¹Dwi Novrina Nawangsari, ²Ahmad Ni'matullah Al-Baarri, ²Sri Mulyani,
²Anang Mohamad Legowo and ²V. Priyo Bintoro

¹Department of Animal Product Technology,

²Department of Food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University,
Semarang, Indonesia

Corresponding Author: Ahmad Ni'matullah Al-Baari, Department of food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

Resistance of Immobilized Lactoperoxidase Activity from Bovine Whey Against Storage Solutions

¹Dwi Novrina Nawangsari, ²Ahmad Ni'matullah Al-Baarri, ²Sri Mulyani, ²Anang Mohamad Legowo and ²V. Priyo Bintoro

¹Department of Animal Product Technology,

²Department of Food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

Corresponding Author: Ahmad Ni'matullah Al-Baari, Department of food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

ABSTRACT

Lactoperoxidase (LPO) could be simply obtained from whey through immobilization using a cation exchange resin of SP Sepharose. LPO received high attention since the antimicrobial properties of LPO system (LPOS) that are consisted of LPO, SCN^- and H_2O_2 was able to generate OSCN^- for strong antimicrobial agents. This study was done to analyze the immobilization efficiency of LPO onto two types of sepharose: SP-Sepharose Fast Flow (SPFF) and SP-Sepharose Big Beads (SPBB). The remaining of LPO's activity (%) against storage solution was also observed. The whey was obtained from bovine skimmed milk that was coagulated using rennet and acid lactic. The LPO was obtained from whey using SPFF. To analyze the resistance of enzyme activity, the immobilized LPO was stored in pure water, phosphate buffer, milk and whey at 10°C. The activity of LPO was monitored for 10 days. The result indicates that the LPO could be purified from whey and single band has been detected using electrophoresis method. The obtained LPO (35 U mL^{-1}) was attached onto SPFF and SPBB. Maximum immobilization efficiency has been achieved by 0.6 g SP-FF and 0.9 g SP-BB. LPO activity of the immobilized LPO were able to be kept until 5 days when it was stored in whey. Other storage solution remained various LPO activity during storage. This study concluded that IE of LPO attached onto sepharoses might be reached variously depended on the sepharose type and whey as LPO source might be used for maintaining LPO activity.

Key words: Lactoperoxidase, SP sepharose fast flow, SP sepharose big beads, immobilization, remaining activity

INTRODUCTION

Lactoperoxidase (LPO), together with SCN^- and H_2O_2 have been understood to generates intermediate product of OSCN^- as antibacterial agent that has a broad spectrum of antimicrobial effects against bacteria, fungi and viruses. This antibacterial agent could be produced if these three components exist in the medium (Seifu *et al.*, 2005). LPOS has been widely used as a preservative in dairy products and nondairy products (Seifu *et al.*, 2004; Touch *et al.*, 2004; FAO/WHO, 2005; Boots and Floris, 2006; Oghaiki *et al.*, 2007; Fweja *et al.*, 2008; Al-Baarri *et al.*, 2011a).

It has been understood that whey contains large number of LPO therefore purification method of LPO from whey has been well developed (Touch *et al.*, 2004; Zhou and Lim, 2009;

Al-Baarri *et al.*, 2010). SP-Sepharose has been known to provide beneficial effect for the immobilization efficiency since SP-Sepharose almost completely immobilized LPO and reusable (Al-Baarri *et al.*, 2010). Although, LPO was widely used in food application but it still remained the problem of its expensiveness, therefore the immobilization of LPO was needed for the reuse of LPO.

SP Sepharose has been known as immobilization agent for capturing lactoperoxidase (Fee and Chand, 2006; Hayashi *et al.*, 2012). SP Sepharose Fast Flow (SPFF) and SP Sepharose Big Beads (SPBB) were the common resin for immobilization since they provide simply application, long term of use and easy for reuse (Amersham-Bioscience, 2001). If compared to other immobilization agent such as chitosan, SP Sepharoses showed higher capturation of lactoperoxidase (Al-Baarri *et al.*, 2012). In the other hand, SP Sepharose application for immobilization of LPO resulting in the much more expensive of the use of LPO, therefore the efficient use of SP Sepharose to immobilize LPO is required. Based on related literatures, there was no documentation for the efficient use of LPO immobilization using SP Sepharose, therefore this research has been done for analyzing the maximum capturation of LPO onto SP Sepharose. This information might provide the benefit for minimum use of SP Sepharose for LPO immobilization. Since the immobilized LPO allowed the reuse of enzyme, the appropriate storage solution for keeping the enzyme activity is needed. To answer this, this research has also been done for analyzing the remaining of LPO activity after storage.

MATERIALS AND METHODS

Materials: SP Sepharose Fast Flow and SP Sepharose Big Beads were purchased from Amersham Pharmacia Biotech, Sweden (Lot. No. 10029743 and 10081054, respectively). Microbial derived rennet was purchased from Singapore and ABTS or 2,20-azino-bis (3-ethylbenz-thiazoline-6-sulphonic acid) was obtained from Kagawa Science (Lot No. 7ROZC-EC) Tokyo Chemical, Industry Co. Ltd., Japan. Cow's milk was obtained from Faculty of Animal and Agricultural Sciences's farm, Diponegoro University, Semarang, Indonesia. The spectrophotometer (Mini UV-1800, Shimadzu, Japan) was used for analysis enzyme activity. Unless other compounds specified, all other compounds were reagent grade.

Preparation of whey: The whey was prepared as method performed by Al-Baarri *et al.* (2011b) without any modification.

LPO immobilization from whey: The procedure for immobilization of LPO was conducted as the method that was performed by Al-Baarri *et al.* (2010) with minor modifications. SP Sepharose Fast Flow (SPFF) was used as agent for LPO immobilization from whey. Whey at the volume of 1800 mL was eluted through a glass column (3×40 cm) filled with 60 g of SPFF. Prior to elution, SPFF was washed with 300 mL of Phosphate Buffer (PB) (pH 6.8) containing 1 M NaCl to remove unnecessary compounds. The whey was circulated through the column using feedback tubing and a peristaltic pump. The circulation was done at the flow rate of 1.0 mL min⁻¹. After draining the whey away, the resin was washed with 300 mL of 0.4 mM NaCl in 0.1 mM phosphate buffer (pH 7.0) using fraction collector (10 mL per tube) to obtain the solution containing high concentration of LPO. Three groups of fractions (fraction number 1-10, 11-20, 21-30) were analyzed for protein profile using sodium dodecyl sulfate polyacrylamide gel electrophoresis

(SDS-PAGE) to check the purity. Finally, based on the SPS PAGE analysis, the fraction number 21-30 was selected for LPO activity analysis (LPO activity was 35 U mL⁻¹). Thus, this fraction was used throughout experiment.

Determination of captured LPO onto SP sepharoses: SPFF and SPBB (0.1-1.0 g) were washed in 1 M NaCl in PB pH 7.0 and then were placed in the column (1×10 cm). The immobilization process was started with the elution of 1 mL of LPO through column. The flow rate was set into 1 mL min⁻¹ using peristaltic pump. The output was collected for measurement of remained LPO activity in the SP Sepharoses. This experiment was repeated three times and column were wash with serial elution of 1 M NaOH and pure water, respectively. The immobilized LPO was stored in pure water, PB, milk and whey. All storage solutions were sterilized using autoclave at 110°C for 10 min. Immobilized LPO was stored at 4°C for 10 days. The remaining of LPO activity immobilized onto SP-Sepharose was calculated by eluting immobilized LPO using 1 M NaCl in PB pH 7.0. The LPO activity in the elution was analyzed spectrometrically.

LPO activity determination: LPO activity was performed as the following method: 450 µL of 1.0 mM ABTS in 10 mM acetate buffer (pH 4.4) and 450 µL of 0.55 mM H₂O₂ in pure water were gently poured into the cuvette. The enzyme (50 µL) was subsequently added to cuvette. The increase of absorbance at 412 nm measured for 20 sec. One unit of LPO enzymatic activity was expressed as the amount of enzyme needed to oxidize 1 µmol ABTS min⁻¹. The molar extinction coefficient of ABTS at 412 nm was 32,400 M⁻¹ cm⁻¹ (Touch *et al.*, 2004).

Immobilization efficiency: The immobilization efficiency (IE) was calculated as follows:

$$IE (\%) = \frac{E_1}{E_0} \times 100$$

where, E₀ is the LPO activity added to the SP Sepharoses (U mL⁻¹) and E₁ is the LPO activity embedded in the SP Sepharose (U mL⁻¹) (Al-Baarri *et al.*, 2010).

RESULTS AND DISCUSSION

Purification LPO: Whey has a lot of enzymes and it is available in low cost because whey is by product of dairy manufacture so it is the challenge to use whey as enzyme sources including LPO. The LPO activity and band(s) of the solution obtained from the elution of 1 M NaCl in PB pH 7.0 through SPFF containing LPO were checked using spectrophotometer and SDS PAGE, respectively. As mention in methods, SPFF containing LPO was generated from whey that was eluted through SPFF column. The result of LPO activity was 27.7±2.9, 39.5±4.5 and 35.2±3.4 U mL⁻¹ for fraction number 1-10, 11-20, 21-30, respectively (data not presented). The highest of LPO activity was group of fraction number 11-20, however since the band of this group showed two bands indicating two protein was detected, for whole of experiment, group of fraction number 21-30 was used. This group showed single band indicating only LPO that was captured by SPFF.

In this research SPFF was used to obtain LPO since this ion exchange resin has diameter 45-165 µm resulting in the wider of surface area than SPBB (Amersham-Bioscience, 2001). In line with this result, Touch *et al.* (2004) used SPFF for purifying LPO from whey resulting in the good

Fig. 1: Immobilization efficiency (IE) of LPO on two types of resin, i.e., SP Sepharose Big Beads (SPBB) and SP Sepharose Fast Flow (SPFF). Two types of resin (0.1-1.0 g) was used to immobilized 1 mL of LPO (35 U mL^{-1}). Flow rate has been set to 1 mL min^{-1} using peristaltic pump. Values are means of three sets of experiments, Error bars represent standard errors of the mean

ability to catch LPO (108 U mL^{-1}). The activity of LPO in this research was less than that of other researcher since the absence of microfiltration step in this research. It has been known that the microfiltration might concentrate the enzyme resulting in the high activity of LPO.

Immobilization efficiency: Immobilization efficiency plays an important role for determination of immobilization agent. This research determined Immobilization Efficiency (IE) of LPO using SPFF dan SPBB (Fig. 1). The volume of SP Sepharoses used in this experiment had a range from 0.1 to 1.0 g to catch the LPO at initial activity of $35.2 \pm 3.4 \text{ U mL}^{-1}$. The increase of IE was found as an increase of SP Sepharoses's weight. As describe in Fig. 1, when the 0.6 g of SPFF was employed, the IE achieved 100% indicating all of LPO employed was able to be captured by SPFF. When the weight of SPFF was increased, the IE was in steady state maximumly. An increase SPBB from 0.1 to 1.0 g elevated the IE from 38.6 to 100%. However, 0.9 g of SPBB completely captured LPO resulting the IE of 100%. One gram of SPFF was reported to have a maximum capture of LPO in 300 mL whey (equal to 750 U mL^{-1} LPO activity) (Al-Baarri *et al.*, 2010). This can be explained that the capture depended on the quantity of enzymes per milliliter. This research used high activity of LPO resulting in the loss of LPO activity.

Since the SPFF and SPBB provided the maximum IE at 0.6 and 0.9 g, respectively, thus these amounts of SP Sepharoses has been used in the rest of experiment. The LPO immobilized onto SPFF and SPBB was stored in the various storage solutions: Pure water, PB, milk and whey for 10 days at 10°C .

Remaining LPO activity during storage: The percentage of remaining immobilized LPO activity stored at 10°C for 10 days in various storage solutions is shown in Fig. 2a and b. LPO activity of immobilized LPO was measured after purging the LPO attached onto SP Sepharose with 1 M NaCl in PB pH 7.0. The percentage of remaining LPO activity was determined by comparing the LPO activity after storage to the initial immobilized LPO at first day of storage.

The percentage of LPO activity attached onto SPBB and SPFF is shown on Fig. 2a and b, respectively. Based on Fig. 2a, whey was able to maintain 100% enzyme activity of LPO attached onto SPBB within 4 days. The extension of storage time resulted in the remarkable

Fig. 2(a-b): (a) Percentage of remaining LPO activity attached onto SP Sepharose Big Beads (SPBB) during 10 days of storage at 10°C in various storage solutions. This experiment used 1 g of SPBB for immobilizing 1 mL of LPO (35 U mL⁻¹). Immediately after immobilization, SPBB then was stored in sterilized pure water, phosphate buffer, milk and whey for 10 days at 10°C. Values are mean of three sets of experiments. Error bars represent standard errors of the mean and (b) Percentage of remaining LPO activity attached onto SP Sepharose Fast Flow (SPFF) during 10 days of storage at 10°C in various storage solution. This experiment used 1 g of SPFF for immobilizing 1 mL of LPO (35 U mL⁻¹). After immobilization, SPFF was stored in sterilized pure water, phosphate buffer, milk and whey. Values are mean of three sets of experiments. Error bars represent standard errors of the mean

reduction of remaining LPO activity. Milk was able to maintain the LPO activity at seven days of storage even though the remaining of LPO activity at that time was very negligible in amount (2.3%).

As previously mentioned, whey was able to completely keep LPO activity within 4 days of storage. This can be explained that whey components support the activity of LPO. It has been studied that LPO activity might be inhibited by casein (Singh *et al.*, 2009) while the casein has been removed from whey.

The remaining of LPO activity attached onto SPFF during 10 days of storage at 10°C is shown on Fig. 2b. PB was able to maintain 100% of LPO activity within 5 days of storage. The milk and whey were able to keep 100% of LPO activity within 3 days of storage. The elevation of storage time until 5 days in whey could keep enzyme activity although the remaining enzyme activity was small amount (10.08%). Based on the availability, also since LPO was simply derived from whey, the storage of immobilized LPO in whey should keep its activity up to 5 days of storage. Therefore it is suggested that immobilized LPO should be stored in whey.

CONCLUSION

The results can be concluded that LPO could be purified from whey using fraction number 21-30. One milliliter of LPO (35 U mL^{-1}) could be completely immobilized onto 0.6 g SPFF or 0.9 g SPBB (immobilization efficiency was 100%). Among various storage solution, whey was able to keep 100% of LPO activity up to 5 days of storage.

ACKNOWLEDGMENTS

This study was supported by funds from the Ministry of Research and Technology of Republic of Indonesia for the project "Insentif Riset SINAS" with Grant Project No. RT-2013-2163.

REFERENCES

- Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2010. Scale-up studies on immobilization of lactoperoxidase using milk whey for producing antimicrobial agent. *J. Indonesian Trop. Anim. Agric.*, 35: 185-191.
- Al-Baarri, A.N., M. Hayashi, M. Ogawa and S. Hayakawa, 2011a. Effects of mono-and disaccharides on the antimicrobial activity of bovine lactoperoxidase system. *J. Food Prot.*, 74: 134-139.
- Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2011b. Application of lactoperoxidase system using bovine whey and the effect of storage condition on lactoperoxidase activity. *Int. J. Dairy Sci.*, 6: 72-78.
- Al-Baarri, A.N., M. Ogawa, T. Visalsok and S. Hayakawa, 2012. Lactoperoxidase immobilized onto various beads for producing natural preservatives solution. *J. Applied Food Technol.*, 1: 4-6.
- Amersham-Bioscience, 2001. Use of sodium hydroxide for cleaning and sanitizing chromatography media and systems. Application Note 18-1124-57 AD, Process Chromatography, Amersham Bioscience, USA. <http://risk.management6.com/Use-of-sodium-hydroxide-for-cleaning-and-sanitizing-chromatography-download-w15138.pdf>.
- Boots, J.W. and R. Floris, 2006. Lactoperoxidase: From catalytic mechanism to practical applications. *Int. Dairy J.*, 16: 1272-1276.
- FAO/WHO, 2005. Benefits and potential risks of the lactoperoxidase system of raw milk preservation. Report of an FAO/WHO Technical Meeting, November 28-December 2, 2005, FAO Headquarters, Rome, Italy, pp: 1-73.
- Fee, C.J. and A. Chand, 2006. Capture of lactoferrin and lactoperoxidase from raw whole milk by cation exchange chromatography. *Separation Purification Technol.*, 48: 143-149.
- Fweja, L.W.T., M.J. Lewis and A.S. Grandison, 2008. Challenge testing the lactoperoxidase system against a range of bacteria using different activation agents. *J. Dairy Sci.*, 91: 2566-2574.
- Hayashi, M., S. Naknukool, S. Hayakawa, M. Ogawa and A.B.A. Ni'matullah, 2012. Enhancement of antimicrobial activity of a lactoperoxidase system by carrot extract and α -carotene. *Food Chem.*, 130: 541-546.
- Oghaiki, N.A., F. Fonteh, P. Kamga, S. Mendi and H. Imele, 2007. Activation of the lactoperoxidase system as a method of preserving raw milk in areas without cooling facilities. *Afr. J. Food Agric. Nutr. Dev.*, 7: 1-14.
- Seifu, E., E.M. Buys and E.F. Donkin, 2004. Quality aspects of Gouda cheese made from goat milk preserved by the lactoperoxidase system. *Int. Dairy J.*, 14: 581-589.

- Seifu, E., E.M. Buys and E.F. Donkin, 2005. Significance of the lactoperoxidase system in the dairy industry and its potential applications: A review. *Trends Food Sci. Technol.*, 16: 137-154.
- Singh, A.K., N. Singh, S. Sharma, K. Shin and M. Takase *et al.*, 2009. Inhibition of lactoperoxidase by its own catalytic product: Crystal structure of the hypothiocyanate-inhibited bovine lactoperoxidase at 2.3-Å resolution. *Biophys. J.*, 96: 646-654.
- Touch, V., S. Hayakawa, S. Yamada and S. Kaneko, 2004. Effects of a lactoperoxidase-thiocyanate-hydrogen peroxide system on *Salmonella enteritidis* in animal or vegetable foods. *Int. J. Food Microbiol.*, 93: 175-183.
- Zhou, Y. and L.T. Lim, 2009. Activation of lactoperoxidase system in milk by glucose oxidase immobilized in electrospun polylactide microfibers. *J. Food Sci.*, 74: C170-C176.

Alert

HOME

JOURNALS

AUTHORS

SUBSCRIBERS

CONTACT

Search

International Journal of Dairy Science

Publisher: Academic Journals Inc., USA

eISSN: 1811-9751
pISSN: 1811-9743

International Journal of Dairy Science is a high quality peer-reviewed scientific journal dedicated to publish cutting edge research work on all aspects of dairy sciences. Scope of the journal includes: Biochemistry, breeding, economics, engineering, environment, food science, genetics, microbiology, nutrition, pathology, physiology, processing, public health quality assurance, sanitation, microbiology and bacteriology.

Submit your next paper to International Journal of Dairy Science via [online submission system](#).

Editor-in-Chief: [Hussein Azzaz Murad](#)

Editor-in-Chief

Hussein Azzaz Murad
National Research Center, Egypt

REGIONAL EDITORS

Abd El-Kader Mahmoud Kholif
National Research Centre, Egypt

ASSOCIATE EDITORS

Ahmed Eid Kholif
National Research Centre, Egypt

Othman El-Mahdy Sayed Othman
National Research Center, Egypt

Mahmoud Rushdi Abd Ellah
Assiut University, Egypt

Navigation

- [Online First](#)
- [Current Issue](#)
- [Previous Issues](#)
- [Editorial Board](#)
- [Submit a Manuscript](#)
- [Guide to Authors](#)
- [Article Processing Charges](#)
- [Subscribe to E-alerts](#)

International Journal of Dairy Science

Google Scholar

Indexed In

- [ASCI-Database](#)
- [Asian Digital Library](#)
- [Cambridge Scientific Abstract](#)
- [Chemical Abstract Services](#)
- [FSTA](#)
- [Google Scholar](#)
- [SCIMAGO](#)

Dr. Ashraf Mohamed Abdel Rahman Abu-Seida
Cairo University, Egypt

Dr. Wahid Mohamed Ahmed
National Research Center, Egypt

Zuhair Bani Ismail
Purdue University, USA

TECHNICAL EDITORS

Mostafa Sayed Abd El-Latif Khattab
National Research Center, Egypt

Jinming Huang
Shandong Academy of Agricultural Sciences, China

Mervat Ibrahim Foda
National Research Center, Egypt

Veerasamy Sejian
National Institute of Animal Nutrition and Physiology, India

A. Yasotha
Madras Veterinary College, India

Dr. Alaa Abdel-Moneam Ghazy Mohamed
National Research Center, Egypt

Sumeet Sharma
University of Alberta, Canada

Ebubekir Ceylan
Hakkari University, Turkey

Suleyman Cilek
Kirikkale University, Turkey

Karakus Kadir
Yuzuncu Yil University, Turkey

Karima Galal Abdel Hameed
South Valley University, Egypt

Richard Osei Amponsah
University of Ghana, Ghana

[Home](#) · [Journals](#) · [For Authors](#) ·
[For Subscribers](#) · [ASCI](#)

© Science Alert. All Rights Reserved

Search SciAlert website

SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

International Journal of Dairy Science

Country[Pakistan](#) - [SJR Ranking of Pakistan](#)**Subject Area and Category**[Agricultural and Biological Sciences](#)
[Animal Science and Zoology](#)[Veterinary](#)
[Food Animals](#)**Publisher**[Asian Network for Scientific Information](#)**Publication type**[Journals](#)**ISSN**

18119743, 18119751

Coverage

2006-ongoing

Scope

International Journal of Dairy Science is dedicated to disseminate the international original research on all aspect of dairy science. International Journal of Dairy Science publishes original scientific research on all aspects of dairy science including: animal husbandry, the physiology, biochemistry and endocrinology of lactation, milk production, composition, preservation, processing and separation, biotechnology and food science, properties of milk proteins and other components, dairy products such as cheese, fermented milks and spreads, relevant studies in bacteriology, enzymology and immunology, the use of milk products in other foods; and the development of methods relevant to these subjects.

[Homepage](#) [Join the conversation about this journal](#)

14

H Index

Quartiles

Show this widget in your own website

Just copy the code below and paste within your html code:

International Journal of Dairy Science

Q4

Animal Science and Zoology

best quartile

SJR 2018

0.18

powered by scimagojr.com

←

<https://www.scimagojr.com>

Sources

Title

Enter title

Find sources

Title: International Journal Of Dairy Science x

i

CiteScore metrics for journals and serials

CiteScore metrics from Scopus are:

• Comprehensive

• Transparent

• Current and free

Use this page to find a source and view associated metrics. Use qualitative as well as quantitative metrics when presenting your research impact. Always use more than one quantitative metric. Learn more about CiteScore.

Citations in 2018

Documents from 3 years

2014

2015

2016

2017

2018

2019

Filter refine list

Apply Clear filters

1 result

Export to Excel

Download Scopus Source List

Learn more about Scopus Source List

View metrics for year: 2018

Display options

☐ Display only Open Access journals

Counts for previous 3 years

- ☐ No minimum selected
- ☐ Minimum citations
- ☐ Minimum documents

Citescore highest quartile

- ☐ Show only titles in top 10 percent
- ☐ 1st quartile
- ☐ 2nd quartile
- ☐ 3rd quartile
- ☐ 4th quartile

Source type

- ☐ Journals
- ☐ Book Series
- ☐ Conference Proceedings
- ☐ Trade Publications

Apply Clear filters

Source title ↓	CiteScore ↓	Highest percentile ↓	Citations 2018 ↓	Documents 2015-17 ↓	% Cited ↓	SNIP ↓	
International Journal of Dairy Science	0.56	32% 20/29 Food Animals	50	90	42	0.741

Top of page

1 document result

Search within results...

Analyze search results

Show all abstracts Sort on: Date (newest)

Refine results

Limit to Exclude

☐ All ☐ RIS export Download View citation overview View cited by Save to list

Access type

☐ Open Access (1)

Year

☐ 2014 (1)

Author name

- ☐ Al-Baarri, A.N.M. (1)
- ☐ Legowo, A.M. (1)
- ☐ Mulyani, S. (1)
- ☐ Nawangsari, D.N. (1)
- ☐ Priyo Bintoro, V.P. (1)

Subject area

- ☐ Agricultural and Biological Sciences (1)
- ☐ Veterinary (1)

Publication stage

Document type

Source title

Keyword

Affiliation

Funding sponsor

Country/territory

☐ Indonesia (1)

Source type

Language

Limit to Exclude

	Document title	Authors	Year	Source	Cited by
☐ 1	Resistance of immobilized lactoperoxidase activity from bovine whey against storage solutions Open Access	Nawangsari, D.N., Al-Baarri, A.N., Mulyani, S., Legowo, A.M., Priyo Bintoro, V.P.	2014	International Journal of Dairy Science 9(2), pp. 56-62	1
View abstract View at Publisher Related documents					

Display: 20 1

results per page

Top of page

BUKTI KORESPONDENSI SAYA

COVER LETTER FOR SUBMISSION OF NEW MANUSCRIPTS

To International Journal of Dairy Science's Editor

Subject: **SUBMISSION OF NEW MANUSCRIPT FOR EVALUATION**

I am enclosing herewith a manuscript entitled "Resistance of Immobilized Lactoperoxidase Activity From Bovine Whey Against Storage Solutions" submitted to "**International Journal of Dairy Science**" for possible evaluation.

With the submission of this manuscript I would like to undertake that the above mentioned manuscript has not been published elsewhere, accepted for publication elsewhere or under editorial review for publication elsewhere; and that my Institute's [**Diponegoro University, Semarang, Indonesia**] representative is fully aware of this submission.

Select Type of Submitted manuscript:

- Original Article ✓
- ~~Review Article~~
- ~~Mini Review~~
- ~~Short Communication~~
- ~~Clinical Article~~
- ~~Perspective~~
- ~~Editorial~~
- Any other (specify the type of manuscript)

For the Editor-in-Chief,

I would like to disclose the following information about the project:

RESISTANCE OF IMMOBILIZED LACTOPEROXIDASE ACTIVITY FROM BOVINE WHEY AGAINST STORAGE SOLUTIONS

The research project was conducted under the supervision of:

1. Ahmad Ni'matullah Al-Baarri, PhD (team leader, corresponding author)
2. Sri Mulyani, MP.
3. Prof. Anang Mohamad Legowo
4. Prof. V. Priyo Bintoro

and the project was run as my M.Sc project

This research project was conducted from July 2013 to February 2013

My Research Project was fully sponsored by Ministry of Research and Technology of Republic of Indonesia with grant number RT-2013-2163.

Detail of the each author with his/her contribution in this paper is as under:

Name of the author and e-mail ID	Types of contribution
Ahmad Ni'matullah Al-Baarri, PhD	Co-author and technical supervisor
Sri Mulyani	Co-author and technical supervisor
Prof. Anang Mohamad Legowo	Co-author and data analysis
Prof. V. Priyo Bintoro	Co-author and data interpretation

I would also like to share the following information with Editor-in-Chief

I have the following similar manuscripts already published from this project:

No, I have no article yet in this journal.
This is the first original article, but my supervisor has an accepted article on International Journal of Dairy Science 6 (1): 72-78,2011 entitled:
Application of Lactoperoxidase System Using Bovine Whey and the Effect of Storage Condition on Lactoperoxidase Activity
My supervisor was also published other article in other international journal.

For quick understanding about the importance of the project following are the significant findings of my submitted article?

Both of SP-Sepharose (Fast Flow and Big Beads) have been known as ion exchange resins for immobilizing lactoperoxidase. Since the price of both compounds are expensive, the efficient

use is required. This article showed the optimum amount of lactoperoxidase might be attached to resin and the analysis of enzyme activity in various storage solutions.

How findings of this research work are unique in their nature?

Many investigator used ion exchange resin to immobilize lactoperoxidase, however none investigator mentioned the optimum condition for immobilizing lactoperoxidase.

This research answered those questions and provide the information for the optimum use of sepharose resin to immobilize lactoperoxidase.

Since in the large scale production and utilization of lactoperoxidase is required, the storage solution are the key to keep the enzyme activity. This article showed the best storage solution to keep the enzyme activity.

A paragraph explaining why your manuscript is appropriate for the selected journal

This original article is about lactoperoxidase that naturally obtained from fresh milk. The enzyme has great benefit to the long time preservation of milk and dairy products. Since this journal concerns about dairy science, I believe this journal is appropriate for this article.

Last update on August 10, 2011

To: International Journal of Dairy Science

Formatted: Left

We have received the response from reviewer 1 and reviewer 2 to our article, however we found the question from reviewer 2 only:

Al Baarri, 2011 missed , Also the result of LPO activity why not presented in a table , Also The captions of the figures must be written at the top of the figures and must be clear and explain the figures .

Formatted: Left, Indent: Left: 1 cm, Right: 1.92 cm

Formatted: Font: Not Bold

Formatted: Left

Our responses:

1. Al-Baarri, 2011 has been changed to Al-Baarri, 2011a. We are sorry, it should be added with "a" after "2011"
2. LPO activity did not appear in a table since we would like to show the differences between treatments (storage). We believe that the description through percentage in a figure is much easier to be understood
3. The caption has been moved into the top of figure and the explanation of figure has been completed.

Formatted: Left, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.63 cm + Indent at: 1.27 cm

Formatted: Left

Sincerely,

Ahmad Ni'matullah Al-Baarri

Formatted: Font: Not Bold

COVER LETTER FOR SUBMISSION OF NEW MANUSCRIPTS

To International Journal of Dairy Science's Editor

Subject: **SUBMISSION OF NEW MANUSCRIPT FOR EVALUATION**

I am enclosing herewith a manuscript entitled "Resistance of Immobilized Lactoperoxidase Activity From Bovine Whey Against Storage Solutions" submitted to "International Journal of Dairy Science" for possible evaluation.

With the submission of this manuscript I would like to undertake that the above mentioned manuscript has not been published elsewhere, accepted for publication elsewhere or under editorial review for publication elsewhere; and that my Institute's [Diponegoro University, Semarang, Indonesia] representative is fully aware of this submission.

Select Type of Submitted manuscript:

- Original Article ✓
- Review Article
- Mini Review
- Short Communication
- Clinical Article
- Perspective
- Editorial
- Any other (specify the type of manuscript)

For the Editor-in-Chief,

I would like to disclose the following information about the project:

RESISTANCE OF IMMOBILIZED LACTOPEROXIDASE ACTIVITY FROM BOVINE WHEY AGAINST STORAGE SOLUTIONS

The research project was conducted under the supervision of:

1. Ahmad Ni'matullah Al-Baarri, PhD
2. Sri Mulyani, MP.
3. Prof. Anang Mohamad Legowo
4. Prof. V. Priyo Bintoro

and the project was run as my M.Sc project

This research project was conducted from July 2013 to February 2013

My Research Project was fully sponsored by Ministry of Research and Technology of Republic of Indonesia with grant number RT-2013-2163.

Detail of the each author with his/her contribution in this paper is as under:

<u>Name of the author and e-mail ID</u>	<u>Types of contribution</u>
<u>Ahmad Ni'matullah Al-Baarri, PhD</u>	<u>Co-author, and article content</u>
<u>Sri Mulyani</u>	<u>Co-author and technical supervisor</u>
<u>Prof. Anang Mohamad Legowo</u>	<u>Co-author, data analysis, article interpretation</u>
<u>Prof. V. Priyo Bintoro</u>	<u>Co-author, data interpretation, English guidance</u>

I would also like to share the following information with Editor-in-Chief

I have the following similar manuscripts already published from this project:

No, I have no article yet in this journal.

This is the first original article, but my supervisor has an accepted article on International Journal of Dairy Science 6 (1): 72-78,2011 entitled:

Application of Lactoperoxidase System Using Bovine Whey and the Effect of Storage Condition on Lactoperoxidase Activity

My supervisor was also published other article in other international journal.

For quick understanding about the importance of the project following are the significant findings of my submitted article?

Both of SP-Sepharose (Fast Flow and Big Beads) have been known as ion exchange resins for immobilizing lactoperoxidase. Since the price of both compounds are expensive, the efficient use is required. This article showed the optimum amount of lactoperoxidase might be attached to resin and the analysis of enzyme activity in various storage solutions.

How findings of this research work are unique in their nature?

Many investigator used ion exchange resin to immobilize lactoperoxidase, however none investigator mentioned the optimum condition for immobilizing lactoperoxidase.

This research answered those questions and provide the information for the optimum use of sepharose resin to immobilize lactoperoxidase.

Since in the large scale production and utilization of lactoperoxidase is required, the storage solution are the key to keep the enzyme activity. This article showed the best storage solution to keep the enzyme activity.

A paragraph explaining why your manuscript is appropriate for the selected journal

This original article is about lactoperoxidase that naturally obtained from fresh milk. The enzyme has great benefit to the long time preservation of milk and dairy products. Since this journal concerns about dairy science, I believe this journal is appropriate for this article.

Last update on August 10, 2011

Evaluation Sheet for Article No.[57374-IJDS-AJ]			
1	Internal Reviewers Comments	Incorporated	
		<div>Yes</div>	<div>No</div>
	<ul style="list-style-type: none">The cover letter for this manuscript is not provided by the author. Download sample cover letter from the link and submit duly filled Cover Letter with all your new submissions http://scialert.net/coverletter.doc	<div>The cover letter has been composed</div>	
Abstract: <ul style="list-style-type: none">End the paragraph of abstract section with a clear cut conclusion of study, but length of abstract section should not exceed from more than 250 words.Check the spelling of yellow highlighted words.	<div>The abstract has been modify and the author has been provided comments</div>		
Methodology: <ul style="list-style-type: none">Don't use personal pronouns throughout the text.Provide some detail of used statistical analysis under separate subheading, in methodology section.Don't start a sentence with a numeric value.	<div>Personal pronouns has been removed</div> <div>Since the statistical analysis has not been used, the subheading for statistical analysis was not provided</div> <div>Numeric starting sentences has been corrected</div>		

<p>Results and Discussion:</p> <ul style="list-style-type: none"> Some numeric values mentioned in text are not in accordance with their respective tabular data. For your convenience some of those values have been yellow highlighted. Correct all data irregularities by comparing numeric values given in text with their respective tabular data. Author has cited Figure 3 in text but didn't provide it in the submitted manuscript. Provide this missing figure and make sure that all figures have been cited in text in consecutive numerical order. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation. Provide correct reference of all figures in text. Delete table 1 and explain its data in text, as it does not contain significant amount of data that needs to be presented in tabular form. Define all abbreviations used in figures in their respective captions. The given captions of your figures 1 and 2 have not been written properly. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such captions for all these figures in which all of their main parts have been properly explained. Assign an alphabet to each sub-part of Figure 2 and explain each figure part in figure caption by referring to its respective alphabet. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such caption in which all main parts of this figure have been properly explained. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation. In submitted manuscript Figure 2 has been presented but Author didn't cite this figure in text. Cite this figure in text at appropriate place with suitable explanation and make sure that all figures have been cited in text in consecutive numerical order. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation. 	<p>Numeric value as starting sentences has been corrected</p> <p>The number of figure has been revised and the double check has been done</p> <p>The suggestion for deleting the reference has been made.</p> <p>Table 1 has been deleted and the content has been cited in the text</p> <p>The correct reference of figure has been made.</p> <p>Authors checked all figure number.</p> <p>All abbreviation in the figure has been</p>	
---	---	--

	defined completely All figure has been explained individually and cited properly. Authors has been revised the number of figure. Figure 2 has been separated as figure 2a and 2b. These figures has been cited in the article properly.	
--	--	--

Note:		
--------------	--	--

<p>Note: Please provide line number in case you incorporated the comments. In case you did not incorporate any comments please provide suitable reasons.</p> <p>***Please NOTE: It is requested to please modify this uploaded file according to the suggested modifications and send modified files as Revised Article for further processing. Please don't remove hyper linking from this file and all other working done by Editorial Office.</p> <p>***Please provide DOI or URL or Pub Med. No for journal, conference, proceeding and workshop reference. ISBN No of book and book chapter references. Thesis type of references may be deleted from list and text.</p> <p>***It is difficult for us to identify the modifications, which you have done according to the internal comments. Therefore it is requested to please change the text color (RED) where you have incorporated or</p>		
--	--	--

modified the manuscript.

Ref#17

RESISTANCE OF IMMOBILIZED LACTOPEROXIDASE ACTIVITY FROM BOVINE WHEY AGAINST STORAGE SOLUTIONS

Dwi Novrina Nawangsari¹⁾, Ahmad Ni'matullah Al-Baarri²⁾, Sri Mulyani²⁾, Anang Mohamad Legowo²⁾, V. Privo Bintoro²⁾

1)Department of Animal Product Technology; Faculty of Animal and Agricultural Sciences; Diponegoro University, Semarang, Indonesia

2)Department of Food Technology, Faculty of Animal and Agricultural Sciences; Diponegoro University, Semarang, Indonesia

ABSTRACT

Lactoperoxidase (LPO) could be simply obtained from whey through immobilization using a cation exchange resin of SP Sepharose. LPO received high attention since the antimicrobial properties of LPO system (LPOS) that are consisted of LPO, SCN⁻, and H₂O₂ was able to generate OSCN⁻ for strong antimicrobial agents. This study was done to analyze the immobilization efficiency of LPO onto two types of sepharose: SP-Sepharose Fast Flow (SPFF) and SP-Sepharose Big Beads (SPBB). The remaining of LPO's activity (%) against storage solution was also observed. The whey was obtained from bovine skimmed milk that was coagulated using rennet and acid lactic. The LPO was obtained from whey using SPFF. To analyze the resistance of enzyme activity, the immobilized LPO was stored in pure water, phosphate buffer, milk, and whey at 10°C. The activity of LPO was monitored for 10 days. The result indicates that the LPO could be purified from whey and single band has been detected using electrophoresis method. The obtained LPO (35 U/ml) was attached onto SPFF and SPBB. Maximum immobilization efficiency has been achieved by 0.6 g SP-FF and 0.9 g SP-BB. LPO activity of the immobilized LPO were able to be kept until 5 days when it was stored in whey. Other storage solution remained various LPO activity during storage. This research concluded that IE of LPO attached onto sepharoses might be reached variously depended on the sepharose type and whey as LPO source might be used for maintaining LPO activity.

Key words: Lactoperoxidase, SP Sepharose Fast Flow, SP Sepharose Big Beads, immobilization, remaining activity.

Deleted: 1

Deleted:

Deleted: and

Commented [ANA1]: Please allow us to add the co-authors since this author has a big contribution to our experiments and article writing. The contribution of authors have been mentioned in cover letter.

Commented [O2]: The cover letter for this manuscript is not provided by the author. Download sample cover letter from the link and submit duly filled Cover Letter with all your new submissions <http://scialert.net/coverletter.doc>

Commented [ANA3]: Cover letter has been made and author has sent this cover letter along with this article.

Commented [O4]: End the paragraph of abstract section with a clear cut conclusion of study, but length of abstract section should not exceed from more than 250 words.

Commented [ANA5]: Cover letter has been added and sent via email

Commented [O6]: Check the spelling of yellow highlighted words.

Deleted: remaining

Deleted: immobilized LPO

Deleted: It was concluded that

Deleted: were able to achieved 100% of immobilization efficiency (IE). LPO

Deleted: onto Sepharoses

Commented [ANA7]: Conclusion has been added and the total word counts is 245 (less than 250)

INTRODUCTION

Lactoperoxidase (LPO), together with SCN^- and H_2O_2 have been understood to generates intermediate product of OSCN^- as antibacterial agent that has a broad spectrum of antimicrobial effects against bacteria, fungi and viruses. This antibacterial agent could be produced if these three components exist in the medium (Seifu et al., 2005, Al-Baarri, 2011a). LPOS has been widely used as a preservative in dairy products and nondairy products (Seifu et al., 2004, Touch et al., 2004, FAO/WHO, 2005, Boots and Floris, 2006, Oghaiki et al, 2007, Fweja et al., 2008, Al-Baarri et al., 2011a).

It has been understood that whey contains large number of LPO therefore purification method of LPO from whey has been well developed (Touch et al., 2004, Zhou and Lim, 2009, Al-Baarri et al., 2010). SP-Sepharose has been known to provide beneficial effect for the immobilization efficiency since SP-Sepharose almost completely immobilized LPO and reusable (Al-Baarri et al., 2010). Although LPOS was widely used in food application but it still remained the problem of its expensiveness, therefore the immobilization of LPO was needed for the reuse of LPO.

SP Sepharose has been known as immobilization agent for capturing lactoperoxidase (Fee and Chand, 2006; Hayashi et al., 2012). SP Sepharose Fast Flow (SPFF) and SP Sepharose Big Beads (SPBB) were the common resin for immobilization since they provide simply application, long term of use, and easy for reuse (Amersham-Bioscience, 2001). If compare to other immobilization agent such as chitosan, SP Sepharoses showed higher capturization of lactoperoxidase (Al-Baarri et al., 2012). In the other hand, SP Sepharose application for immobilization of LPO resulting in the much more expensive of the use of LPO, therefore the efficient use of SP Sepharose to immobilize LPO is required. Based on related literatures, there was no documentation for the efficient use of LPO immobilization using SP Sepharose, therefore this research has been done for analyzing the maximum capturization of LPO onto SP Sepharose. This information might provide the benefit for minimum use of SP Sepharose for LPO immobilization. Since the immobilized LPO allowed the reuse of enzyme, the appropriate storage solution for keeping the enzyme activity is needed. To answer this, this research has also been done for analyzing the remaining of LPO activity after storage.

MATERIALS AND METHODS

Materials

Deleted: s

Deleted: appropriate

Commented [O8]: Don't use personal pronouns throughout the text.

Deleted:

Commented [ANA9]: I do not have idea to replace this word, so, I did not change this word.

Commented [O10]: Provide some detail of used statistical analysis under separate subheading, in methodology section.

Deleted: descriptive analysis was used for describing activity of LPO in all treatment.

SP Sepharose Fast Flow, and SP Sepharose Big Beads, were purchased from Amersham Pharmacia Biotech, Sweden (Lot. No.10029743 and No. 10081054, respectively). Microbial derived rennet was purchased from Singapore and ABTS or 2,20-azino-bis (3-ethylbenz-thiazoline-6-sulphonic acid), was obtained from Kagawa Science (Lot No. 7ROZC-EC) Tokyo Chemical, Industry Co. Ltd. Japan. Cow's milk was obtained from Faculty of Animal and Agricultural Sciences's farm, Diponegoro University, Semarang, Indonesia. The spectrophotometer (Mini UV-1800, Shimadzu, Japan) was used for analysis enzyme activity. Unless other compounds specified, all other compounds were reagent grade.

Deleted: SP-FF

Deleted: SP-BB

Commented [ANA11]: ABTS has been replaced as start of sentence

Commented [O12]: Don't start a sentence with a numeric value.

Deleted: or ABTS

Preparation of whey

The whey was prepared as method performed by Al-Baarri et. al. (2011b) without any modification.

LPO Immobilization from Whey

The procedure for immobilization of LPO was conducted as the method that was performed by Al-Baarri et al., 2010 with minor modifications. SP Sepharose Fast Flow (SPFF) was used as agent for LPO immobilization from whey. Whey at the volume of 1800 ml was eluted through a glass column (3 x 40 cm) filled with 60 g of SPFF. Prior to elution, SPFF was washed with 300 ml of phosphate buffer (PB) (pH 6,8) containing 1 M NaCl to remove unnecessary compounds. The whey was circulated through the column using feedback tubing and a peristaltic pump. The circulation was done at the flow rate of 1.0 ml/min. After draining the whey away, the resin was washed with 300 ml of 0.4 mM NaCl in 0.1 mM phosphate buffer (pH 7.0) using fraction collector (10 ml per tube) to obtain the solution containing high concentration of LPO. Three groups of fractions (fraction number 1–10, 11–20, 21–30) were analyzed for protein profile using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS PAGE) to check the purity. Finally, based on the SDS PAGE analysis, the fraction number 21–30 was selected for LPO activity analysis (LPO activity was 35 U/ml). Thus, this fraction, was used throughout experiment.

Deleted: immobilization

Deleted: a

Deleted: -

Deleted: group

Deleted: SDS PAGE

Deleted: analyzed

Deleted: and

Determination of Captured LPO onto SP Sepharoses

SPFF and SPBB (0.1–1.0 g) were washed in 1 M NaCl in PB pH 7.0 and then were placed in the column (1x10 cm). The immobilization process was started with the elution of 1 ml of LPO through column. The flow rate was set into 1 ml/min, using peristaltic pump. The output was collected for measurement of remained LPO activity in the SP Sepharoses. This

Deleted: -

Deleted: -

Deleted: ute

experiment was repeated three times and column were wash with serial elution of 1 M NaOH and pure water, respectively. The immobilized LPO was stored in milk, whey, pure water and whey. All storage solutions were sterilized using autoclave at 110°C for 10 minutes. Immobilized LPO was stored at 4°C for 10 days. The remaining of LPO activity immobilized onto SP-Sepharose was calculated by eluting immobilized LPO using 1 M NaCl in PB pH 7.0. [The LPO activity in the elution was analyzed spectrometrically.](#)

LPO Activity Determination

LPO activity was performed as the following method: 450µl of 1.0 mM ABTS in 10 mM acetate buffer (pH 4.4) and 450 µl of 0.55 mM H₂O₂ in pure water were gently poured into the cuvette. The enzyme (50 µl) was subsequently added to cuvette. The increase of absorbance at 412 nm measured for 20 second. One unit of LPO enzymatic activity was expressed as the amount of enzyme needed to oxidize 1 µmol ABTS/min. The molar extinction coefficient of ABTS at 412nm was 32.400 M⁻¹ cm⁻¹ ([Touch et al., 2004](#)).

Immobilization Efficiency

The immobilization efficiency (IE) was calculated as follows: IE(%)= E1/E0 x 100, where E0 is the LPO activity added to the SP Sepharoses (U/ml) and E1 is the LPO activity embedded in the SP Sepharose (U/ml) ([Al-Baarri et al., 2010](#)).

RESULTS AND DISSCUSSION

Purification LPO

Whey has a lot of enzymes and it is available in low cost because whey is by product of dairy manufacture so it is the challenge to use whey as enzyme sources including LPO. [The LPO activity and band \(s\) of the solution obtained from the elution of 1 M NaCl in PB pH 7.0 through SPFF containing LPO were checked using spectrophotometer and SDS PAGE, respectively.](#) As mention in methods, SPFF containing LPO was generated from whey that was eluted through SPFF column. The result of LPO activity was 27.7±2.9, 39.5±4.5, and 35.2±3.4 U/ml for fraction number 1–10, 11–20, 21–30, respectively [\(data not presented\)](#). [The highest of LPO activity was group of fraction number 11–20, however since the band of this group showed two bands indicating two protein was detected, for whole of experiment, group of fraction number 21–30 was used. This group showed single band indicating only LPO that was captured by SPFF.](#)

Deleted: Table 1 shows t

Deleted: y

Commented [O13]: Some numeric values mentioned in text are not in accordance with their respective tabular data. For your convenience some of those values have been yellow highlighted. Correct all data irregularities by comparing numeric values given in text with their respective tabular data.

Deleted: As shown in Table 1, t

In this research SPFF was used to obtain LPO since this ion exchange resin has diameter 45–165 μm resulting in the wider of surface area than SPBB (Amersham-Bioscience, 2001). In line with this result, Touch et al., 2004 used SPFF for purifying LPO from whey resulting in the good ability to catch LPO (108 U/ml). The activity of LPO in this research was less than that of other researcher since the absence of microfiltration step in this research. It has been known that the microfiltration might concentrate the enzyme resulting in the high activity of LPO.

Immobilization Efficiency

Immobilization efficiency plays an important role for determination of immobilization agent. This research determined immobilization efficiency (IE) of LPO using SPFF dan SPBB (Figure 1). The volume of SP Sepharoses used in this experiment had a range from 0.1 to 1.0 g to catch the LPO at initial activity of 35.2 ± 3.4 U/ml. The increase of IE was found as an increase of SP Sepharoses's weight. As describe in Figure 1, when the 0.6 g of SPFF was employed, the IE achieved 100% indicating all of LPO employed was able to be captured by SPFF. When the weight of SPFF was increased, the IE was in steady state maximumly. An increase SPBB from 0.1 to 1.0 g elevated the IE from 38.6 to 100%. However, 0.9 g of SPBB completely captured LPO resulting the IE of 100%. One gram of SPFF was reported to have a maximum capture of LPO in 300 ml whey (equal to 750 U/ml LPO activity) (Al-Baarri et al., 2010). This can be explained that the capture depended on the quantity of enzymes per mililiter. This research used high activity of LPO resulting in the loss of LPO activity.

Since the SPFF and SPBB provided the maximum IE at 0.6 and 0.9 g, respectively, thus these amounts of SP Sepharoses has been used in the rest of experiment. The LPO immobilized onto SPFF and SPBB was stored in the various storage solutions: pure water, PB, milk, and whey for 10 days in 10°C .

Remaining LPO Activity During Storage

The percentage of remaining immobilized LPO activity stored at 10°C for 10 days in various storage solutions is shown in Figure 2a and 2b. LPO activity of immobilized LPO was measured after purging the LPO attached onto SP Sepharose with 1 M NaCl in PB pH 7.0. The percentage of remaining LPO activity was determined by comparing the LPO activity after storage to the initial immobilized LPO at first day of storage.

The percentage of LPO activity attached onto SPBB and SPFF is shown on Figure 2a and Figure 2b, respectively. Based on Figure 2a, whey was able to maintain 100% enzyme

Deleted: Figure 1 shows w

Deleted: might be

Deleted: these amount

Deleted: p

Commented [O14]: Author has cited Figure 3 in text but didn't provide it in the submitted manuscript. Provide this missing figure and make sure that all figures have been cited in text in consecutive numerical order. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Commented [O15]: Provide correct reference of all figures in text.

Commented [ANA16]: I have double checked about this.

Deleted: 2

Deleted: LPO

activity of LPO attached onto SPBB within 4 days. The extension of storage time resulted in the remarkable reduction of remaining LPO activity. Milk was able to maintain the LPO activity at seven days of storage even though the remaining of LPO activity at that time was very negligible in amount (2.3%).

As previously mentioned, whey was able to completely keep LPO activity within 4 days of storage. This can be explained that whey components support the activity of LPO. It has been studied that LPO activity might be inhibited by casein (Singh et al., 2009) while the casein has been removed from whey.

The remaining of LPO activity attached onto SPFF during 10 days of storage at 10°C is shown on Figure 2b. PB was able to maintain 100% of LPO activity within 5 days of storage. The milk and whey were able to keep 100% of LPO activity within 3 days of storage. The elevation of storage time until 5 days in whey could keep enzyme activity although the remaining enzyme activity was small amount (10.08%). Based on the availability, also since LPO was simply derived from whey, the storage of immobilized LPO in whey should keep its activity up to 5 days of storage. Therefore it is suggested that immobilized LPO should be stored in whey.

CONCLUSION

The results can be concluded that LPO could be purified from whey using fraction number 21–30. One milliliter of LPO (35 U/ml) could be completely immobilized onto 0.6 g SPFF or 0.9 g SPBB (immobilization efficiency was 100%). Among various storage solution, whey was able to keep 100% of LPO activity up to 5 days of storage.

ACKNOWLEDGMENT

This work was supported by funds from the Ministry of Research and Technology of Republic of Indonesia for the project "Insentif Riset SINAS" with grant project number RT-2013-2163.

REFERENCES

Al-Baarri, A.N., M. Hayashi, M. Ogawa and S. Hayakawa, 2011a. Effects of mono- and disaccharides on the antimicrobial activity of bovine lactoperoxidase system. J. Food Prot., 74: 134-139.

1174452ja

Commented [ANA17]: I have double check for the explanation of figure and proper figure

Deleted: ,

Formatted: Indent: First line: 0 cm

Formatted: Level 1, Indent: First line: 0 cm

Formatted: Indent: First line: 0 cm

Commented [ANA18]: Please allow us to explain the our acknowledgment

Formatted: Level 1, Indent: First line: 0 cm

Deleted: Al-Baarri, A.N., 2011. Lactoperoxidase Activity on Bovine Whey at Critical Temperature Storage. Unpublished data.

Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2010. Scale-up studies on immobilization of lactoperoxidase using milk whey for producing antimicrobial agent. J. Indonesian Trop. Anim. Agric., 35: 185-191.

1174454ja

Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2011b. Application of lactoperoxidase system using bovine whey and the effect of storage condition on lactoperoxidase activity. Int. J. Dairy Sci., 6: 72-78.

671120ja

Al-Baarri, A.N., M. Ogawa, T. Visalsok and S. Hayakawa, 2012. Lactoperoxidase immobilized onto various beads for producing natural preservatives solution. J. Applied Food Technol., 1: 4-6.

1174455ja

Amersham-Bioscience, 2001. Use of sodium hydroxide for cleaning and sanitizing chromatography media and systems. Application Note 18-1124-57 AD, Process Chromatography, Amersham Bioscience, USA.

62223an

Oghaiki, N.A., F. Fonteh, P. Kamga, S. Mendi and H. Imele, 2007. Activation of the lactoperoxidase system as a method of preserving raw milk in areas without cooling facilities. Afr. J. Food Agric. Nutr. Dev., 7: 1-14.

42889ja

Boots, J.W. and R. Floris, 2006. Lactoperoxidase: From catalytic mechanism to practical applications. Int. Dairy J., 16: 1272-1276.

595412ja

FAO/WHO, 2005. Benefits and potential risks of the lactoperoxidase system of raw milk preservation. Report of an FAO/WHO Technical Meeting, November 28-December 2, 2005, FAO Headquarters, Rome, Italy, pp: 1-73.

58223an

Fee, C.J. and A. Chand, 2006. Capture of lactoferrin and lactoperoxidase from raw whole milk by cation exchange chromatography. Separation Purification Technol., 48, 143-149.

1174461ja

Fweja, L.W.T., M.J. Lewis and A.S. Grandison, 2008. Challenge testing the lactoperoxidase system against a range of bacteria using different activation agents. J. Dairy Sci., 91: 2566-2574.

433822ja

Hayashi, M., S. Naknukool, S. Hayakawa, M. Ogawa and A.B.A. Ni'matulah, 2012. Enhancement of antimicrobial activity of a lactoperoxidase system by carrot extract and β -carotene. Food Chem., 130: 541-546.

1174469ja

Seifu, E., E.M. Buys and E.F. Donkin, 2004. Quality aspects of Gouda cheese made from goat milk preserved by the lactoperoxidase system. Int. Dairy J., 14: 581-589.

1181348ja

Seifu, E., E.M. Buys and E.F. Donkin, 2005. Significance of the lactoperoxidase system in the dairy industry and its potential applications: A review. Trends Food Sci. Technol., 16: 137-154.

571875ja

Singh, A.K., N. Singh, S. Sharma, K. Shin and M. Takase et al., 2009. Inhibition of lactoperoxidase by its own catalytic product: Crystal structure of the hypothiocyanate-inhibited bovine lactoperoxidase at 2.3-Å resolution. Biophys. J., 96: 646-654.

568418ja

Touch, V., S. Hayakawa, S. Yamada and S. Kaneko, 2004. Effects of a lactoperoxidase-thiocyanate-hydrogen peroxide system on *Salmonella enteritidis* in animal or vegetable foods. Int. J. Food Microbiol., 93: 175-183.

78030ja

Zhou, Y. and L.T. Lim, 2009. Activation of lactoperoxidase system in milk by glucose oxidase immobilized in electrospun polylactide microfibers. J. Food Sci., 74: C170-C176.

568445ja

Shahzaib 28-8-2013, **{Add by saira: 28-08-13}**

Hypered by: Hina 28-8-13

Figure 1. Immobilization efficiency (IE) of LPO on two types of resin, i.e. SP Sepharose Big Beads (SPBB) and SP Sepharose Fast Flow (SPFF). Two types of resin (0.1–1.0 g) was used to immobilized 1 ml of LPO (35 U/ml). Flow rate has been set to 1 ml/min using peristaltic pump. Values are means of three sets of experiments. Error bars represent standard errors of the mean.

Commented [ANA21]: Table 1 has been deleted and the content has been explained in the text.

Commented [O22]: Define all abbreviations used in figures in their respective captions.

Commented [O23]: The given captions of your figures 1 and 2 have not been written properly. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such captions for all these figures in which all of their main parts have been properly explained.

Formatted: Indent: Left: 0 cm, First line: 0 cm

Deleted:

Deleted:

Deleted: Figure 1. Immobilization efficiency of SPBB and SPFF using various weight.

Figure 2a. The percentage of remaining LPO activity attached onto SP Sepharose Big Beads (SPBB) during 10 days of storage at 10°C in various storage solutions. This experiment used 1 g of SPBB for immobilizing 1 ml of LPO (35 U/ml). Immediately after immobilization, SPBB then was stored in sterilized pure water, phosphate buffer, milk, and whey for 10 days at 10°C. Values are mean of three sets of experiments. Error bars represent standard errors of the mean

Figure 2b The percentage of remaining LPO activity attached onto SP Sepharose Fast Flow (SPFF) during 10 days of storage at 10°C in various storage solution. This experiment used 1 g of SPFF for immobilizing 1 ml of LPO (35 U/ml). After immobilization, SPFF was stored in sterilized pure water, phosphate buffer, milk, and whey. Values are mean of three sets of experiments. Error bars represent standard errors of the mean

Formatted: Indent: Left: 0 cm, First line: 0 cm

Deleted:

Commented [FT26]: Assign an alphabet to each sub-part of Figure 2 and explain each figure part in figure caption by referring to its respective alphabet. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such caption in which all main parts of this figure have been properly explained. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Commented [O27]: In submitted manuscript Figure 2 has been presented but Author didn't cite this figure in text. Cite this figure in text at appropriate place with suitable explanation and make sure that all figures have been cited in text in consecutive numerical order. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Commented [ANA28]: Self explanatory has been added

Commented [ANA29]: This figure has been presented in the article.

Deleted: Figure 2a. The percentage of LPO activity attached onto SPBB during ten days of storage in various storage solutions

Deleted: Figure 2b The percentage of LPO activity attached onto SPFF during ten days of storage in various storage solution.

Resistance of Immobilized Lactoperoxidase Activity from Bovine Whey Against Storage Solutions

¹Dwi Novrina Nawangsari, ²Ahmad Ni'matullah Al-Baarri, ²Sri Mulyani, ²Anang Mohamad Legowo and ²V. Priyo Bintoro

¹Department of Animal Product Technology,

²Department of Food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

Corresponding Author: Ahmad Ni'matullah Al-Baari, Department of food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

ABSTRACT

Lactoperoxidase (LPO) could be simply obtained from whey through immobilization using a cation exchange resin of SP Sepharose. LPO received high attention since the antimicrobial properties of LPO system (LPOS) that are consisted of LPO, SCN^- and H_2O_2 was able to generate OSCN^- for strong antimicrobial agents. This study was done to analyze the immobilization efficiency of LPO onto two types of sepharose: SP-Sepharose Fast Flow (SPFF) and SP-Sepharose Big Beads (SPBB). The remaining of LPO's activity (%) against storage solution was also observed. The whey was obtained from bovine skimmed milk that was coagulated using rennet and acid lactic. The LPO was obtained from whey using SPFF. To analyze the resistance of enzyme activity, the immobilized LPO was stored in pure water, phosphate buffer, milk and whey at 10°C. The activity of LPO was monitored for 10 days. The result indicates that the LPO could be purified from whey and single band has been detected using electrophoresis method. The obtained LPO (35 U mL^{-1}) was attached onto SPFF and SPBB. Maximum immobilization efficiency has been achieved by 0.6 g SP-FF and 0.9 g SP-BB. LPO activity of the immobilized LPO were able to be kept until 5 days when it was stored in whey. Other storage solution remained various LPO activity during storage. This study concluded that IE of LPO attached onto sepharoses might be reached variously depended on the sepharose type and whey as LPO source might be used for maintaining LPO activity.

Key words: Lactoperoxidase, SP sepharose fast flow, SP sepharose big beads, immobilization, remaining activity

INTRODUCTION

Lactoperoxidase (LPO), together with SCN^- and H_2O_2 have been understood to generates intermediate product of OSCN^- as antibacterial agent that has a broad spectrum of antimicrobial effects against bacteria, fungi and viruses. This antibacterial agent could be produced if these three components exist in the medium (Seifu *et al.*, 2005). LPOS has been widely used as a preservative in dairy products and nondairy products (Seifu *et al.*, 2004; Touch *et al.*, 2004; FAO/WHO, 2005; Boots and Floris, 2006; Oghaiki *et al.*, 2007; Fweja *et al.*, 2008; Al-Baarri *et al.*, 2011a).

It has been understood that whey contains large number of LPO therefore purification method of LPO from whey has been well developed (Touch *et al.*, 2004; Zhou and Lim, 2009;

Al-Baarri *et al.*, 2010). SP-Sepharose has been known to provide beneficial effect for the immobilization efficiency since SP-Sepharose almost completely immobilized LPO and reusable (Al-Baarri *et al.*, 2010). Although, LPO was widely used in food application but it still remained the problem of its expensiveness, therefore the immobilization of LPO was needed for the reuse of LPO.

SP Sepharose has been known as immobilization agent for capturing lactoperoxidase (Fee and Chand, 2006; Hayashi *et al.*, 2012). SP Sepharose Fast Flow (SPFF) and SP Sepharose Big Beads (SPBB) were the common resin for immobilization since they provide simply application, long term of use and easy for reuse (Amersham-Bioscience, 2001). If compare to other immobilization agent such as chitosan, SP Sepharoses showed higher capturization of lactoperoxidase (Al-Baarri *et al.*, 2012). In the other hand, SP Sepharose application for immobilization of LPO resulting in the much more expensive of the use of LPO, therefore the efficient use of SP Sepharose to immobilize LPO is required. Based on related literatures, there was no documentation for the efficient use of LPO immobilization using SP Sepharose, therefore this research has been done for analyzing the maximum capturization of LPO onto SP Sepharose. This information might provide the benefit for minimum use of SP Sepharose for LPO immobilization. Since the immobilized LPO allowed the reuse of enzyme, the appropriate storage solution for keeping the enzyme activity is needed. To answer this, this research has also been done for analyzing the remaining of LPO activity after storage.

MATERIALS AND METHODS

Materials: SP Sepharose Fast Flow and SP Sepharose Big Beads were purchased from Amersham Pharmacia Biotech, Sweden (Lot. No. 10029743 and 10081054, respectively). Microbial derived rennet was purchased from Singapore and ABTS or 2,20-azino-bis (3-ethylbenz-thiazoline-6-sulphonic acid) was obtained from Kagawa Science (Lot No. 7ROZC-EC) Tokyo Chemical, Industry Co. Ltd., Japan. Cow's milk was obtained from Faculty of Animal and Agricultural Sciences's farm, Diponegoro University, Semarang, Indonesia. The spectrophotometer (Mini UV-1800, Shimadzu, Japan) was used for analysis enzyme activity. Unless other compounds specified, all other compounds were reagent grade.

Preparation of whey: The whey was prepared as method performed by Al-Baarri *et al.* (2011b) without any modification.

LPO immobilization from whey: The procedure for immobilization of LPO was conducted as the method that was performed by Al-Baarri *et al.* (2010) with minor modifications. SP Sepharose Fast Flow (SPFF) was used as agent for LPO immobilization from whey. Whey at the volume of 1800 mL was eluted through a glass column (3×40 cm) filled with 60 g of SPFF. Prior to elution, SPFF was washed with 300 mL of Phosphate Buffer (PB) (pH 6.8) containing 1 M NaCl to remove unnecessary compounds. The whey was circulated through the column using feedback tubing and a peristaltic pump. The circulation was done at the flow rate of 1.0 mL min⁻¹. After draining the whey away, the resin was washed with 300 mL of 0.4 mM NaCl in 0.1 mM phosphate buffer (pH 7.0) using fraction collector (10 mL per tube) to obtain the solution containing high concentration of LPO. Three groups of fractions (fraction number 1-10, 11-20, 21-30) were analyzed for protein profile using sodium dodecyl sulfate polyacrylamide gel electrophoresis

(SDS-PAGE) to check the purity. Finally, based on the SPS PAGE analysis, the fraction number 21-30 was selected for LPO activity analysis (LPO activity was 35 U mL⁻¹). Thus, this fraction was used throughout experiment.

Determination of captured LPO onto SP sepharoses: SPFF and SPBB (0.1-1.0 g) were washed in 1 M NaCl in PB pH 7.0 and then were placed in the column (1×10 cm). The immobilization process was started with the elution of 1 mL of LPO through column. The flow rate was set into 1 mL min⁻¹ using peristaltic pump. The output was collected for measurement of remained LPO activity in the SP Sepharoses. This experiment was repeated three times and column were wash with serial elution of 1 M NaOH and pure water, respectively. The immobilized LPO was stored in pure water, PB, milk and whey. All storage solutions were sterilized using autoclave at 110°C for 10 min. Immobilized LPO was stored at 4°C for 10 days. The remaining of LPO activity immobilized onto SP-Sepharose was calculated by eluting immobilized LPO using 1 M NaCl in PB pH 7.0. The LPO activity in the elution was analyzed spectrometrically.

LPO activity determination: LPO activity was performed as the following method: 450 µL of 1.0 mM ABTS in 10 mM acetate buffer (pH 4.4) and 450 µL of 0.55 mM H₂O₂ in pure water were gently poured into the cuvette. The enzyme (50 µL) was subsequently added to cuvette. The increase of absorbance at 412 nm measured for 20 sec. One unit of LPO enzymatic activity was expressed as the amount of enzyme needed to oxidize 1 µmol ABTS min⁻¹. The molar extinction coefficient of ABTS at 412 nm was 32.400 M⁻¹ cm⁻¹ (Touch *et al.*, 2004).

Immobilization efficiency: The immobilization efficiency (IE) was calculated as follows:

$$IE (\%) = \frac{E_1}{E_0} \times 100$$

where, E₀ is the LPO activity added to the SP Sepharoses (U mL⁻¹) and E₁ is the LPO activity embedded in the SP Sepharose (U mL⁻¹) (Al-Baarri *et al.*, 2010).

RESULTS AND DISCUSSION

Purification LPO: Whey has a lot of enzymes and it is available in low cost because whey is by product of dairy manufacture so it is the challenge to use whey as enzyme sources including LPO. The LPO activity and band(s) of the solution obtained from the elution of 1 M NaCl in PB pH 7.0 through SPFF containing LPO were checked using spectrophotometer and SDS PAGE, respectively. As mention in methods, SPFF containing LPO was generated from whey that was eluted through SPFF column. The result of LPO activity was 27.7±2.9, 39.5±4.5 and 35.2±3.4 U mL⁻¹ for fraction number 1-10, 11-20, 21-30, respectively (data not presented). The highest of LPO activity was group of fraction number 11-20, however since the band of this group showed two bands indicating two protein was detected, for whole of experiment, group of fraction number 21-30 was used. This group showed single band indicating only LPO that was captured by SPFF.

In this research SPFF was used to obtain LPO since this ion exchange resin has diameter 45-165 µm resulting in the wider of surface area than SPBB (Amersham-Bioscience, 2001). In line with this result, Touch *et al.* (2004) used SPFF for purifying LPO from whey resulting in the good

Fig. 1: Immobilization efficiency (IE) of LPO on two types of resin, i.e., SP Sepharose Big Beads (SPBB) and SP Sepharose Fast Flow (SPFF). Two types of resin (0.1-1.0 g) was used to immobilized 1 mL of LPO (35 U mL^{-1}). Flow rate has been set to 1 mL min^{-1} using peristaltic pump. Values are means of three sets of experiments, Error bars represent standard errors of the mean

ability to catch LPO (108 U mL^{-1}). The activity of LPO in this research was less than that of other researcher since the absence of microfiltration step in this research. It has been known that the microfiltration might concentrate the enzyme resulting in the high activity of LPO.

Immobilization efficiency: Immobilization efficiency plays an important role for determination of immobilization agent. This research determined Immobilization Efficiency (IE) of LPO using SPFF dan SPBB (Fig. 1). The volume of SP Sepharoses used in this experiment had a range from 0.1 to 1.0 g to catch the LPO at initial activity of $35.2 \pm 3.4 \text{ U mL}^{-1}$. The increase of IE was found as an increase of SP Sepharoses's weight. As describe in Fig. 1, when the 0.6 g of SPFF was employed, the IE achieved 100% indicating all of LPO employed was able to be captured by SPFF. When the weight of SPFF was increased, the IE was in steady state maximumly. An increase SPBB from 0.1 to 1.0 g elevated the IE from 38.6 to 100%. However, 0.9 g of SPBB completely captured LPO resulting the IE of 100%. One gram of SPFF was reported to have a maximum capture of LPO in 300 mL whey (equal to 750 U mL^{-1} LPO activity) (Al-Baarri *et al.*, 2010). This can be explained that the capture depended on the quantity of enzymes per milliliter. This research used high activity of LPO resulting in the loss of LPO activity.

Since the SPFF and SPBB provided the maximum IE at 0.6 and 0.9 g, respectively, thus these amounts of SP Sepharoses has been used in the rest of experiment. The LPO immobilized onto SPFF and SPBB was stored in the various storage solutions: Pure water, PB, milk and whey for 10 days at 10°C .

Remaining LPO activity during storage: The percentage of remaining immobilized LPO activity stored at 10°C for 10 days in various storage solutions is shown in Fig. 2a and b. LPO activity of immobilized LPO was measured after purging the LPO attached onto SP Sepharose with 1 M NaCl in PB pH 7.0. The percentage of remaining LPO activity was determined by comparing the LPO activity after storage to the initial immobilized LPO at first day of storage.

The percentage of LPO activity attached onto SPBB and SPFF is shown on Fig. 2a and b, respectively. Based on Fig. 2a, whey was able to maintain 100% enzyme activity of LPO attached onto SPBB within 4 days. The extension of storage time resulted in the remarkable

Fig. 2(a-b): (a) Percentage of remaining LPO activity attached onto SP Sepharose Big Beads (SPBB) during 10 days of storage at 10°C in various storage solutions. This experiment used 1 g of SPBB for immobilizing 1 mL of LPO (35 U mL⁻¹). Immediately after immobilization, SPBB then was stored in sterilized pure water, phosphate buffer, milk and whey for 10 days at 10°C. Values are mean of three sets of experiments. Error bars represent standard errors of the mean and (b) Percentage of remaining LPO activity attached onto SP Sepharose Fast Flow (SPFF) during 10 days of storage at 10°C in various storage solution. This experiment used 1 g of SPFF for immobilizing 1 mL of LPO (35 U mL⁻¹). After immobilization, SPFF was stored in sterilized pure water, phosphate buffer, milk and whey. Values are mean of three sets of experiments. Error bars represent standard errors of the mean

reduction of remaining LPO activity. Milk was able to maintain the LPO activity at seven days of storage even though the remaining of LPO activity at that time was very negligible in amount (2.3%).

As previously mentioned, whey was able to completely keep LPO activity within 4 days of storage. This can be explained that whey components support the activity of LPO. It has been studied that LPO activity might be inhibited by casein (Singh *et al.*, 2009) while the casein has been removed from whey.

The remaining of LPO activity attached onto SPFF during 10 days of storage at 10°C is shown on Fig. 2b. PB was able to maintain 100% of LPO activity within 5 days of storage. The milk and whey were able to keep 100% of LPO activity within 3 days of storage. The elevation of storage time until 5 days in whey could keep enzyme activity although the remaining enzyme activity was small amount (10.08%). Based on the availability, also since LPO was simply derived from whey, the storage of immobilized LPO in whey should keep its activity up to 5 days of storage. Therefore it is suggested that immobilized LPO should be stored in whey.

CONCLUSION

The results can be concluded that LPO could be purified from whey using fraction number 21-30. One milliliter of LPO (35 U mL^{-1}) could be completely immobilized onto 0.6 g SPFF or 0.9 g SPBB (immobilization efficiency was 100%). Among various storage solution, whey was able to keep 100% of LPO activity up to 5 days of storage.

ACKNOWLEDGMENTS

This study was supported by funds from the Ministry of Research and Technology of Republic of Indonesia for the project "Insentif Riset SINAS" with Grant Project No. RT-2013-2163.

REFERENCES

- Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2010. Scale-up studies on immobilization of lactoperoxidase using milk whey for producing antimicrobial agent. *J. Indonesian Trop. Anim. Agric.*, 35: 185-191.
- Al-Baarri, A.N., M. Hayashi, M. Ogawa and S. Hayakawa, 2011a. Effects of mono-and disaccharides on the antimicrobial activity of bovine lactoperoxidase system. *J. Food Prot.*, 74: 134-139.
- Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2011b. Application of lactoperoxidase system using bovine whey and the effect of storage condition on lactoperoxidase activity. *Int. J. Dairy Sci.*, 6: 72-78.
- Al-Baarri, A.N., M. Ogawa, T. Visalsok and S. Hayakawa, 2012. Lactoperoxidase immobilized onto various beads for producing natural preservatives solution. *J. Applied Food Technol.*, 1: 4-6.
- Amersham-Bioscience, 2001. Use of sodium hydroxide for cleaning and sanitizing chromatography media and systems. Application Note 18-1124-57 AD, Process Chromatography, Amersham Bioscience, USA. <http://risk.management6.com/Use-of-sodium-hydroxide-for-cleaning-and-sanitizing-chromatography-download-w15138.pdf>.
- Boots, J.W. and R. Floris, 2006. Lactoperoxidase: From catalytic mechanism to practical applications. *Int. Dairy J.*, 16: 1272-1276.
- FAO/WHO, 2005. Benefits and potential risks of the lactoperoxidase system of raw milk preservation. Report of an FAO/WHO Technical Meeting, November 28-December 2, 2005, FAO Headquarters, Rome, Italy, pp: 1-73.
- Fee, C.J. and A. Chand, 2006. Capture of lactoferrin and lactoperoxidase from raw whole milk by cation exchange chromatography. *Separation Purification Technol.*, 48: 143-149.
- Fweja, L.W.T., M.J. Lewis and A.S. Grandison, 2008. Challenge testing the lactoperoxidase system against a range of bacteria using different activation agents. *J. Dairy Sci.*, 91: 2566-2574.
- Hayashi, M., S. Naknukool, S. Hayakawa, M. Ogawa and A.B.A. Ni'matullah, 2012. Enhancement of antimicrobial activity of a lactoperoxidase system by carrot extract and α -carotene. *Food Chem.*, 130: 541-546.
- Oghaiki, N.A., F. Fonteh, P. Kamga, S. Mendi and H. Imele, 2007. Activation of the lactoperoxidase system as a method of preserving raw milk in areas without cooling facilities. *Afr. J. Food Agric. Nutr. Dev.*, 7: 1-14.
- Seifu, E., E.M. Buys and E.F. Donkin, 2004. Quality aspects of Gouda cheese made from goat milk preserved by the lactoperoxidase system. *Int. Dairy J.*, 14: 581-589.

- Seifu, E., E.M. Buys and E.F. Donkin, 2005. Significance of the lactoperoxidase system in the dairy industry and its potential applications: A review. *Trends Food Sci. Technol.*, 16: 137-154.
- Singh, A.K., N. Singh, S. Sharma, K. Shin and M. Takase *et al.*, 2009. Inhibition of lactoperoxidase by its own catalytic product: Crystal structure of the hypothiocyanate-inhibited bovine lactoperoxidase at 2.3-Å resolution. *Biophys. J.*, 96: 646-654.
- Touch, V., S. Hayakawa, S. Yamada and S. Kaneko, 2004. Effects of a lactoperoxidase-thiocyanate-hydrogen peroxide system on *Salmonella enteritidis* in animal or vegetable foods. *Int. J. Food Microbiol.*, 93: 175-183.
- Zhou, Y. and L.T. Lim, 2009. Activation of lactoperoxidase system in milk by glucose oxidase immobilized in electrospun polylactide microfibers. *J. Food Sci.*, 74: C170-C176.

Resistance of Immobilized Lactoperoxidase Activity from Bovine Whey Against Storage Solutions

¹Dwi Novrina Nawangsari, ²Ahmad Ni'matullah Al-Baarri, ²Sri Mulyani,

²Anang Mohamad Legowo and ²V. Priyo Bintoro

The complete address for author 1: Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

¹Department of Animal Product Technology,

²Department of Food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

Corresponding author should be addressed to: Ahmad Ni'matullah Al-Baarri, Department of Food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia (albari@undip.ac.id)

Corresponding Author: Dwi Novrina Nawangsari, Department of Animal Product Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia

ABSTRACT

Lactoperoxidase (LPO) could be simply obtained from whey through immobilization using a cation exchange resin of SP Sepharose. LPO received high attention since the antimicrobial properties of LPO system (LPOS) that are consisted of LPO, SCN^- and H_2O_2 was able to generate OSCN^- for strong antimicrobial agents. This study was done to analyze the immobilization efficiency of LPO onto two types of sepharose: SP-Sepharose Fast Flow (SPFF) and SP-Sepharose Big Beads (SPBB). The remaining of LPO's activity (%) against storage solution was also observed. The whey was obtained from bovine skimmed milk that was coagulated using rennet and acid lactic. The LPO was obtained from whey using SPFF. To analyze the resistance of enzyme activity, the immobilized LPO was stored in pure water, phosphate buffer, milk and whey at 10°C . The activity of LPO was monitored for 10 days. The result indicates that the LPO could be purified from whey and single band has been detected using electrophoresis method. The obtained LPO (35 U mL^{-1}) was attached onto SPFF and SPBB. Maximum immobilization efficiency has been achieved by 0.6 g SP-FF and 0.9 g SP-BB. LPO activity of the immobilized LPO were able to be kept until 5 days when it was stored in whey. Other storage solution remained various LPO activity during storage. This study concluded that IE of LPO attached onto sepharoses might be reached variously depended on the sepharose type and whey as LPO source might be used for maintaining LPO activity.

Key words: Lactoperoxidase, SP sepharose fast flow, SP sepharose big beads, immobilization, remaining activity

INTRODUCTION

Lactoperoxidase (LPO), together with SCN^- and H_2O_2 have been understood to generates intermediate product of OSCN^- as antibacterial agent that has a broad spectrum of antimicrobial effects against bacteria, fungi and viruses. This antibacterial agent could be produced if these three components exist in the medium (Seifu *et al.*, 2005). LPOS has been widely used as a preservative in dairy products and nondairy products (Seifu *et al.*, 2004; Touch *et al.*, 2004; FAO/WHO, 2005; Boots and Floris, 2006; Oghaiki *et al.*, 2007; Fweja *et al.*, 2008; Al-Baarri *et al.*, 2011a).

It has been understood that whey contains large number of LPO therefore purification method of LPO from whey has been well developed (Touch *et al.*, 2004; Zhou and Lim, 2009;

Al-Baarri *et al.*, 2010). SP-Sepharose has been known to provide beneficial effect for the immobilization efficiency since SP-Sepharose almost completely immobilized LPO and reusable (Al-Baarri *et al.*, 2010). Although, LPO was widely used in food application but it still remained the problem of its expensiveness, therefore the immobilization of LPO was needed for the reuse of LPO.

SP Sepharose has been known as immobilization agent for capturing lactoperoxidase (Fee and Chand, 2006; Hayashi *et al.*, 2012). SP Sepharose Fast Flow (SPFF) and SP Sepharose Big Beads (SPBB) were the common resin for immobilization since they provide simply application, long term of use and easy for reuse (Amersham-Bioscience, 2001). If compare to other immobilization agent such as chitosan, SP Sepharoses showed higher capturization of lactoperoxidase (Al-Baarri *et al.*, 2012). In the other hand, SP Sepharose application for immobilization of LPO resulting in the much more expensive of the use of LPO, therefore the efficient use of SP Sepharose to immobilize LPO is required. Based on related literatures, there was no documentation for the efficient use of LPO immobilization using SP Sepharose, therefore this research has been done for analyzing the maximum capturization of LPO onto SP Sepharose. This information might provide the benefit for minimum use of SP Sepharose for LPO immobilization. Since the immobilized LPO allowed the reuse of enzyme, the appropriate storage solution for keeping the enzyme activity is needed. To answer this, this research has also been done for analyzing the remaining of LPO activity after storage.

MATERIALS AND METHODS

Materials: SP Sepharose Fast Flow and SP Sepharose Big Beads were purchased from Amersham Pharmacia Biotech, Sweden (Lot. No. 10029743 and 10081054, respectively). Microbial derived rennet was purchased from Singapore and ABTS or 2,20-azino-bis (3-ethylbenz-thiazoline-6-sulphonic acid) was obtained from Kagawa Science (Lot No. 7ROZC-EC) Tokyo Chemical, Industry Co. Ltd. Japan. Cow's milk was obtained from Faculty of Animal and Agricultural Sciences's farm, Diponegoro University, Semarang, Indonesia. The spectrophotometer (Mini UV-1800, Shimadzu, Japan) was used for analysis enzyme activity. Unless other compounds specified, all other compounds were reagent grade.

Preparation of whey: The whey was prepared as method performed by Al-Baarri *et al.* (2011b) without any modification.

LPO immobilization from whey: The procedure for immobilization of LPO was conducted as the method that was performed by Al-Baarri *et al.* (2010) with minor modifications. SP Sepharose Fast Flow (SPFF) was used as agent for LPO immobilization from whey. Whey at the volume of 1800 mL was eluted through a glass column (3×40 cm) filled with 60 g of SPFF. Prior to elution, SPFF was washed with 300 mL of Phosphate Buffer (PB) (pH 6.8) containing 1 M NaCl to remove unnecessary compounds. The whey was circulated through the column using feedback tubing and a peristaltic pump. The circulation was done at the flow rate of 1.0 mL min⁻¹. After draining the whey away, the resin was washed with 300 mL of 0.4 mM NaCl in 0.1 mM phosphate buffer (pH 7.0) using fraction collector (10 mL per tube) to obtain the solution containing high concentration of LPO. Three groups of fractions (fraction number 1-10, 11-20, 21-30) were analyzed for protein profile using sodium dodecyl sulfate polyacrylamide gel electrophoresis

(SDS-PAGE) to check the purity. Finally, based on the SPS PAGE analysis, the fraction number 21-30 was selected for LPO activity analysis (LPO activity was 35 U mL⁻¹). Thus, this fraction was used throughout experiment.

Determination of captured LPO onto SP sepharoses: SPFF and SPBB (0.1-1.0 g) were washed in 1 M NaCl in PB pH 7.0 and then were placed in the column (1×10 cm). The immobilization process was started with the elution of 1 mL of LPO through column. The flow rate was set into 1 mL min⁻¹ using peristaltic pump. The column was washed with serial elution of 1 M NaOH and 1 M Phosphate Buffer (PB) pH 7.0, milk, and whey. The immobilized LPO was stored in milk, whey, pure water and whey. The storage bottles were sterilized using autoclave at 110°C for 10 min. Immobilized LPO was stored at 4°C for 10 days. The remaining of LPO activity immobilized onto SP-Sepharose was calculated by eluting immobilized LPO using 1 M NaCl in PB pH 7.0. The LPO activity in the elution was analyzed spectrometrically.

It should be:
pure water,
1 M Phosphate Buffer (PB) pH
7.0, milk, and whey

LPO activity determination: LPO activity was performed as the following method: 450 µL of 1.0 mM ABTS in 10 mM acetate buffer (pH 4.4) and 450 µL of 0.55 mM H₂O₂ in pure water were gently poured into the cuvette. The enzyme (50 µL) was subsequently added to cuvette. The increase of absorbance at 412 nm measured for 20 sec. One unit of LPO enzymatic activity was expressed as the amount of enzyme needed to oxidize 1 µmol ABTS min⁻¹. The molar extinction coefficient of ABTS at 412 nm was 32.400 M⁻¹ cm⁻¹ (Touch *et al.*, 2004).

Immobilization efficiency: The immobilization efficiency (IE) was calculated as follows:

$$IE (\%) = \frac{E_1}{E_0} \times 100$$

where, E₀ is the LPO activity added to the SP Sepharoses (U mL⁻¹) and E₁ is the LPO activity embedded in the SP Sepharose (U mL⁻¹) (Al-Baarri *et al.*, 2010).

RESULTS AND DISCUSSION

Purification LPO: Whey has a lot of enzymes and it is available in low cost because whey is by product of dairy manufacture so it is the challenge to use whey as enzyme sources including LPO. The LPO activity and band(s) of the solution obtained from the elution of 1 M NaCl in PB pH 7.0 through SPFF containing LPO were checked using spectrophotometer and SDS PAGE, respectively. As mention in methods, SPFF containing LPO was generated from whey that was eluted through SPFF column. The result of LPO activity was 27.7±2.9, 39.5±4.5 and 35.2±3.4 U mL⁻¹ for fraction number 1-10, 11-20, 21-30, respectively (data not presented). The highest of LPO activity was group of fraction number 11-20, however since the band of this group showed two bands indicating two protein was detected, for whole of experiment, group of fraction number 21-30 was used. This group showed single band indicating only LPO that was captured by SPFF.

In this research SPFF was used to obtain LPO since this ion exchange resin has diameter 45-165 µm resulting in the wider of surface area than SPBB (Amersham-Bioscience, 2001). In line with this result, Touch *et al.* (2004) used SPFF for purifying LPO from whey resulting in the good

Fig. 1: Immobilization efficiency (IE) of LPO on two types of resin, i.e., SP Sepharose Big Beads (SPBB) and SP Sepharose Fast Flow (SPFF). Two types of resin (0.1-1.0 g) was used to immobilized 1 mL of LPO (35 U mL^{-1}). Flow rate has been set to 1 mL min^{-1} using peristaltic pump. Values are means of three sets of experiments, Error bars represent standard errors of the mean

ability to catch LPO (108 U mL^{-1}). The activity of LPO in this research was less than that of other researcher since the absence of microfiltration step in this research. It has been known that the microfiltration might concentrate the enzyme resulting in the high activity of LPO.

Immobilization efficiency: Immobilization efficiency plays an important role for determination of immobilization agent. This research determined Immobilization Efficiency (IE) of LPO using SPFF dan SPBB (Fig. 1). The volume of SP Sepharoses used in this experiment had a range from 0.1 to 1.0 g to catch the LPO at initial activity of $35.2 \pm 3.4 \text{ U mL}^{-1}$. The increase of IE was found as an increase of SP Sepharoses's weight. As describe in Fig. 1, when the 0.6 g of SPFF was employed, the IE achieved 100% indicating all of LPO employed was able to be captured by SPFF. When the weight of SPFF was increased, the IE was in steady state maximumly. An increase SPBB from 0.1 to 1.0 g elevated the IE from 38.6 to 100%. However, 0.9 g of SPBB completely captured LPO resulting the IE of 100%. One gram of SPFF was reported to have a maximum capture of LPO in 300 mL whey (equal to 750 U mL^{-1} LPO activity) (Al-Baarri *et al.*, 2010). This can be explained that the capture depended on the quantity of enzymes per milliliter. This result was due to the loss of LPO resulting in the loss of LPO activity.

Since the SPFF and SPBB provided the maximum IE at 0.6 and 0.9 g, respectively, the same amounts of SP Sepharoses has been used in the rest of experiment. The immobilized LPO on SPFF and SPBB was stored in the various storage solutions: Pure water, PB, milk and whey for 10 days at 10°C .

It should be:
pure water,
1 M Phosphate Buffer (PB) pH
7.0, milk, and whey

Remaining LPO activity during storage: The percentage of remaining immobilized LPO activity stored at 10°C for 10 days in various storage solutions is shown in Fig. 2a and b. LPO activity of immobilized LPO was measured after purging the LPO attached onto SP Sepharose with 1 M NaCl in PB pH 7.0. The percentage of remaining LPO activity was determined by comparing the LPO activity after storage to the initial immobilized LPO at first day of storage.

The percentage of LPO activity attached onto SPBB and SPFF is shown on Fig. 2a and b, respectively. Based on Fig. 2a, whey was able to maintain 100% enzyme activity of LPO attached onto SPBB within 4 days. The extension of storage time resulted in the remarkable

Fig. 2(a-b): (a) Percentage of remaining LPO activity attached onto SP Sepharose Big Beads (SPBB) during 10 days of storage at 10°C in various storage solutions. This experiment used 1 g of SPBB for immobilizing 1 mL of LPO (35 U mL⁻¹). Immediately after immobilization, SPBB then was stored in sterilized pure water, phosphate buffer, milk and whey for 10 days at 10°C. Values are mean of three sets of experiments. Error bars represent standard errors of the mean and (b) Percentage of remaining LPO activity attached onto SP Sepharose Fast Flow (SPFF) during 10 days of storage at 10°C in various storage solution. This experiment used 1 g of SPFF for immobilizing 1 mL of LPO (35 U mL⁻¹). After immobilization, SPFF was stored in sterilized pure water, phosphate buffer, milk and whey. Values are mean of three sets of experiments. Error bars represent standard errors of the mean

reduction of remaining LPO activity. Milk was able to maintain the LPO activity at seven days of storage even though the remaining of LPO activity at that time was very negligible in amount (2.3%).

As previously mentioned, whey was able to completely keep LPO activity within 4 days of storage. This can be explained that whey components support the activity of LPO. It has been studied that LPO activity might be inhibited by casein (Singh *et al.*, 2009) while the casein has been removed from whey.

The remaining of LPO activity attached onto SPFF during 10 days of storage at 10°C is shown on Fig. 2b. PB was able to maintain 100% of LPO activity within 5 days of storage. The milk and whey were able to keep 100% of LPO activity within 3 days of storage. The elevation of storage time until 5 days in whey could keep enzyme activity although the remaining enzyme activity was small amount (10.08%). Based on the availability, also since LPO was simply derived from whey, the storage of immobilized LPO in whey should keep its activity up to 5 days of storage. Therefore it is suggested that immobilized LPO should be stored in whey.

CONCLUSION

The results can be concluded that LPO could be purified from whey using fraction number 21-30. One milliliter of LPO (35 U mL^{-1}) could be completely immobilized onto 0.6 g SPFF or 0.9 g SPBB (immobilization efficiency was 100%). Among various storage solution, whey was able to keep 100% of LPO activity up to 5 days of storage.

ACKNOWLEDGMENTS

This study was supported by funds from the Ministry of Research and Technology of Republic of Indonesia for the project "Insentif Riset SINAS" with Grant Project No. RT-2013-2163.

REFERENCES

- Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2010. Scale-up studies on immobilization of lactoperoxidase using milk whey for producing antimicrobial agent. *J. Indonesian Trop. Anim. Agric.*, 35: 185-191.
- Al-Baarri, A.N., M. Hayashi, M. Ogawa and S. Hayakawa, 2011a. Effects of mono-and disaccharides on the antimicrobial activity of bovine lactoperoxidase system. *J. Food Prot.*, 74: 134-139.
- Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2011b. Application of lactoperoxidase system using bovine whey and the effect of storage condition on lactoperoxidase activity. *Int. J. Dairy Sci.*, 6: 72-78.
- Al-Baarri, A.N., M. Ogawa, T. Visalsok and S. Hayakawa, 2012. Lactoperoxidase immobilized onto various beads for producing natural preservatives solution. *J. Applied Food Technol.*, 1: 4-6.
- Amersham-Bioscience, 2001. Use of sodium hydroxide for cleaning and sanitizing chromatography media and systems. Application Note 18-1124-57 AD, Process Chromatography, Amersham Bioscience, USA. <http://risk.management6.com/Use-of-sodium-hydroxide-for-cleaning-and-sanitizing-chromatography-download-w15138.pdf>.
- Boots, J.W. and R. Floris, 2006. Lactoperoxidase: From catalytic mechanism to practical applications. *Int. Dairy J.*, 16: 1272-1276.
- FAO/WHO, 2005. Benefits and potential risks of the lactoperoxidase system of raw milk preservation. Report of an FAO/WHO Technical Meeting, November 28-December 2, 2005, FAO Headquarters, Rome, Italy, pp: 1-73.
- Fee, C.J. and A. Chand, 2006. Capture of lactoferrin and lactoperoxidase from raw whole milk by cation exchange chromatography. *Separation Purification Technol.*, 48: 143-149.
- Fweja, L.W.T., M.J. Lewis and A.S. Grandison, 2008. Challenge testing the lactoperoxidase system against a range of bacteria using different activation agents. *J. Dairy Sci.*, 91: 2566-2574.
- Hayashi, M., S. Naknukool, S. Hayakawa, M. Ogawa and A.B.A. Ni'matullah, 2012. Enhancement of antimicrobial activity of a lactoperoxidase system by carrot extract and α -carotene. *Food Chem.*, 130: 541-546.
- Oghaiki, N.A., F. Fonteh, P. Kamga, S. Mendi and H. Imele, 2007. Activation of the lactoperoxidase system as a method of preserving raw milk in areas without cooling facilities. *Afr. J. Food Agric. Nutr. Dev.*, 7: 1-14.
- Seifu, E., E.M. Buys and E.F. Donkin, 2004. Quality aspects of Gouda cheese made from goat milk preserved by the lactoperoxidase system. *Int. Dairy J.*, 14: 581-589.

- Seifu, E., E.M. Buys and E.F. Donkin, 2005. Significance of the lactoperoxidase system in the dairy industry and its potential applications: A review. *Trends Food Sci. Technol.*, 16: 137-154.
- Singh, A.K., N. Singh, S. Sharma, K. Shin and M. Takase *et al.*, 2009. Inhibition of lactoperoxidase by its own catalytic product: Crystal structure of the hypothiocyanate-inhibited bovine lactoperoxidase at 2.3-Å resolution. *Biophys. J.*, 96: 646-654.
- Touch, V., S. Hayakawa, S. Yamada and S. Kaneko, 2004. Effects of a lactoperoxidase-thiocyanate-hydrogen peroxide system on *Salmonella enteritidis* in animal or vegetable foods. *Int. J. Food Microbiol.*, 93: 175-183.
- Zhou, Y. and L.T. Lim, 2009. Activation of lactoperoxidase system in milk by glucose oxidase immobilized in electrospun polylactide microfibers. *J. Food Sci.*, 74: C170-C176.

Dear The Editor of
International Journal of Dairy Science

Thank you for your kind to process our article to publish at your journal. We received the final proof of our article entitled: "Resistance of Immobilized Lactoperoxidase Activity from Bovine Whey Against Storage Solutions" (article number 57374) for second time.

In this time, we could not see our revision (the article seems similar to previous article). Through this letter, we hope the correction for our final proof's article are able to be received.

Here we listed the correction for final proof.

Position	It was written	It should be written
Page 1 on the Corresponding Author	Dwi Novrina Nawangsari, Department of Animal Product Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia	Ahmad Ni'matullah Al-Baarri, Department of Food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Indonesia (albari@undip.ac.id)
Page 3 on the sub title: Determination of Captured LPO ...	The immobilized LPO was stored in milk, whey, pure water and whey	The immobilized LPO was stored in pure water, PB, milk, and whey

Again, thank you very much for your attention and we hope this correction are able to be implemented in our article.

Sincerely,

Ahmad Ni'matullah Al-Baarri
Author

SCIENCE ALERT

www.scialert.com

P.O.Box 126208 Deira Dubai, UAE.

Jan 23, 2014

Mr. Ahmad N AL-Baarri,
Diponegoro University

Subject: Acceptance Letter for Article No. 57374-IJDS-AJ

It's a great pleasure for us to inform you that below mentioned manuscript has been accepted for publication in International Journal of Dairy Science as Research Article on the recommendation of the reviewers.

Title: RESISTANCE OF IMMOBILIZED LACTOPEROXIDASE ACTIVITY FROM BOVINE WHEY AGAINST STORAGE SOLUTIONS

Author's Name: Dwi Novrina Nawangsari, Ahmad Nimatullah Al-Baarri and Sri Mulyani

Receiving Date: August 02, 2013

Revised on: October 26, 2013

Accepted on: October 31, 2013

Published on: January 20, 2014

Regards

M. Imran Pasha
Publication Manager

RESISTANCE OF IMMOBILIZED LACTOPEROXIDASE ACTIVITY FROM BOVINE WHEY AGAINST STORAGE SOLUTIONS

by Dwi Novrina Nawangsari

Submission date: 04-Sep-2019 02:55AM (UTC+0700)

Submission ID: 1166759461

File name: 57374-IJDS-AJ-evaluated-Hypered_RAW_MANUSCRIPT.doc (331.5K)

Word count: 2915

Character count: 15230

To: International Journal of Dairy Science

Formatted: Left

We have received the response from reviewer 1 and reviewer 2 to our article, however we found the question from reviewer 2 only:

Al-Baarri, 2011 missed. Also the result of LPO activity why not presented in a table
Also The captions of the figures must be written at the top of the figures and
must be clear and explain the figures.

Formatted: Left, Indent: Left: 0.39", Right: 0.76"

Formatted: Font: Not Bold

Formatted: Left

Our responses:

- Al-Baarri, 2011 has been changed to Al-Baarri, 2011a. We are sorry, it should be added with "a" after "2011"
- LPO activity did not appear in a table since we would like to show the differences between treatments (storage). We believe that the description through percentage in a figure is much easier to be understood
- The caption has been moved into the top of figure and the explanation of figure has been completed.

Formatted: Left, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Left

Sincerely,

Ahmad Ni'matullah Al-Baarri

Formatted: Font: Not Bold

[Last update on August 10, 2011](#)

RESISTANCE OF IMMOBILIZED LACTOPEROXIDASE ACTIVITY FROM BOVINE WHEY AGAINST STORAGE SOLUTIONS

Dwi Novrina Nawangsari¹⁾, Ahmad Ni'matullah Al-Baarri²⁾, and Sri Mulyani²⁾, Anang Mohamad Legowo²⁾, V. Privo Bintoro²⁾

1)Department of Animal Product Technology; Faculty of Animal and Agricultural Sciences; Diponegoro University, Semarang, Indonesia

2)Department of Food Technology, Faculty of Animal and Agricultural Sciences; Diponegoro University, Semarang, Indonesia

ABSTRACT

Lactoperoxidase (LPO) could be simply obtained from whey through immobilization using a cation exchange resin of SP Sepharose. LPO received high attention since the antimicrobial properties of LPO system (LPOS) that are consisted of LPO, SCN⁻, and H₂O₂ was able to generate OSCN⁻ for strong antimicrobial agents. This study was done to analyze the immobilization efficiency of LPO onto two types of sepharose: SP-Sepharose Fast Flow (SPFF) and SP-Sepharose Big Beads (SPBB). The remaining of LPO's activity (%) against storage solution was also observed. The whey was obtained from bovine skimmed milk that was coagulated using rennet and acid lactic. The LPO was obtained from whey using SPFF. To analyze the remaining-resistence of immobilized LPO-enzyme activity, the immobilized LPO was stored in pure water, phosphate buffer, milk, and whey at 10°C. The activity of LPO was monitored for 10 days. The result indicates that the LPO could be purified from whey and single band has been detected using electrophoresis method. The obtained LPO (35 U/ml) was attached onto SPFF and SPBB. Maximum immobilization efficiency has been achieved. It was concluded that by 0.6 g SP-FF and 0.9 g SP-BB were able to achieved 100% of immobilization efficiency (IE). LPO activity of the immobilized LPO onto Sepharoses were able to be kept until 5 days when it was stored in whey. Other storage solution remained various LPO activity during storage. This research concluded that IE of LPO attached onto sepharoses might be reached variously depended on the sepharose type and whey as LPO source might be used for maintaining LPO activity.

Key words: Lactoperoxidase, SP Sepharose Fast Flow, SP Sepharose Big Beads, immobilization, remaining activity.

Commented [ANA1]: Please allow us to add the co-authors since this author has a big contribution to our experiments and article writing. The contribution of authors have been mentioned in cover letter.

Commented [O2]: The cover letter for this manuscript is not provided by the author. Download sample cover letter from the link and submit duly filled Cover Letter with all your new submissions <http://scialert.net/coverletter.doc>

Commented [ANA3]: Cover letter has been made and author has sent this cover letter along with this article.

Commented [O4]: End the paragraph of abstract section with a clear cut conclusion of study, but length of abstract section should not exceed from more than 250 words.

Commented [ANA5]: Cover letter has been added and sent via email

Commented [O6]: Check the spelling of **yellow** highlighted words.

Commented [ANA7]: Conclusion has been added and the total word counts is 245 (less than 250)

INTRODUCTION

Lactoperoxidase (LPO), together with SCN^- and H_2O_2 have been understood to generates intermediate product of OSCN^- as antibacterial agent that has a broad spectrum of antimicrobial effects against bacteria, fungi and viruses. This antibacterial agent could be produced if these three components exists in the medium (Seifu et al., 2005, Al-Baarri, 2011a). LPOS has been widely used as a preservative in dairy products and nondairy products (Seifu et al., 2004, Touch et al., 2004, FAO/WHO, 2005, Boots and Floris, 2006, Oghaiki et al, 2007, Fweja et al., 2008, Al-Baarri et al., 2011a).

It has been understood that whey contains large number of LPO therefore purification method of LPO from whey has been well developed (Touch et al., 2004, Zhou and Lim, 2009, Al-Baarri et al., 2010). SP-Sepharose has been known to provide beneficial effect for the immobilization efficiency since SP-Sepharose almost completely immobilized LPO and reusable (Al-Baarri et al., 2010). Although LPOS was widely used in food application but it still remained the problem of its expensiveness, therefore the immobilization of LPO was needed for the reuse of LPO.

SP Sepharose has been known as appropriate-immobilization agent for capturing lactoperoxidase (Fee and Chand, 2006; Hayashi et al., 2012). SP Sepharose Fast Flow (SPFF) and SP Sepharose Big Beads (SPBB) were the common resin for immobilization since they provide simply application, long term of use, and easy for reuse (Amersham-Bioscience, 2001). If compare to other immobilization agent such as chitosan, SP Sepharoses showed higher capturization of lactoperoxidase (Al-Baarri et al., 2012). In the other hand, SP Sepharose application for immobilization of LPO resulting in the much more expensive of the use of LPO, therefore the efficient use of SP Sepharose to immobilize LPO is required. Based on related literatures-, there was no documentation for the efficient use of LPO immobilization using SP Sepharose, therefore this research has been done for analyzing the maximum capturization of LPO onto SP Sepharose. This information might provide the benefit for minimum use of SP Sepharose for LPO immobilization. Since the immobilized LPO allowed the reuse of enzyme, the appropriate storage solution for keeping the enzyme activity is needed. To answer this, this research has also been done for analyzing the remaining of LPO activity after storage.

MATERIALS AND METHODS

descriptive analysis was used for describing activity of LPO in all treatment.

Commented [08]: Don't use personal pronouns throughout the text.

Commented [ANA9]: I do not have idea to replace this word, so, I did not change this word.

Commented [O10]: Provide some detail of sued statistical analysis under separate subheading, in methodology section.

SP Sepharose Fast Flow~~SP-FF~~ and SP Sepharose Big Beads ~~SP-BB~~ were purchased from Amersham Pharmacia Biotech, Sweden (Lot. No.10029743 and No. 10081054, respectively). Microbial derived rennet was purchased from Singapore and ABTS or 2,20-azino-bis (3-ethylbenz-thiazoline-6-sulphonic acid) ~~or ABTS~~ was obtained from Kagawa Science (Lot No. 7ROZC-EC) Tokyo Chemical, Industry Co. Ltd. Japan. Cow's milk was obtained from Faculty of Animal and Agricultural Sciences's farm, Diponegoro University, Semarang, Indonesia. The spectrophotometer (Mini UV-1800, Shimadzu, Japan) was used for analysis enzyme activity. Unless other compounds specified, all other compounds were reagent grade.

Preparation of whey

The whey was prepared as method performed by [Al-Baarri et. al. \(2011b\)](#) without any modification.

LPO Immobilization from Whey

The procedure for immobilization of LPO ~~immobilization~~ was conducted as the method that was performed by [Al-Baarri et al., 2010](#) with minor modifications. SP Sepharose Fast Flow (SPFF) was used as ~~a~~ agent for LPO immobilization from whey. Whey at the volume of 1800 ml was eluted through a glass column (3 x 40 cm) filled with 60 g of SP-FF. Prior to elution, SPFF was washed with 300 ml of phosphate buffer (PB) (pH 6,8) containing 1 M NaCl to remove unnecessary compounds. The whey was circulated through the column using feedback tubing and a peristaltic pump. The circulation was done at the flow rate of 1.0 ml/min. After draining the whey away, the resin was washed with 300 ml of 0.4 mM NaCl in 0.1 mM phosphate buffer (pH 7.0) using fraction collector (10 ml per tube) to obtain the solution containing high concentration of LPO. Three ~~group~~groups of fractions (fraction number 1–10, 11–20, 21–30) were analyzed for protein profile using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS PAGE) ~~SDS PAGE~~ to check the purity. Finally, based on the SPS PAGE analysis, the fraction number 21–30 was analyzed-selected for LPO activity analysis (LPO activity was 35 U/ml). Thus, this fraction ~~and~~ was used throughout experiment.

Determination of Captured LPO onto SP Sepharoses

SP-FF and SP-BB (0.1–1.0 g) were washed in 1 M NaCl in PB pH 7.0 and then were placed in the column (1x10 cm). The immobilization process was started with the elution of 1 ml of LPO through column. The flow rate was set into 1 ml/min ~~ute~~ using peristaltic pump.

Commented [ANA11]: ABTS has been replaced as start of sentence

Commented [O12]: Don't start a sentence with a numeric value.

The output was collected for measurement of remained LPO activity in the SP Sepharoses. This experiment was repeated three times and column were wash with serial elution of 1 M NaOH and pure water, respectively. The immobilized LPO was stored in milk, whey, pure water and whey. All storage solutions were sterilized using autoclave at 110°C for 10 minutes. Immobilized LPO was stored at 4°C for 10 days. The remaining of LPO activity immobilized onto SP-Sepharose was calculated by eluting immobilized LPO using 1 M NaCl in PB pH 7.0. The LPO activity in the elution was analyzed spectrometrically.

LPO Activity Determination

LPO activity was performed as the following method: 450µl of 1.0 mM ABTS in 10 mM acetate buffer (pH 4.4) and 450 µl of 0.55 mM H₂O₂ in pure water were gently poured into the cuvette. The enzyme (50 µl) was subsequently added to cuvette. The increase of absorbance at 412 nm measured for 20 second. One unit of LPO enzymatic activity was expressed as the amount of enzyme needed to oxidize 1 µmol ABTS/min. The molar extinction coefficient of ABTS at 412nm was 32.400 M⁻¹ cm⁻¹ (Touch et al., 2004).

Immobilization Efficiency

The immobilization efficiency (IE) was calculated as follows: $IE(\%) = E1/E0 \times 100$, where E0 is the LPO activity added to the SP Sepharoses (U/ml) and E1 is the LPO activity embedded in the SP Sepharose (U/ml) (Al-Baarri et al., 2010).

RESULTS AND DISCUSSION

Purification LPO

Whey has a lot of enzymes and it is available in low cost because whey is by product of dairy manufacture so it is the challenge to use whey as enzyme sources including LPO. Table 1 shows The LPO activity and band (s) of the solution obtained from the elution of 1 M NaCl in PB pH 7.0 through SPFF containing LPO were checked using spectrophotometer and SDS PAGE, respectively. As mention in methods, SPFF containing LPO was generated from whey that was eluted through SPFF column. The result of LPO activity was 27.7±2.9, 39.5±4.5, and 35.2±3.4 U/ml for fraction number 1–10, 11–20, 21–30, respectively (data not presented). As shown in Table 1 The highest of LPO activity was group of fraction number 11–20, however since the band of this group showed two bands indicating two protein was detected, for whole of experiment, group of fraction number 21–30 was used. This group showed single band indicating only LPO that was captured by SPFF.

Commented [O13]: Some numeric values mentioned in text are not in accordance with their respective tabular data. For your convenience some of those values have been yellow highlighted. Correct all data irregularities by comparing numeric values given in text with their respective tabular data.

In this research SPFF was used to obtain LPO since this ion exchange resin has diameter 45–165 μm resulting in the wider of surface area than SPBB (Amersham-Bioscience, 2001). In line with this result, Touch et al., 2004 used SPFF for purifying LPO from whey resulting in the good ability to catch LPO (108 U/ml). The activity of LPO in this research was less than that of other researcher since the absence of microfiltration step in this research. It has been known that the microfiltration might concentrate the enzyme resulting in the high activity of LPO.

Immobilization Efficiency

Immobilization efficiency plays an important role for determination of immobilization agent. This research determined immobilization efficiency (IE) of LPO using SPFF dan SPBB (Figure 1). The volume of SP Sepharoses used in this experiment had a range from 0.1 to 1.0 g to catch the LPO at initial activity of 35.2 ± 3.4 U/ml. The increase of IE was found as an increase of SP Sepharoses's weight. As describe in Figure 1, Figure 1 shows wwhen the 0.6 g of SPFF was employed, the IE achieved 100% indicating all of LPO employed was able to be captured by SPFF. When the weight of SPFF was increased, the IE was in steady state maximumly. An increase SPBB from 0.1 to 1.0 g elevated the IE from 38.6 to 100%. However, 0.9 g of SPBB completely captured LPO resulting the IE of 100%. One gram of SPFF was reported to have a maximum capture of LPO in 300 ml whey (equal to 750 U/ml LPO activity) (Al-Baarri et al., 2010). This can be explained that the capture might be depended on the quantity of enzymes per mililiter. This research used high activity of LPO resulting in the loss of LPO activity.

Since the SPFF and SPBB provided the maximum IE at 0.6 and 0.9 g, respectively, thus these amount these amounts of SP Sepharoses has been used in the rest of experiment. The LPO immobilized onto SPFF and SPBB was stored in the various storage solutions: pure water, PB, milk, and whey for 10 days in 10°C .

Remaining LPO Activity During Storage

The percentage of remaining immobilized LPO activity stored at 10°C for 10 days in various storage solutions is shown in Figure 22a and 2b. LPO activity of immobilized LPO was measured after purging the LPO attached onto SP Sepharose with 1 M NaCl in PB pH 7.0. The percentage of remaining LPO activity was determined by comparing the LPO activity after storage to the initial immobilized LPO at first day of storage.

Commented [O14]: Author has cited Figure 3 in text but didn't provide it in the submitted manuscript. Provide this missing figure and make sure that all figures have been cited in text in consecutive numerical order. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Commented [O15]: Provide correct reference of all figures in text.

Commented [ANA16]: I have double checked about this.

The percentage of LPO activity attached onto SPBB and SPFF is shown on Figure 2a and Figure 2b, respectively. Based on Figure 2a, whey was able to maintain 100% LPO enzyme activity of LPO attached onto SPBB within 4 days. The extension of storage time resulted in the remarkable reduction of remaining LPO activity. Milk was able to maintain the LPO activity at seven days of storage even though the remaining of LPO activity at that time was very negligible in amount (2.3%).

As previously mentioned, whey was able to completely keep LPO activity within 4 days of storage. This can be explained that whey components support the activity of LPO. It has been studied that LPO activity might be inhibited by casein (Singh et al., 2009) while the casein has been removed from whey.

The remaining of LPO activity attached onto SPFF during 10 days of storage at 10°C is shown on Figure 2b. PB was able to maintain 100% of LPO activity within 5 days of storage. The milk and whey were able to keep 100% of LPO activity within 3 days of storage. The elevation of storage time until 5 days in whey could keep enzyme activity although the remaining enzyme activity was small amount (10.08%). Based on the availability, also, since LPO was simply derived from whey, the storage of immobilized LPO in whey should keep its activity up to 5 days of storage. Therefore it is suggested that immobilized LPO should be stored in whey.

CONCLUSION

The results can be concluded that LPO could be purified from whey using fraction number 21–30. One milliliter of LPO (35 U/ml) could be completely immobilized onto 0.6 g SPFF or 0.9 g SPBB (immobilization efficiency was 100%). Among various storage solution, whey was able to keep 100% of LPO activity up to 5 days of storage.

ACKNOWLEDGMENT

This work was supported by funds from the Ministry of Research and Technology of Republic of Indonesia for the project "Insentif Riset SINAS" with grant project number RT-2013-2163.

REFERENCES

Al-Barazi, A.N., 2011. Lactoperoxidase Activity on Bovine Whey at Critical Temperature Storage. Unpublished data.
disaccharides on the antimicrobial activity of bovine lactoperoxidase system. J. Food Prot., 74: 134-139.

Commented [ANA17]: I have double check for the explanation of figure and proper figure

Formatted: Indent: First line: 0"

Formatted: Level 1, Indent: First line: 0"

Formatted: Indent: First line: 0"

Commented [ANA18]: Please allow us to explain the our acknowledgment

Formatted: Level 1, Indent: First line: 0"

Commented [G19]: Please remove or replace it because this reference is against format. (Unpublished data).

1174452ja

Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2010. Scale-up studies on immobilization of lactoperoxidase using milk whey for producing antimicrobial agent. J. Indonesian Trop. Anim. Agric., 35: 185-191.

1174454ja

Al-Baarri, A.N., M. Ogawa and S. Hayakawa, 2011b. Application of lactoperoxidase system using bovine whey and the effect of storage condition on lactoperoxidase activity. Int. J. Dairy Sci., 6: 72-78.

671120ja

Al-Baarri, A.N., M. Ogawa, T. Visalsok and S. Hayakawa, 2012. Lactoperoxidase immobilized onto various beads for producing natural preservatives solution. J. Applied Food Technol., 1: 4-6.

1174455ja

Amersham-Bioscience, 2001. Use of sodium hydroxide for cleaning and sanitizing chromatography media and systems. Application Note 18-1124-57 AD, Process Chromatography, Amersham Bioscience, USA.

62223an

Oghaiki, N.A., F. Fonteh, P. Kamga, S. Mendi and H. Imele, 2007. Activation of the lactoperoxidase system as a method of preserving raw milk in areas without cooling facilities. Afr. J. Food Agric. Nutr. Dev., 7: 1-14.

42889ja

Boots, J.W. and R. Floris, 2006. Lactoperoxidase: From catalytic mechanism to practical applications. Int. Dairy J., 16: 1272-1276.

595412ja

FAO/WHO, 2005. Benefits and potential risks of the lactoperoxidase system of raw milk preservation. Report of an FAO/WHO Technical Meeting, November 28-December 2, 2005, FAO Headquarters, Rome, Italy, pp: 1-73.

58223an

Fee, C.J. and A. Chand, 2006. Capture of lactoferrin and lactoperoxidase from raw whole milk by cation exchange chromatography. Separation Purification Technol., 48, 143-149.

1174461ja

Fweja, L.W.T., M.J. Lewis and A.S. Grandison, 2008. Challenge testing the lactoperoxidase system against a range of bacteria using different activation agents. J. Dairy Sci., 91: 2566-2574.

433822ja

Hayashi, M., S. Naknukool, S. Hayakawa, M. Ogawa and A.B.A. Ni'matulah, 2012. Enhancement of antimicrobial activity of a lactoperoxidase system by carrot extract and β -carotene. Food Chem., 130: 541-546.

1174469ja

Seifu, E., E.M. Buys and E.F. Donkin, 2004. Quality aspects of Gouda cheese made from goat milk preserved by the lactoperoxidase system. Int. Dairy J., 14: 581-589.

1181348ja

Seifu, E., E.M. Buys and E.F. Donkin, 2005. Significance of the lactoperoxidase system in the dairy industry and its potential applications: A review. Trends Food Sci. Technol., 16: 137-154.

571875ja

Singh, A.K., N. Singh, S. Sharma, K. Shin and M. Takase et al., 2009. Inhibition of lactoperoxidase by its own catalytic product: Crystal structure of the hypothiocyanate-inhibited bovine lactoperoxidase at 2.3-A resolution. Biophys. J., 96: 646-654.

568418ja

Touch, V., S. Hayakawa, S. Yamada and S. Kaneko, 2004. Effects of a lactoperoxidase-thiocyanate-hydrogen peroxide system on *Salmonella enteritidis* in animal or vegetable foods. Int. J. Food Microbiol., 93: 175-183.

78030ja

Zhou, Y. and L.T. Lim, 2009. Activation of lactoperoxidase system in milk by glucose oxidase immobilized in electrospun polylactide microfibers. J. Food Sci., 74: C170-C176.

568445ja

Shahzaib 28-8-2013, {Add by saira: 28-08-13}

Hypered by: Hina 28-8-13

Table 1. LPO activities of solution obtained from dilution of 1 M of NaCl in PB pH 7.0 through captured LPO from bovine whey onto SP Sepharose Fast Flow in three groups of fraction number. Data were obtained from three replications.

Commented [O20]: Delete table 1 and explain its data in text, as it does not contain significant amount of data that needs to be presented in tabular form.

Figure 1. Immobilization efficiency (IE) of LPO on two types of resin, i.e. SP Sepharose Big Beads (SPBB) and SP Sepharose Fast Flow (SPFF). Two types of resin (0.1–1.0 g) was used to immobilized 1 ml of LPO (35 U/ml). Flow rate has been set to 1 ml/min using peristaltic pump. Values are means of three sets of experiments. Error bars represent standard errors of the mean.

Commented [ANA21]: Table 1 has been deleted and the content has been explained in the text.

Commented [O22]: Define all abbreviations used in figures in their respective captions.

Commented [O23]: The given captions of your figures 1 and 2 have not been written properly. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such captions for all these figures in which all of their main parts have been properly explained.

Formatted: Indent: Left: 0", First line: 0"

Commented [O24]: Define all abbreviations used in figures in their respective captions.

Commented [O25]: The given captions of your figures 1 and 2 have not been written properly. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such captions for all these figures in which all of their main parts have been properly explained.

Formatted: Indent: Left: 0", First line: 0"

Figure 1. Immobilization efficiency of SPBB and SPFF using various weight.

Figure 2a. The percentage of remaining LPO activity attached onto SP Sepharose Big Beads (SPBB) during 10 days of storage at 10°C in various storage solutions. This experiment used 1 g of SPBB for immobilizing 1 ml of LPO (35 U/ml). Immediately after immobilization, SPBB was stored in sterilized pure water, phosphate buffer, milk, and whey for 10 days at 10°C. Values are mean of three sets of experiments. Error bars represent standard errors of the mean.

Commented [FT26]: Assign an alphabet to each sub-part of Figure 2 and explain each figure part in figure caption by referring to its respective alphabet. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such caption in which all main parts of this figure have been properly explained. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Commented [O27]: In submitted manuscript Figure 2 has been presented but Author didn't cite this figure in text. Cite this figure in text at appropriate place with suitable explanation and make sure that all figures have been cited in text in consecutive numerical order. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Figure 2b The percentage of remaining LPO activity attached onto SP Sepharose Fast Flow (SPFF) during 10 days of storage at 10°C in various storage solution. This

Commented [ANA28]: Self explanatory has been added

Commented [ANA29]: This figure has been presented in the article.

experiment used 1 g of SPFF for immobilizing 1 ml of LPO (35 U/ml). After immobilization, SPFF was stored in sterilized pure water, phosphate buffer, milk, and whey. Values are mean of three sets of experiments. Error bars represent standard errors of the mean

Figure 2a- The percentage of LPO activity attached onto SPBB during ten days of storage in various storage solutions

Figure 2b The percentage of LPO activity attached onto SPFF during ten days of storage in various storage solution.

Commented [FT30]: Assign an alphabet to each sub-part of Figure 2 and explain each figure part in figure caption by referring to its respective alphabet. A figure should be self-explanatory and its caption plays a very important role in this regard. Provide such caption in which all main parts of this figure have been properly explained. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Commented [O31]: In submitted manuscript Figure 2 has been presented but Author didn't cite this figure in text. Cite this figure in text at appropriate place with suitable explanation and make sure that all figures have been cited in text in consecutive numerical order. All cited figures must be presented in the manuscript and all presented figures must be cited in the text at appropriate places along with their suitable explanation.

Commented [ANA32]: Self explanatory has been added

Commented [ANA33]: This figure has been presented in the article.

RESISTANCE OF IMMOBILIZED LACTOPEROXIDASE ACTIVITY FROM BOVINE WHEY AGAINST STORAGE SOLUTIONS

ORIGINALITY REPORT

13%	%	13%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

3%
★ Amir Golmohamadi, Matthew J. Morra, Inna Popova, Caleb I. Nindo. "Optimizing the use of Sinapis alba seed meal extracts as a source of thiocyanate (SCN –) for the lactoperoxidase system", LWT - Food Science and Technology, 2016
Publication

Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	Off		