

# The Browning Value Changes

*by Ahmad Nimatullah Al-baarri*

---

**Submission date:** 14-Nov-2019 11:26AM (UTC+0700)

**Submission ID:** 1213460568

**File name:** ctral\_Analysis\_on\_the\_Maillard\_Reaction\_Product\_From\_Glucose.pdf (590.22K)

**Word count:** 1693

**Character count:** 9313

1

PAPER • OPEN ACCESS

2

## The browning value changes and spectral analysis on the Maillard reaction product from glucose and methionine model system

4

To cite this article: A.N. Al-Baari *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **102** 012003

View the [article online](#) for updates and enhancements.

2

## The browning value changes and spectral analysis on the Maillard reaction product from glucose and methionine model system

A.N. Al Baarri<sup>1\*</sup>, A.M. Legowo<sup>1</sup>, Widayat<sup>2</sup>

<sup>1</sup>Food Technology Department, Faculty of Animal and Agricultural Sciences, Diponegoro University, Indonesia

<sup>2</sup>Chemical Engineering Department, Faculty of Engineering, Diponegoro University, Indonesia

\*email: [albari@undip.ac.id](mailto:albari@undip.ac.id)

**Abstract.** D-glucose has been understood to provide the various effect on the reactivity in Maillard reaction resulting in the changes in physical performance of food product. Therefore this research was done to analyse physical appearance of Maillard reaction product made of D-glucose and methionine as a model system. The changes in browning value and spectral analysis model system were determined. The glucose-methionine model system was produced through the heating treatment at 50°C and RH 70% for 24 hours. The data were collected for every three hour using spectrophotometer. As result, browning value was elevated with the increase of heating time and remarkably high if compare to the D-glucose only. Furthermore, the spectral analysis showed that methionine turned the pattern of peak appearance. As conclusion, methionine raised the browning value and changed the pattern of spectral analysis in Maillard reaction model system.

**Keywords:** D-glucose, methionine, browning value, spectral analysis, Maillard reaction

### 1. Introduction

Maillard reaction has been widely known as a three step of chemical reaction involving both carbonyl groups of reducing sugars and free amino groups of proteins that contributes to a very important role in food industry that applies thermal process. These three steps are early, intermediate and final step [1]. Maillard reaction has been recognized as critical effects on food properties, such as colour, texture, and flavour that might provide attractive odour such as caramel-like and burning [2]. The generation of these senses are rely on the applied amino acid and sugar. The reactivity of Maillard reaction was reported as sugar variety dependent [3, 4] and applied of protein [5].

Besides their generation in specific odour, Maillard reaction provides beneficial functions as antioxidant [6]. The utilization of glucose might generate the production of 3-deoxy-glucosone, as one of the intermediate products of the Maillard reaction. This compound acts as a precursor of 5-hydroxymethylfurfural generation, resulting in the high antioxidant activity and brown colour of Maillard reaction product [2].



1 Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

Since glucose has been widely known as ingredient in Maillard reaction, this study aimed to determine the browning value and spectral analysis in Maillard reaction between D-glucose and methionine.

## 2. Material and method

D-glucose was obtained from Sigma Aldrich US. Methionine was purchased from Cheil Jedang Indonesia, Co. Ltd with the purity index of 99%. Unless otherwise specified, all other chemicals were reagent grade.

### 2.1 Preparation of D-glucose-methionine model system

The preparation method of D-glucose-methionine model system has been adopted from other researcher with some modifications [7]. The D-glucose as a control and 1:1 D-glucose-methionine were dissolved in 400 mL of 10 mM carbonate buffer (pH 9) solution. The solution was then heated at 50°C and RH 70% for 24 hours.

### 2.2 Spectral analysis of Maillard reaction product

The heated samples were diluted with phosphate buffer until 200  $\mu$ L in microtube to measure the emission spectrum starting from 190 to 620 nm using spectrophotometer (UV-1280; Shimadzu, Kyoto, Japan). This spectroscopic measurement method has been adopted from previous researcher [7].

### 2.3 Browning value measurement

The browning value of the MRPs was measured according to the method of earlier researcher [7] without any modification using spectrophotometer at 420 nm.


### 2.4 Data analysis

The results were reported in figures and the description upon figure was provided. The data were obtained from at least three time of replication.


## 3. Result and discussion

Figure 1 shows the browning value through the determination of absorbance at 420 nm <sup>2</sup> on the Maillard reaction product using D-glucose and methionine model system when was heated at 50°C for 24 hour. It can be described that the increase of both sample was able to be detected clearly. D-glucose-methionine model system showed remarkably higher (about two times higher) in the absorbance value than D-glucose only. As seen on this figure, the Maillard reaction model system generating absorbance at 1.218 while sugar alone only generated the absorbance at 0.644. The higher browning value might able to be recognized starting at heating time of 9 hour that was resulting the value of 0.660 while sugar-only-sample provided about a half of absorbance (0.338).

In this research, methionine was chosen to generate the Maillard reaction in model system, due to its high reactivity [5]. D-glucose was applied as the control because it widely used in Maillard reaction process. Hexoses-amino acid system model generated its specific reactivity based on the sugar and amino acid that was used [8]. Methionine was known as the high reactivity of amino acid and it reduced at about 60% when Maillard processing was applied [9]. Spectral analysis showed similar patterns or peak appearance on the D-glucose model system when the heating time of 15 and 18 hour was applied (Fig. 2a). However, the remarkable differences in the pattern or peak appearance was occurred when the methionine was applied (Fig. 2b). This could be seen at the appearance of peak at 18-h heating time. As can be seen, no peak was detected at the D-glucose-methionine model system on 18-h heating time. Therefore, the presence of amino acid in the model system turned the spectral appearance remarkably.



**Figure 1.** Browning development through the determination of absorbance at 420 nm on Maillard reaction product of D-glucose and D-glucose-methionine model system that was heated at 50°C for 24 hour.



**Figure 2.** Spectral analysis of Maillard reaction product of the D-glucose (a) and D-glucose-methionine model system (b) that was heated at 50°C.

It was recognized that the heating process might reduce remarkably the luminosity ( $L^*$ ) in Maillard reaction system [10]. This reduction in the  $L^*$  value was due to the formation of brown pigments from Maillard reaction that might interference the spectral analysis. Brown pigments was easily detected at 420 nm [11], therefore based on the obtained spectral analysis at 420 nm, the change on the absorbance at those wavelength might be detected with ease.

#### 4. Conclusion

As conclusion, the browning value of D-glucose-methionine model system were higher than D-glucose model system and it could be seen starting at the 9-h of heating time. The heating time at 18 h

showed the change in peak appearance and pattern of spectral analysis. This result may provide the beneficial information about Maillard reaction process during 24 hour.

### 5. Reference

- [1]. Jiang W, He X, Yang H, Xiang X, Hu S, Li S and Liu Y 2017 Histamine reduction by Maillard reaction with glucose, *Food Control* **82**(Supplement C) 136-144.
- [2]. Asikin Y, Kamiya A, Mizu M, Takara K, Tamaki H and Wada K 2014 Changes in the physicochemical characteristics, including flavour components and Maillard reaction products, of non-centrifugal cane brown sugar during storage, *Food Chemistry* **149** 170–177.
- [3]. Sun Y, Hayakawa S, Puangmanee and K. Izumori 2006 Chemical properties and antioxidative activity of glycated  $\alpha$ -lactalbumin with a rare sugar, D-allose, by maillard reaction, *Food Chemistry* **95** 509-517.
- [4]. Sun Y, Hayakawa S, Chuamanochan M , Fujimoto M, Innun A and Izumori K 2006 Antioxidant effect of maillard reaction products obtained from ovalbumin and different D-Aldohexoses, *Bioscience, Biotechnology, and Biochemistry* **70**(3) 598-605.
- [5]. Suminar AT, Al-Baari AN and Legowo AM 2017 Demonstration of physical phenomena and scavenging activity from d-psicose and methionine maillard reaction products, *Potravinarstvo Slovák Journal of Food Sciences* **11**(1) 417–424.
- [6]. Chen X.-M and Kitts D D 2011 Antioxidant and anti-inflammatory activities of Maillard reaction products isolated from sugar–amino acid model systems *Journal of Agricultural and Food Chemistry* **59** 11294–11303.
- [7]. Arum SK, Legowo AM, Al-Baari AN, Hayakawa S and Ogawa M 2017 Prolongation the Shelf Life of Indonesian Soft Cheese (Dangke) Derived from Bovine Milk using Lactoperoxidase System and Lysozyme, *Czech Journal of Food Science Article* in Press.
- [8]. Tan T.-C, AlKarkhi AFM and Easa AM 2012 Assessment of the ribose-induced Maillard reaction as a means of gelatine powder identification and quality control, *Food Chemistry* **134** 2430-2436
- [9]. Hou C, Wu S, Xia Y, Phillips G O and Cui SW 2017 A novel emulsifier prepared from Acacia seyal polysaccharide through Maillard reaction with casein peptides, *Food Hydrocolloids* **69** 236–241.
- [10].Cai L, Li D, Dong Z, Cao A, Lin and Li J. 2016 Change regularity of the characteristics of Maillard reaction products derived from xylose and Chinese shrimp waste hydrolysates, *LWT - Food Science and Technology* **65** 908–916.
- [11].Delgado-Andrade C, Seiquer I, Nieto R and Navarro MP 2004 Effects of heated glucose–lysine and glucose–methionine model-systems on mineral solubility, *Food Chemistry* **87** 329–337.

# The Browning Value Changes

ORIGINALITY REPORT

<b>19%</b>	%	<b>19%</b>	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

- 1** A N Al-Baarri, A M Legowo, Widayat, S B M Abduh, M Hadipernata, Wisnubroto, D K Ardianti, M N Susanto, M Yusuf, E K Demasta. "Determination Hypoiodous Acid (HIO) By Peroxidase System Using Peroxidase Enzyme", IOP Conference Series: Earth and Environmental Science, 2018  
Publication **7%**
- 2** Luiz Gustavo Gonçalves Rodrigues, Simone Mazzutti, Luciano Vitali, Gustavo A. Micke, Sandra Regina Salvador Ferreira. "Recovery of bioactive phenolic compounds from papaya seeds agroindustrial residue using subcritical water extraction", Biocatalysis and Agricultural Biotechnology, 2019  
Publication **5%**
- 3** J. A. Marengo, C. A. Nobre, G. Sampaio, L. F. Salazar, L. S. Borma. "Chapter 9 Climate change in the Amazon Basin: Tipping points, changes in extremes, and impacts on natural and human systems", Springer Science and **3%**

4

A N Al-Baarri, A M Legowo, Widayat, S B M  
Abduh, F P Lestari, D Desnasari, I P M Santoso.  
"Potential of L-fucose isolated from Brown  
Seaweeds as Promising Natural Emulsifier  
compare to Carboxymethyl Cellulose (CMC)",  
IOP Conference Series: Earth and  
Environmental Science, 2018

2%

Publication

---

5

Luyun Cai, Dongmei Li, Zhijian Dong, Ailing  
Cao, Hong Lin, Jianrong Li. "Change regularity  
of the characteristics of Maillard reaction  
products derived from xylose and Chinese  
shrimp waste hydrolysates", LWT - Food  
Science and Technology, 2016

2%

Publication

---

Exclude quotes

Off

Exclude matches

< 20 words

Exclude bibliography

On