

Effect of Brusato/Biolarvicide Administration on Behavioral Response of *Aedes aegypti* and its Toxicity on Vero Cells

by Dwi Sutiningsih


Submission date: 10-Jun-2019 08:42AM (UTC+0700)

Submission ID: 1141848225

File name: Effect_of_Brusato/Biolarvicide.pdf (761.5K)

Word count: 6257

Character count: 32686



Journal of Biological Sciences

ISSN 1727-3048

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>



Research Article

Effect of Brusatol Biolarvicide Administration on Behavioral Response of *Aedes aegypti* and its Toxicity on Vero Cells¹Dwi Sutningsih and ²Nurjazuli¹Department of Epidemiology and Tropical Disease, Faculty of Public Health, University of Diponegoro, Semarang, Indonesia²Department of Environmental Health, Faculty of Public Health, University of Diponegoro, Semarang, Indonesia

Abstract

Background and Objective: Brusatol is one of quassinoid, derived from Makassar fruit (*Brucea javanica* (L.) Merr.) and belong to the family of Simaroubaceae. Brusatol has been proven to have insecticidal and anti-feeding activity on larvae of *Spodoptera exigua*. This study aimed to investigate the larvicidal activity and behavioral response to the larvae of *A. aegypti* after administration of brusatol and to demonstrate cytotoxicity activity in vero cells. **Materials and Methods:** Larvicidal activity was monitored using bioassay test. The behavioral response of larval *A. aegypti* was observed after treatment with various brusatol concentrations and compared directly with control every 10, 30 min and 1, 5 and 24 h. The examination of brusatol cytotoxicity on vero cells performed by MTT (3-[4,5-dimethylthiazol-2-yl]-2,5 diphenyltetrazolium bromide) method. **Results:** The results of this study showed that the behavioral response of *A. aegypti* larvae after treatment with brusatol has similarity with the behavioral response after treatment with temephos 1 ppm. The behavioral response of *A. aegypti* larvae showed symptoms of a nerve poison that is excitation, convulsions, tremors and paralysis. The lethal concentration of 50 and 90 (LC₅₀, LC₉₀) to *A. aegypti* larvae were 0.669±0.106 and 8.331±0.060 ppm, respectively. Brusatol cytotoxic activity on vero cells is low, with IC₅₀ values worth 1402.029±0.164 mg mL⁻¹. **Conclusion:** The results of this study proved the potential of brusatol as biolarvicide to control the mosquito vector with low levels of cytotoxicity.

Key words: Brusatol, *Aedes aegypti*, vero cell, behaviour, cytotoxicity

⁵Received: February 02, 2017

Accepted: March 08, 2017

Published: March 15, 2017

Citation: Dwi Sutningsih and Nurjazuli, 2017. Effect of brusatol biolarvicide administration on behavioral response of *Aedes aegypti* and its toxicity on vero cells. J. Biol. Sci., 17: 127-135.

Corresponding Author: ⁹Dwi Sutningsih, Department of Epidemiology and Tropical Disease, Faculty of Public Health, University of Diponegoro, Semarang, Indonesia Tel/Fax: 024-7460044

Copyright: © 2017 Dwi Sutningsih and Nurjazuli. This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Dengue Hemorrhagic Fever (DHF) until today is one of the diseases which becomes a serious public health problem in Indonesia. This disease is transmitted by the *Aedes aegypti* mosquito¹. One method to limit the spread of dengue is by using larvicides. In 1980, temephos 1% (abate) established as part of a mass eradication program to eradicate *Aedes aegypti* in Indonesia. But using temephos repeatedly may lead to the emergence of resistance of various species of mosquitoes which are also known as disease's vectors². Development of new insecticides which do not pose a risk of resistance and environmentally more friendly can be obtained through the use of biolarvicide³. Bioinsecticide or biological insecticide is an insecticide, its base ingredients derived from plants that contain chemical substances (bioactive) which is toxic to insects and easy to decompose (biodegradable) in nature so it does not pollute the environment and relatively are safe for humans. In addition, biological insecticide is selective⁴. The toxicity power comes from its toxic substances that contained in biological insecticide. These substances can act as toxins through the digestive tract absorption or through the skin on the animals which have tender body⁵.

Brusatol is a quassinoid derived from *B. javanica* (L.) Merr. fruit. Its molecular formula is $C_{26}H_{32}O_{11}$ with amorphous powder and bitter taste. Zhang *et al.*⁶ have proved that brusatol isolated from *B. javanica* (L.) Merr. has insecticidal and antifeedant properties against instar 3 larvae of *Spodoptera exigua*. Syahputra⁷ and Lina *et al.*⁸ have also demonstrated that the active material contained in *B. javanica* (L.) Merr. extract has larvicidal effect against *Crocidolomia pavonana* (*Lepidoptera: Crambidae*). Extract of *B. javanica* (L.) Merr. can inhibit feeding, decrease the rate of growth and inhibit laying *C. pavonana* imago⁹. These informations showed that the study of *B. javanica* (L.) Merr. and brusatol compound as larvicides/insecticides is limited to the use of agricultural pests. Therefore, this study aimed to investigate the more detail on the brusatol larvicidal activity and the effect on behavior of the larvae of *A. aegypti* after treatment with brusatol concentration and cytotoxicity in normal cells (vero cells).

MATERIALS AND METHODS

Makasar fruit (*B. javanica* (L.) Merr.) was purchased from a wholesaler of medicinal plants (Herbal Merapi Yogyakarta) and further identified in the Laboratory of Pharmaceutical Biology of the Faculty of Pharmacy Gadjah Mada University to obtain confirmation of the simplisia data information. The instar III larvae of *A. aegypti* until the beginning of instar IV

was acquired and developed at the Laboratory of Parasitology, Faculty of Medicine, Gadjah Mada University. Vero cells were obtained from the Laboratory of Parasitology, Faculty of Medicine, University of Gadjah Mada. The isolation and identification of brusatol were performed in the Laboratory of Biology Pharmacy, Faculty of Pharmacy, Gadjah Mada University. The behavior testing of the larvae of *A. aegypti* and the brusatol cytotoxicity on vero cells were undertaken at the Laboratory of Parasitology, Faculty of Medicine, Gadjah Mada University in Yogyakarta.

Extraction and isolation of brusatol: The isolation of brusatol from the seeds of *B. javanica* (L.) Merr. was undertaken using the method of Zhang *et al.*⁶ with a slight modification; *Brucea javanica* (L.) Merr. dried seeds (5 kg) made it into powder, shaken with EtOH-H₂O (15 L), the solvent was evaporated in vacuum and the extracts were combined and concentrated, followed by suspending in H₂O. The aqueous layer was further extracted with petroleum ether, CHCl₃, EtOAc and n-Butanol. The CHCl₃ layer was evaporated under vacuum to afford extracts. This extract was re-suspended with CHCl₃ and then chromatographed on a silica gel column (400 g, 200-300 mesh) by eluting successively with CHCl₃ containing increasing amounts of MeOH (1:0, 50:1, 20:1, 10:1, 5:1 and 0:1). The result in the organic layer is taken and evaporated to obtain a concentrate and then made a solution of MeOH (100-250 mL) at a temperature of 60°C and then stored at room temperature. MeOH solution above when it is allowed to stand at room temperature to form crystalline compounds that expected as brusatol. Further separation is by filtration. The rest of the filtrate/residue is separated by Thin Layer of Chromatography (TLC). The filtrate/residue is evaporated. Further separation is performed if it is necessary.

Larvicidal test: Larvicidal activity test was conducted using bioassay method according to the standard of WHO¹⁰ with a slight modification. Initially it was performed a preliminary test to determine the range of brusatol concentration that can be deadly to larvae of *A. aegypti* on instar III until the beginning of instar IV. Furthermore, the concentrations used in further tests with positive control is temephos 1 ppm whereas the negative control is used only as much as 100 mL of distilled water, repeated 4 times. The larvae of *A. aegypti* on the end of instar III until the beginning of instar IV are used as much as 25 larvae put in each treatment media and control. After 24 h, the larvae of *A. aegypti* that were dead were counted. Temperature, pH of the media and humidity in the room were measured at the beginning and the end of the study.

Behavioral change test: Behavioral change was measured using the method described by Sharma *et al.*¹¹ with a slight modification. The instar III larvae of *A. aegypti* until the beginning of instar IV were put in a plastic cup filled with brusatol with various concentrations (0.33, 0.6, 1, 3, 9 and 27 ppm), each of 25 larvae, which consisted of four replications with positive control is temephos 1% while the negative control is used only as much as 100 mL of distilled water. Change in larval behavior was monitored every 10 min, 30, 1, 5 and 24 h after treatment. Data on larval behavior was observed by speed/activity of movement, physical response, posture and behavior. Data was recorded and photographed using Nikon D5300 camera. Similar observations were performed in the control group and then compared to the treatment group.

Brusatol cytotoxicity test: Brusatol cytotoxicity test in vero cells was conducted using MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) based on the method of Van Meerloo *et al.*¹² with a slight modification. Vero cells were cultured using M199 medium which has added 10% FBS, 2% penicillin-streptomycin and 0.5-1% fungison. Vero cells were taken from liquid nitrogen, heated at 37°C until liquefied. The cell suspension that has liquefied put into conical tube and washed with M199 complete medium. The cell suspension was then transferred into a culture flask and incubated in a 37°C incubator, 5% CO₂. Cell growth was observed with inverted microscope every day until the cells almost meet the base wall of flask. Vero cell harvesting is done after the flask was almost completely full. Cells were washed with PBS and added with 0.25% trypsin for the cells detached from the flask wall. The cell suspension was made by adding complete medium, after which the number of cells were counted with a haemocytometer. Cell is inserted into a 96 well microplate at a density of 2×10^4 cells/well in 100 μ L. Then the culture medium containing brusatol for each variation of the concentration (1200, 600, 300, 150, 75, 37.5 μ g mL⁻¹) was added. Cell cultures were incubated for 24 h in an incubator temperature of 37°C, 5% CO₂. Cell growth was observed using MTT then compared with the control culture (without treatment). Medium discarded after incubation mass ends then added 100 μ L of complete medium and 10 μ L solution of MTT. Furthermore it was incubated for 4 h in an incubator at 37°C, 5% CO₂. Then it was added 100 μ L of 10% SDS in 0.01 M HCl to dissolve formazan which was formed and incubated overnight at room temperature. The test result was read by ELISA at a wavelength of 595 nm.

Data analysis: Probit analysis was conducted on larval mortality after 24 h exposure to different concentrations of brusatol using SPSS version 24. If the data were normally distributed and homogeneous then the data were analyzed using one-way ANOVA test, whereas if the data were not normally distributed and not homogeneous the data were tested using the Kruskal-Wallis test, with a reliability level of 0.05 to determine whether there are differences between the mortality of *A. aegypti* larvae in various concentration of treatment.

The data of viable cell obtained is calculated inhibition percentage using the formula in Eq. 1³²:

$$\frac{AB}{CD} \times 100\% \quad (1)$$

where, A is absorbance of treatment (cells+media +compound), B is absorbance of control samples (media+compound), C is absorbance of control cells (cells+media) and D is absorbance media control.

The value of IC₅₀ was calculated using probit regression analysis using SPSS version 24. The small value of IC₅₀ indicates that the compound has a high cytotoxic effect and vice versa the big value of IC₅₀ suggests that the cytotoxic effect is low.

RESULTS

Larvicidal activity and behavior of *A. aegypti* after treatment with brusatol: The environment considered in this study was the pH of the media, media temperature and humidity. The results of the measurement of media temperature, pH and humidity in the room of the study respectively was 25°C, 7 and 71-72%. The percentage of the mortality of *A. aegypti* larvae after observation for 24 h was presented in Table 1. The lethal concentration of 50 and 90 (LC₅₀ and LC₉₀) calculated by probit analysis using SPSS version 24 were 0.669 ± 0.106 and 8.331 ± 0.060 ppm respectively. Kruskal Wallis test results showed the value of $p = 0.000$ ($p < 0.05$), the average number of deaths of *A. aegypti* larvae was significantly different.

Observation of behavioral response of *A. aegypti* larvae after treatment with brusatol was conducted directly by looking at changes in the physical and behavioral response or body movements of *A. aegypti* larvae as compared with controls. The result of behavioral response observation on the larvae of *A. aegypti* is shown in Table 2. Observation of behavioral responses to larvae of instar III *A. aegypti* and early instar IV after brusatol treatment showed a different pattern of behavior compared to the control. Immediately after brusatol

Table 1: Average of percentage of the mortality of *A. aegypti* larvae in the group treated with brusatol in wide range of concentrations, temephos of 1 ppm and control after 24 h of observation

Brusatol concentration (ppm)	No. of <i>A. aegypti</i> larvae on each test	Mortality of <i>A. aegypti</i> larvae on each test					
		I	II	III	IV	Mean	Mortality \pm SD (%)
0.33	25	12	8	10	10	10.00	40 \pm 1.63*
0.6	25	12	10	10	11	10.75	43 \pm 0.96*
1	25	13	14	12	13	13.00	52 \pm 0.82*
3	25	22	22	23	20	21.75	87 \pm 1.26*
9	25	20	21	23	19	20.75	83 \pm 1.71*
27	25	25	25	25	25	25.00	100 \pm 0.00*
Temephos 1 ppm	25	25	25	25	25	25.00	100 \pm 0.00
Control	25	0	0	0	0	0.00	0 \pm 0.00

*p<0.05 with Kruskal Wallis test, ppm: Part per million, SD: Standard deviation

exposure, all the larvae of *A. aegypti* indicated a natural and powerful movement (natural and vigorous movement). But after 10 min of brusatol exposure, larvae of *A. aegypti* indicated anxiety, excitation and agitation as well as on the provision of temephos 1 ppm. Then the larvae would quickly writhe around 10-15 min and began to sink in the bottom of the water. Most larvae showed heavy anxiety response and very aggressive by biting their gills anal with its mouth so that it looked like a circular motion to form a ring structure (Fig. 1).

Brusatol cytotoxicity on vero cells: The average of alive vero cells observation and the percentage of viable cells after brusatol administration and the control can be seen in Table 3. The percentage of vero cells which were alive reaching 100 \pm 0.014% at the lowest concentration of brusatol which is 37.5 $\mu\text{g mL}^{-1}$, whereas at the highest concentration of brusatol which is 1200 $\mu\text{g mL}^{-1}$, the percentage of viable cells is 49.897 \pm 0.217%. Results of probit analysis using SPSS program version 24 obtained value of IC₅₀ of brusatol on vero cells at 1402.029 \pm 0.164 $\mu\text{g mL}^{-1}$. Kruskal Wallis test results showed the value of p = 0.153 (p>0.05), the average of vero viable cell was not significantly different.

DISCUSSION

Larvicidal activity and behavior of *A. aegypti* (L.) after treatment with brusatol: The environment that is considered in this study is the pH of the media, media temperature and humidity. The levels of pH, media temperature and humidity measured at the beginning and the end of the study. The result of media temperature measurement shows that the temperature of the media is still in the optimal temperature range (25-27°C) for the development of *A. aegypti* that is

25°C. The pH value of the media was also in the optimal pH range (6.5-7) for larval development *A. aegypti* that is 7. The results of the measurement of humidity in the room of the study ranged between 70-71%, that is still within the range of normal humidity for larval development *A. aegypti* that is 60-80%¹⁰. From these results it can be concluded that the mortality of larvae *A. aegypti* is not caused by environmental factors such as pH media, media temperature and humidity.

The results of 24 h observation showed that there is no larval mortality *A. aegypti* on control group, so that the number of dead larvae of *A. aegypti* in the treatment group did not need to be corrected by Abbot formula. At the lowest concentrations of brusatol that is 0.33 ppm larval mortality of *A. aegypti* is occurred by 40 \pm 1.63%. Larval mortality of *A. aegypti* reached 100% at a brusatol concentration of 27 ppm which is equivalent to giving temephos at a concentration of 1 ppm (Table 1). This means that brusatol is effective as a natural larvicide on the larvae of *A. aegypti*. The higher the concentration of brusatol, the higher the mortality of larvae of *A. aegypti*. The mortality of *A. aegypti* in this study is due to the toxic activity from the brusatol compound. Toxicity of brusatol on *A. aegypti* larvae stated with the lethal concentration of 50 and 90 (LC₅₀ and LC₉₀) which amounted to 0.669 \pm 0.106 and 8.331 \pm 0.060 ppm. The smaller the number of lethal concentration the compound is more toxic. Kruskal Wallis test results showed that the average number of deaths of *A. aegypti* larvae is significantly different. The difference in larval mortality of *A. aegypti* larvae is likely to be influenced by several factors, namely instar larvae and differences in the sensitivity of each test larvae. Factor larval instar is related to age of the larvae. Although the larvae used were instar III larvae until the early of instar IV, the varieties of larval age is remained. Early instar III larvae are more susceptible to insecticides than the early instar IV larvae¹³. In addition, differences in larval mortality is influenced also by the

Table 2: Behavior of *A. aegypti* larvae after administration of wide range of brusatol concentrations, 1 ppm of temephos and control over the time period of observation

Time (min)	Brusatol concentration (ppm)	Behaviour	Temephos (1 ppm)	Control
0 (min)	0.33	All larvae showed agile and strong mobility, active movement up and down	All larvae showed agile and strong mobility, active movement up and down	
10 (min)	0.6	Shows their anxiety, excitation, agitation/upheaval	Anxiety, tremor	
30 (h)	1	Anxiety is strong, circular movements and bite anal papillae aggressively and form a ring	Anxiety and getting stronger tremor	All larvae showed agile and strong mobility, active movement up and down
1 (h)	3	Anxiety is strong and fast, tremors, convulsions, almost die and sink	Tremors, convulsions, paralysis, moribund and sinking	
5 (h)	9	Strong Tremor and convulsion, followed by paralysis, most moribund and sink	All larvae die sinking, parallel to the surface of the water	
24 (h)	27	Larvae die sinking and parallel to the surface of the water, smaller body size	All larvae die sinking, smaller body size	

ppm.:Part per million




Fig. 1(a-b): Digital photomicrographs of early IV instar larvae of *A. aegypti*, (a) Control larvae and (b) Larvae treated with brusatol 9 ppm
 Circled larvae show the aggressive anal gill biting behaviour forming ring-shaped structure

differences in the sensitivity of each test larvae. Although using larvae with the same instar and age, each larva has a different level of vulnerability¹⁴.

Brusatol is a toxic substances contained in Makasar fruit (*B. javanica* (L.) Merr.) which gets into the body of *A. aegypti* larvae through the skin, mouth and respiratory tract. From Table 2 it can be seen that difference in the behavioral response of *A. aegypti* larvae between the normal larvae (control) and larvae which got treatment with brusatol and 1 ppm of temephos. About 10 min observation after giving brusatol, *A. aegypti* larvae showed movement agitation, excitation and restlessness, while by giving 1 ppm temephos showed their anxiety and tremor. Meanwhile, normal larvae (control) showed agile and strong mobility. Thirty minutes later looked aggressive and circular movements by biting annals gills and formed a ring on *A. aegypti* larvae after

Table 3: Average Vero cells were alive and the percentage of viable cells after brusatol administration and control

Brusatol concentration ($\mu\text{g mL}^{-1}$)	Absorbance value on each test				Mean	Viable cell \pm SD (%)
	I	II	III	IV		
1200	0.201	0.162	0.248	0.496	0.277	49.897 \pm 0.217
600	0.469	0.233	0.453	0.513	0.417	77.789 \pm 0.181
300	0.473	0.547	0.302	0.496	0.454	85.158 \pm 0.153
150	0.440	0.583	0.545	0.302	0.467	87.741 \pm 0.181*
75	0.517	0.058	0.538	0.422	0.384	97.176 \pm 0.098*
37.5	0.513	0.517	0.829	0.736	0.649	100 \pm 0.045
Cell Control	0.715	0.761	0.767	0.805	0.762	100 \pm 0.037
Media Control	0.044	0.033	0.032	0.035	0.036	0 \pm 0.005

* $p < 0.05$ with Kruskal Wallis test, SD: Standard deviation

giving brusatol. The results are consistent with study of Sharma *et al.*¹¹ who proved that early instar IV of *A. aegypti* larvae shows behavioral agitation, excitation, restlessness and aggressive movement by biting anal gills after exposure of leaf extract of *Achyranthes aspera* and stem extracts of *A. aspera*. These symptoms are similar to symptoms caused by a neurotoxin, meaning that brusatol might act as sitolisin that affect neuromuscular larvae system. Circular movement of the larvae, accompanied by aggressive biting anal gills behavior proved that there are neurotoxic symptoms of brusatol on *A. aegypti* larvae (Fig. 1b). Anal gills on mosquitoes have some role in controlling electrolyte balance needed for survival¹⁵. The effects of brusatol toxic caused electrolyte imbalance in the anal area, causing violent by biting anal gills. The results of this study were similar to results of previous studies which indicate that bioactive plant compounds can act as a neurotoxin although aggressive and un-coordinated movements together with other toxic symptoms were observed at relatively different time intervals¹⁶⁻²⁰. Observation of similar behavior has been reported by Warikoo and Kumar²¹ on the larvae of *A. aegypti* tested with root extract of *Argemone mexicana*. The result of microscopic observation on the early instar IV of *A. aegypti* larvae after treatment by giving concentration of *A. mexicana* extract 91.331 and 156.684 ppm for 24 h showed a structural deformity of anal gills. This damage may cause the function anomalies that may have led to interference in the regulation of the osmotic and ionic²⁰. Malfunctioning of this regulation may lead to the death of the larvae of *A. aegypti*. One hour after treatment with brusatol, larvae of *A. aegypti* showed the movement of restlessness, tremors, convulsions and paralysis that increasingly powerful that eventually most larvae die, it also occurred in the larvae of *A. aegypti* after the administration of 1 ppm of temephos. This means that there was a similarity to the behavior of larvae of *A. aegypti* after administration brusatol with 1 ppm of temephos. Sanjaya and Safaria²² mentioned that the toxic compounds that enter the body can cause a four-stage larvae responses which are excitation,

convulsions (spasticity), paralysis (paralysis) and mortality. Brusatol is a toxic substance that can kill larvae of *A. aegypti*, larvae enters the body through the pores of the larvae's skin, gastrointestinal tract and respiratory tract. The toxic compounds caused the disruption of the digestive system, the respiratory system and the nervous system in larvae¹⁸⁻²⁰. These compounds caused sensory nerve cells that are on the surface of the larval body disturbed. The mechanism of action of brusatol is performed by inhibiting acetylcholinesterase enzyme which serves to hydrolyze the acetylcholine. In the normal situation, the acetylcholine functions to transmit nerve impulses, after that, acetylcholine is hydrolyzed by the acetylcholinesterase enzyme into choline and acetic acid²³. The bound acetylcholinesterase enzyme causes the cumulation of acetylcholine, so that results in disruption and damage to the nervous system²⁴. It makes the muscles continue to contract until exhaustion and then leads to paralysis and death.

Brusatol enters the body of larvae through the skin membrane by diffusion. A number of toxic substances that enter the body cause damage to the skin cells. These toxic compounds hydrolyze the skin membrane by remodeling the proteins of the skin (collagen) into several sections. The damage to the skin cell membrane causes the loss of the skin membrane impermeability. Hence, the other toxic compounds freely enter the body of the larvae. The number of toxic compounds entering the body causes damage to the proteins in the skin membrane so that the function of the skin as a protection of the body is disrupted²⁵.

In addition to the diffusion of the skin membrane, brusatol supposedly enters through the gastrointestinal tract. The gastrointestinal tract of the mosquito larvae consists of three parts, which are the front part (foregut), middle (midgut) and back (hindgut) of the gastrointestinal tract. The process of food digestion and absorption occurs in the middle of the gastrointestinal tract/midgut²². In the middle of the gastrointestinal tract/midgut, it is coated by the epithelial tissue. Toxic substances enter through the mouth into the

middle of the gastrointestinal tract/midgut. These toxic substances cause the lysis of the epithelial cells resulting in the reduction of the tension on the surface of the mucous membranes of the middle of gastrointestinal tract/midgut. Thus, the food digestion and absorption might not occur²⁵.

Another way brusatol enter the body of larvae is through the respiratory tract. Air enters through a siphon attached to the surface of the water. This toxic substance is supposed to cover the medium surface, so it blocks the siphon to get oxygen from the medium surface^{22,25}. The nervous tissue of the larvae is very sensitive towards the lack of oxygen, causing the wilting of the nerves and damage of the siphon. Therefore, the larvae face the difficulty in breathing and eventually dying²⁶.

Brusatol as an antifeedant can inhibit the feeding activity of the larvae of *A. aegypti*. According to Zhang *et al.*⁶, brusatol isolated from *B. javanica* (L.) Merr. have an insecticidal and antifeedant effect to the third instar larvae of *Spodoptera exigua*. Further study showed that brusatol can induce apoptosis in insect cell lines IOZCAS-Spec-II and Sf21 characterized by the DNA fragmentation, activation of caspase-3 and the enzyme cytochrome-c release from mitochondria. Antifeedant is a chemical compound that inhibits the feeding activity of insects but will not destruct, expel or entrap the insects directly. Antifeedant compound only inhibits the appetite (feeding inhibition) in insects. Antifeedant compound is suppressant (suppress the activity of biting) and a deterrent (prevent the insects for eating continuously) towards insects²⁷. As another antifeedant compounds, brusatol is considered to be able to destruct larvae indirectly by causing disturbances in the signal transmission system of the appetite stimulant (phagostimulant) on the insects. The low feeding activities of the larvae caused the energy to be reduced so that the larval growth process is inhibited and eventually die. In addition, the bitter taste of brusatol⁶ also caused the larvae lose their appetite which leads to starving and death of the larvae.

Temephos is one of pesticide belongs to class of organic phosphate compounds that can enter the body through the skin, inhaled through breathing and ingested through the mouth. Organic phosphate poisoning in insects followed by restlessness, hyperexcitability, tremors, convulsions and eventually muscle paralysis (paralysis)²⁸. Mortality of *A. aegypti* larvae reached 100% after 24 h of exposure to brusatol concentration of 27 ppm, while the 1 ppm of temephos giving 100% mortality of larvae of *A. aegypti* reached after 5 h of treatment. This means that the time required to turn off

100% of *A. aegypti* larvae is quickly achieved after administration of 1 ppm of temephos than 27 ppm of brusatol. This is because temephos is more quickly absorbed by the larvae of *A. aegypti* compared with brusatol at the same time. Temephos contains phosphorothionate group which has lipophilic nature so it is able to penetrate to the waxy epicuticular layer of *A. aegypti* larvae which is hydrophilic so it is more toxic than brusatol. Temephos toxicity against mosquito larvae is closely related to how much the molecule is oxidized or it can be said that the more oxidized, toxicity decreases. Temephos inhibits the cholinesterase enzyme both in vertebrates and invertebrates, causing disruption in neural activity as a continuous muscle contractions, convulsions and ultimately insects will die²⁹. Metabolism of temephos namely fosforotionat group (P = S) in the body of the animal was changed to phosphate (P = O) is more potential as anticholinesterases. The larvae of *A. aegypti* was able to change the P = S to P = O ester faster than house flies as well as temephos penetration into larvae is rapid where more than 99% of temephos in the medium is absorbed within 1 h after treatment. After the absorption, temephos is converted into metabolic products which some of these metabolic products are excreted into the water^{28,29}.

Brusatol cytotoxicity on vero cells: Cytotoxic test in this study was conducted to determine the brusatol toxicity on normal cells. Cytotoxic test is used to determine the IC₅₀ value parameters. IC₅₀ value indicates the concentration which produces barriers of cell proliferation as much as 50% and demonstrates the potential toxicity of a compound to cell³⁰. IC₅₀ values may indicate the potential of a compound as cytotoxic. The higher the value of IC₅₀ against vero cells means the compounds are less toxic to normal cells³¹. The end of the cytotoxicity test can provide information on the percentage of cells that are able to survive. Table 3 shows the percentage of viable cells after administration of brusatol growing increasingly at lower concentrations. In brusatol concentration of 1200 µg mL⁻¹, the percentage of viable cells is 49.897±0.217%, while the lowest concentration of 37.5 µg mL⁻¹, the percentage of viable cells is 100±0.045%. Probit analysis results showed IC₅₀ value of brusatol against vero cells is 1402.029±0.164 µg mL⁻¹ but not significantly different (p>0.05). It means brusatol compound has cytotoxic effects categorized as low (low cytotoxicity)²⁶ because it has a value of IC₅₀>30 µg mL⁻¹. Similar observations have been reported by Mangungsong³² who reported that the compound of brusatol isolated from the seed of

B. javanica (L.) Merr. has a low cytotoxic effect on vero cells with IC₅₀ value of 1366.55 µg mL⁻¹.

CONCLUSION

This study can be concluded that the potential effect of brusatol as biolarvicide to control the mosquito vector with low levels of cytotoxicity on normal cells (Vero cells). The behavioral response of *A. aegypti* larvae showed symptoms of a neurotoxic that is excitation, convulsions, tremors and paralysis.

SIGNIFICANCE STATEMENT

Brusatol isolated from the seeds of Makasar fruit (*Brucea javanica* (L.) Merr.) is potentially effective and safe in controlling mosquito vector especially *A. aegypti*. Brusatol biolarvicide is proven significantly to have larvicidal effect on *A. aegypti* with the LC₅₀ of 0.669±0.106 ppm and LC₉₀ that is at 8.331±0.060 ppm.

ACKNOWLEDGMENTS

The authors would like to thank the Rector of University of Diponegoro who has funded this study through PNBIP DIPA University of Diponegoro funding No. SP DIPA-042.01.2.400898 / 2016, dated December 7, 2015 of fiscal year 2016.

REFERENCES

1. Ndione, R.D., O. Faye, M. Ndiaye, A. Dieye and J.M. Afoutou, 2007. Toxic effects of neem products (*Azadirachta indica* A. Juss.) on *Aedes aegypti* Linnaeus 1762 larvae. Afr. J. Biotechnol., 6: 2846-2854.
2. Daniel, 2008. When larvae and adult mosquitoes are immune to insecticides. Farmacia, 7: 43-50.
3. Lailatul, K.L., A. Kadarohman and R. Eko, 2010. Biolarvicidal effectivity of ethanol extract from the waste of Akar Wangi (*Vetiveria zizanioides*) oil distillation towards *Aedes aegypti*, *Culex* sp. and *Anopheles sundaicus* larvae. J. Sains Teknologi Kimia, 1: 59-65.
4. Moehammadi, N., 2005. The biolarvicidal potential of the extract of *Ageratum conyzoides* Linn and *Saccopetalum horsfieldii* Benn leaves against *Aedes aegypti* L. larvae. J. Biol. Res., 10: 1-4.
5. Isman, M.B., 2006. Botanical insecticides, deterrents and repellents in modern agriculture and an increasingly regulated world. Annu. Rev. Entomol., 51: 45-66.
6. Zhang, L., X. Feng, D. Ma, J. Yang, H. Jiang, Y. Zhang and W. He, 2013. Brusatol isolated from *Brucea javanica* (L.) Merr. induces apoptotic death of insect cell lines. Pesticide Biochem. Physiol., 107: 18-24.
7. Syahputra, E., 2008. [Bioactivity of *Brucea javanica* fruit as nabatical insecticide for agricultural insect pest]. Bul. Littro, 19: 56-67.
8. Lina, E.C., Dadang, S. Manuwoto, G. Syahbirin and D. Prijono, 2013. Synergistic action of mixed extracts of *Brucea javanica* (Simaroubaceae), *Piper aduncum* (Piperaceae) and *Tephrosia vogelii* (Leguminosae) against cabbage head caterpillar, *Crociodolomia pavonana*. J. Biopesticides, 6: 77-83.
9. Lina, E.C., Arnetti, D. Prijono and Dadang, 2010. Insecticidal activity of *Brucea javanica* L. Merr against cabbage pests *Crociodolomia pavonana* (Lepidoptera: Crambidae). Manggaro, 11: 36-39.
10. WHO., 2005. Guidelines for laboratory and field testing of mosquito larvicides. WHO/CDS/WHOPES/GCDPP/2005.13. http://libdoc.who.int/hq/2005/WHO_CDS_WHOPES_GCDP_P_2005.13.pdf
11. Sharma, A., S. Kumar and P. Tripathi, 2015. Impact of *Achyranthes aspera* leaf and stem extracts on the survival, morphology and behaviour of an Indian strain of dengue vector, *Aedes aegypti* L. (Diptera: Culicidae). J. Mosquito Res., Vol. 5. 10.5376/jmr.2015.05.0007
12. Van Meerloo, J., G.J.L. Kaspers and J. Cloos, 2011. Cell sensitivity assays: The MTT assay. Methods Mol. Biol., 731: 237-245.
13. Rueda, L.M., K.J. Patel, R.C. Axtell and R.E. Stinner, 1990. Temperature-dependent development and survival rates of *Culex quinquefasciatus* and *Aedes aegypti* (Diptera: Culicidae). J. Med. Entomol., 27: 892-898.
14. Barrera, R., M. Amador and G.G. Clark, 2006. Ecological factors influencing *Aedes aegypti* (Diptera: Culicidae) productivity in artificial containers in Salinas, Puerto Rico. J. Med. Entomol., 43: 484-492.
15. Becker, N., D. Petric, M. Zgomba, C. Boase, M. Minoo, C. Dahl and A. Kaiser, 2010. Mosquitoes and their Control. 2nd Edn., Springer, New York, ISBN: 9783540928744, Pages: 577.
16. Choochote, W., B. Tuetun, D. Kanjanapothi, E. Rattanachanpichai and U. Chaithong et al, 2004. Potential of crude seed extract of celery, *Apium graveolens* L., against the mosquito *Aedes aegypti* (L.) (Diptera: Culicidae). J. Vector Ecol., 29: 340-346.
17. Choochote, W., U. Chaithong, K. Kamsuk, E. Rattanachanpichai and A. Jitpakdi et al, 2006. Adulticidal activity against *Stegomyia aegypti* (Diptera: Culicidae) of three Piper spp. Rev. Inst. Med. Trop. S. Paulo, 48: 33-37.
18. Dharmagadda, V.S.S., S.N. Naik, P.K. Mittal and P. Vasudevan, 2005. Larvicidal activity of *Tagetes patula* essential oil against three mosquito species. Bioresour. Technol., 96: 1235-1240.

19. Chaithong, U., W. Choochote, K. Kamsuk, A. Jitpakdi and P. Tippawangkosol *et al.*, 2006. Larvicidal effect of pepper plants on *Aedes aegypti* (L.) (Diptera: Culicidae). *J. Vector Ecol.*, 31: 138-144.
20. Kumar, S., R. Warikoo and N. Wahab, 2010. Larvicidal potential of ethanolic extracts of dried fruits of three species of peppercorns against different instars of an Indian strain of dengue fever mosquito, *Aedes aegypti* L. (Diptera: Culicidae). *Parasitol. Res.*, 107: 901-907.
21. Warikoo, R. and S. Kumar, 2013. Impact of *Argemone mexicana* extracts on the larval, morphological and behavioral response of dengue vector, *Aedes aegypti* L. (Diptera: Culicidae). *Parasitol. Res.*, 112: 3477-3484.
22. Sanjaya, Y. and T. Safaria, 2006. [Toxicity of poison from spider *Nephila* sp. to *Aedes aegypti* L. larvae]. *J. Biodiversitas*, 7: 191-194.
23. Yu, S.J., 2008. *The Toxicology and Biochemistry of Insecticides*. CRC Press/Taylor and Francis Group, New York, USA., ISBN-13: 9781420059755, Pages: 276.
24. Boublik, Y., P. Saint-Aguet, A. Lougarre, M. Arnaud, F. Villatte, S. Estrada-Mondaca and D. Fournier, 2002. Acetylcholinesterase engineering for detection of insecticide residues. *Protein Eng.*, 15: 43-50.
25. Kim, I.H., R. Suzuki, Y. Hitotsuyanagi and K. Takeya, 2003. Three novel quassinoids, javanicolides A and B and javanoside A, from seeds of *Brucea javanica*. *Tetrahedron*, 59: 9985-9989.
26. Lu, F.C. and S. Kacew, 2002. *Lu's Basic Toxicology: Fundamentals, Targets Organ and Risk Assessment*. 4th Edn., Taylor and Francis, New York, Pages: 386.
27. Kringer, R., 2010. *Handbook of Pesticide Toxicology*. Academic Press, California, pp: 120-132.
28. Govindachari, T.R., G.N.K. Kumari, G. Gopalakrishnan, G. Suresh, S.D. Wesley and T. Sreelatha, 2001. Insect antifeedant and growth regulating activities of quassinoids from *Samadera indica*. *Fitoterapia*, 72: 568-571.
29. Uthai, U., P. Rattanapreechachai and L. Chohanadisai, 2011. Bioassay and effective concentration of Temephos against *Aedes aegypti* larvae and the adverse effect upon indigenous predators: *Toxorhynchites splendens* and *Micronecta* sp. *Asia J. Public Health*, 2: 67-77.
30. Meiyanto, E., Sismindari, L. Candra and Moordiani, 2003. [Antiproliferative effect of the bark and leaves of erythrina fusca lour against hela cells]. *Majalah Farmasi Indonesia*, 14: 124-131.
31. Muhtadi, 2008. Pemisahan fraksi dan senyawa-senyawa yang berkhasiat antiplasmodium dari ekstrak metanol kulit kayu mimba (*Azadirachta indica* Juss). *J. Penelit Sains Dan Teknol.*, 9: 117-136.
32. Mangungsong, S., 2012. Aktivitas senyawa semisintetik kuasinoid dari Buah Makasar (*Brucea javanica* (L.) Merr.) sebagai antikanker dengan target protein p53, bcl-2, kaspase-3, COX-2 dan c-Myc. Ph.D. Thesis, Universitas Gadjah Mada, Indonesia.

Effect of BrusatolBiolarvicide Administration on BEhavioral Response of Aedes aegypti and its Toxicity on Vero Cells

ORIGINALITY REPORT

17%

SIMILARITY INDEX

10%

INTERNET SOURCES

9%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Udayana University Student Paper	2%
2	Lan Zhang, Xuehuan Feng, Dejun Ma, Jingjing Yang, Hongyun Jiang, Yanning Zhang, Weizhi He. "Brusatol isolated from Brucea javanica (L.) Merr. induces apoptotic death of insect cell lines", Pesticide Biochemistry and Physiology, 2013 Publication	2%
3	eprints.undip.ac.id Internet Source	2%
4	biopublisher.ca Internet Source	1%
5	docsdrive.com Internet Source	1%
6	Dharmagadda, V.S.S.. "Larvicidal activity of Tagetes patula essential oil against three mosquito species", Bioresource Technology, 200507 Publication	1%

7	Isnatin Miladiyah, Jumina Jumina, Sofia Mubarika Haryana, Mustofa Mustofa. "Biological activity, quantitative structure–activity relationship analysis, and molecular docking of xanthone derivatives as anticancer drugs", Drug Design, Development and Therapy, 2018 Publication	1 %
8	Submitted to Politeknik Kesehatan Kemenkes Semarang Student Paper	<1 %
9	media.neliti.com Internet Source	<1 %
10	Suwandi, Jhons Fatriyadi, Supargiyono, Widya Asmara, and Hari Kusnanto. "Mapping and Prevalence of Malaria Falciparum Patients with ACT Failed Therapy, in Hanura Public Health Center, Pesawaran, Lampung, Indonesia", Open Journal of Epidemiology, 2014. Publication	<1 %
11	link.springer.com Internet Source	<1 %
12	scholar.unand.ac.id Internet Source	<1 %
13	www.j3.jstage.jst.go.jp Internet Source	<1 %

14

repository.seafdec.org.ph

Internet Source

<1 %

15

Arif Susanto. "Assessment of Diesel Particulate Matter Exposure of Underground Miners in Indonesia", Journal of Ecological Engineering, 2018

Publication

<1 %

16

www.latamjpharm.org

Internet Source

<1 %

17

Ik Hwi Kim, Satoru Takashima, Yukio Hitotsuyanagi, Tomoyo Hasuda, Koichi Takeya. " New Quassinoids, Javanicolides C and D and Javanicosides B–F, from Seeds of ", Journal of Natural Products, 2004

Publication

<1 %

18

Kurniawan Teguh Martono, Cahya Setya Utama, Bambang Sulistiyanto, Merry Christiyanto. "Utilization of social media in livestock product marketing group of cattle", 2016 3rd International Conference on Information Technology, Computer, and Electrical Engineering (ICITACEE), 2016

Publication

<1 %

19

Submitted to University of Dehli

Student Paper

<1 %

20

www.hec.gov.pk

Internet Source

<1 %

21	Submitted to Universitas Indonesia Student Paper	<1 %
22	Submitted to Turun yliopisto Student Paper	<1 %
23	Submitted to University of Hong Kong Student Paper	<1 %
24	"Natural Products", Springer Nature, 2013 Publication	<1 %
25	www.mysciencework.com Internet Source	<1 %
26	ejomr.org Internet Source	<1 %
27	www.entomoljournal.com Internet Source	<1 %
28	Submitted to Universiti Putra Malaysia Student Paper	<1 %
29	benthamopen.com Internet Source	<1 %
30	Submitted to Touro College Student Paper	<1 %
31	Hye-Mi Park, Il-Kwon Park. "Larvicidal activity of <i>Amyris balsamifera</i> , <i>Daucus carota</i> and <i>Pogostemon cablin</i> essential oils and their components against <i>Culex pipiens pallens</i> ", <i>Journal of Asia-Pacific Entomology</i> , 2012 Publication	<1 %

32

www.ijcrbp.com

Internet Source

<1 %

33

Submitted to Liverpool John Moores University

Student Paper

<1 %

34

www.science.gov

Internet Source

<1 %

35

Rujun Ma, Hongru Li, Yu Zhang, Ying Lin, Xuhua Qiu, Min Xie, Bing Yao. "The toxic effects and possible mechanisms of Brusatol on mouse oocytes", PLOS ONE, 2017

Publication

<1 %

36

Paweł Czerniewicz, Grzegorz Chrzanowski, Iwona Sprawka, Hubert Sytykiewicz.

"Aphicidal activity of selected Asteraceae essential oils and their effect on enzyme activities of the green peach aphid, *Myzus persicae* (Sulzer)", Pesticide Biochemistry and Physiology, 2018

Publication

<1 %

37

pingpdf.com

Internet Source

<1 %

38

Langdon, C.J.. "Survival, Pb-uptake and behaviour of three species of earthworm in Pb treated soils determined using an OECD-style toxicity test and a soil avoidance test", Environmental Pollution, 200511

Publication

<1 %

39	Submitted to School of Business and Management ITB Student Paper	<1 %
40	www.repository.ugm.ac.id Internet Source	<1 %
41	K. Pavanand. " <i>In Vitro</i> Antimalarial Activity of <i>Brucea javanica</i> against Multi-Drug Resistant <i>Plasmodium falciparum</i> ", <i>Planta Medica</i> , 04/1986 Publication	<1 %
42	www.spandidos-publications.com Internet Source	<1 %
43	Submitted to Delhi University Student Paper	<1 %
44	Submitted to The Hong Kong Polytechnic University Student Paper	<1 %
45	Raymond, Demba Ndione, Faye Omar, Ndiaye Mady, Dieye Abdoulaye, and Marie Afoutou Joseacute. "Toxic effects of neem products (<i>Azadirachta indica</i> A. Juss) on <i>Aedes aegypti</i> Linnaeus 1762 larvae", <i>AFRICAN JOURNAL OF BIOTECHNOLOGY</i> , 2007. Publication	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Effect of BrusatoI Biolarvicide Administration on BEhavioral Response of Aedes aegypti and its Toxicity on Vero Cells

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10
