

## **URUAE Int'l Conference Proceedings**

**5th International Conference on Computer, Electronics and  
Manufacturing Engineering (ICCEME-16)**

**5th International Conference on Food, Agricultural and  
Biological Sciences (ICFABS-2016)**

**&**

**International Conference on Nanotechnology, Environmental  
and Civil Engineering (ICNECE-2016)**



**Dec. 25-26, 2016 Bangkok (Thailand)**

**Editors:**

**Prof. Dr. Naime Arslan**

**Dr. Thaweesak Yingthawornsuk**

**ISBN 978-93-84468-88-0**

**ORGANISED BY :**



***5th International Conference on Computer, Electronics and  
Manufacturing Engineering (ICCEME-16)***

***5th International Conference on Food, Agricultural and  
Biological Sciences (ICFABS-2016)***

**&**

***International Conference on Nanotechnology, Environmental  
and Civil Engineering (ICNECE-2016)***

**Dec. 25-26, 2016**

**Bangkok (Thailand)**

**Editors:**

**Prof. Dr. Naime Arslan**

**Dr. Thaweesak Yingthawornsuk**

**ISBN 978-93-84468-88-0**

**Organized By:**



**URUAE.ORG**

## Program Committee (ICCEME-16)

- Prof. Dr. Barbara Marchetti, Engineering Faculty Università degli Studi eCampus, Italy  
Prof. Dr. Abdelkader Adia, Department of Computer Science, University of Oran, Algeria  
Prof. Dr. BOUDEN Toufik, Ndi Laboratory, Automatic Department, Jijel University, Algeria  
Prof. Dr. DALAH Mohamed, Department of Mathematics, Faculty of Sciences, University Constantine 1, Algeria  
Prof. Dr. Dimiter Velov, University of National and World Economy, Sofia, Bulgaria  
Prof. Dr. Hai Ho, Kennesaw State University, Southern Polytechnic College of Engineering and Engineering Technology, USA  
Prof. Dr. Hufang DENG, South China University of Technology, China  
Prof. Dr. Jimmy (C.M.) Kao, Distinguished Professor, National Sun Yat-Sen University Coordinator of Environmental Engineering Program, National Science Council, Taiwan  
Prof. Dr. Kei Eguchi, Department of Information Electronics, Fukuoka Institute of Technology, Japan  
Prof. Dr. Maher Mohamed Nofel, Mechanical Engineering Department, Taibah University, KSA  
Prof. Dr. Miroslav V. Tomić, University of Belgrade, Faculty of Mechanical Engineering, Serbia  
Prof. Dr. Mohd Nasir Taib, Universiti Teknologi MARA (UiTM), Malaysia  
Prof. Dr. Osman ADIGUZEL, Department of Physics, Firat University, Elazığ, Turkey  
Prof. Dr. Valentina Mihaela Pomazan, University Ovidius of Constanta, Faculty of Mechanical and Industrial Engineering, Constanta, Romania  
Prof. Jemel Antidze, Vekua Scientific Institute of Applied Mathematics, Tbilisi State University, Tbilisi, Georgia  
Prof. Mihai Caramihai, University POLITEHNICA Bucharest, Romania  
Prof. Okba KAZAR, Computer Sciences Department, University of Biskra, Algeria  
Prof. Samy Oraby, Manufacturing Engineering Technology, College Technological Studies, PAAET, Kuwait  
Prof. Susantha N. De Silva, Institute of Automotive Engineers, Sri Lanka  
Prof. Biljana Radulovic, University of Novi Sad, Serbia  
Prof. Dae-sun Hong, Changwon National University, Korea  
Asso. Prof. Dr. Nechoi A. Al-Sayid, Faculty of Computer Science, Applied Science University, Jordan  
Asso. Prof. Dr. Florin Negoescu, Department of Machine Manufacturing Technology, Technical University "Gheorghe Asachi" of Iasi, Romania  
Asso. Prof. Dr. Nayef Ghazem, Associate Professor, United Arab Emirates University, UAE  
Asso. Prof. Ir Nik Ghazali Nik Daud, Department of Electrical & Electronic Engineering, Faculty of Engineering, National Defence University of Malaysia  
Asso. Prof. Shih-Chuan Yeh, Department of Electronic Engineering, De Lin Institute of Technology, Taiwan  
Assoc. Prof. Yuan-Wei Tseng, Department of Electrical Engineering, I-Shou University, Taiwan, R.O.C  
Asst. Prof. I. Erika Böyükdaşkın, Wichita State University, USA  
Dr. Malik Abdulrazzaq Alseedi, Department of Electrical, College of Engineering, University of Misan, Iraq  
Dr. Prof. Akos Lakatos, University of Debrecen, Faculty of Engineering, Debrecen, Hungary  
Dr. Prof. Ayman Batiha, International Sustainability Institute, Cairo, Egypt  
Dr. Yu-Kuang Zhao, Associate Professor, Department of Refrigeration, Air Condition and Energy Engineering, National Chin-Yi University of Technology, Taiwan

# Program Committee (ICFABS-2016)

- Prof. Asep K. Supriatna, Padjadjaran University, Indonesia
- Prof. Dr. Alaa H. Wadie Al-Fatlawi, Environmental Engineering Department, College of Engineering, University of Babylon, Iraq
- Prof. Dr. Assam El-Osta, Baker IDI Heart & Diabetes Institute, Melbourne, Australia
- Prof. Dr. Bülent TOPCUOĞLU, Akdeniz University, Vocational School of Technical Sciences, Antalya, Turkey
- Prof. Dr. Chairil Anwar, Medical Faculty of Sriwijaya University, Indonesia
- Prof. Chung, Ying-Chien, Department of Biological Science and Technology, China University of Science and Technology, Taiwan (ROC)
- Prof. Dr. Emhemmed Abdullah Alhibshi, Chemistry department, Faculty of Science, Aljabel Algharbi University, Libya
- Prof. Dr. Gary Sweeney, York University, Toronto, Canada
- Prof. Dr. Geraldine Apelo-Quinones, Department of Biological Science, College of Science, University of Eastern Philippines (UEP), Philippines
- Prof. Dr. Gopal Natesan, Department of Medicinal Chemistry, Faculty of Pharmacy, MAHSA University, Malaysia
- Prof. Dr. Heecheul Kim, Professor, Kyung Hee University, Yongin-si, Gyeonggi-do, Republic of Korea
- Prof. Dr. Mohamed Abd Ellatif Said, College of Medicine, King Khalid University, Abha, Saudi Arabia & Mansoura Faculty of Medicine, Mansoura University, Egypt
- Prof. Dr. Mohamed Elbagemi, Department of Chemistry, Faculty of science, University of Misurata, Iraq
- Prof. Dr. Rahim Ahmadi, Islamic Azad University, Hamedan Branch, Iran; International Avicenna College, Hungary, European Union
- Prof. Dr. Samia M. Sanad, Zoology Department, Zagazig University, Egypt
- Prof. Dr. Sawsan Sajid Al-Jubori, Professor, Department of Biology, Biotechnology College of science, Al – Mustansiriya University, Iraq
- Prof. Dr. Tank Shantilal K., Department of Biosciences, V.N.South Gujarat University, Surat, India
- Prof. Dr. Yuei-An Liou, National Central University, Taiwan
- Prof. Dr. Sanaâ SAOIABI, Department of Chemistry, Faculty of Sciences, Mohammed V University Rabat, Morocco
- Prof. Francisco Javier Navarro Domínguez, Department of Chemical Engineering, University of Huelva, Campus el Carmen, Spain
- Prof. Jamaan S. Ajarem, King Saud University, Saudi Arabia
- Prof. Dr. Yalcin Kaya, Engineering Faculty Genetic Engineering Department, Turkey
- Assoc. Prof. Dogan Ozdemir, Ishik University, Erbil, Iraq
- Assoc. Prof. Ahmed Saber Abu-zaiton, Al-albayt University, Jordan
- Assoc. Prof. Dr. Kumaran Narayanan, Monash University Malaysia, Malaysia
- Asst. Prof. Dr. Afaf Ghais Abadi Ahmed, Chemical Engineering Department, Faculty of Engineering, Sudan
- Assoc. Prof. Dr. Khalid Rehman Hakeem, Faculty of Forestry, Universiti Putra Malaysia
- Assoc. Prof. Dr. Trt Retnantnpih Soeprbowati, University of Diponegoro, Semarang, Indonesia
- Assoc. Prof. Dr. Alaa Kareem Niamah, Food Science Department, Agriculture college, Basrah university, Iraq
- Assoc. Prof. Mahad Baawain, Sultan Qaboos University, Oman
- Assoc. Prof. Mariam Gaidamashvili, Associate Professor, Department of Biology, Faculty of Exact and Natural Sciences, Javakhishvili Tbilisi State University, Georgia
- Assoc. Prof. Ahmed Abdel-Megeed, Department of Botany and Microbiology, Faculty of Science, King Saud University, Riyadh, Saudi Arabia
- Assoc. Prof. Dr. Rym SALAH TAZDAÏT, Mouloud Mammeri University of Tizi-Ouzou, Algeria
- Assoc. Prof. Dr. Tri Retnaningsih Soeprbowati, Biology, University of Diponegoro, Semarang, Indonesia
- Assoc. Prof. Yuji Sakai, Department of Environmental and Energy Chemistry, Kogakuin University, Japan
- Assoc. Prof. Ass Prof. Sello Alfred LIKUKU, Department of Basic Sciences, Botswana College of Agriculture, Botswana

# Program Committee (ICNECE-2016)

Prof. Dr. Ahmed Helmy Othman, Civil Engineering Department, College of Engineering, Najran University, Najran, KSA

Prof. Dr. Alaa H. Wadie Al-Fatlawi, Environmental Engineering Department, College of Engineering, Babylon University, Iraq

Prof. Dr. Belhadj Safia, Department of Agronomy, Faculty of Natural Sciences and Life, University of Djelfa, Algeria

Prof. Dr. Bozena Lozowicka, Plant Protection Institute –National Research Institute, Chelmonskiego, Bialystok, Poland

Prof. Dr. José Manuel Gómez-Soberón, Department of Architectural Technology, Superior School of Building, Technical University of Catalonia, Spain

Prof. Dr. Mostafa Abd-El-Azeem Abo-El-Soud Amer, Agricultural Botany Department Faculty of Agriculture (Saba-Basha), Alexandria University Egypt

Prof. Dr. Wen-Pei Sung, Integrated Research Center for Green Living Technologies, Dept. of Landscape Architecture, National Chin-Yi University of Technology, Taiwan

Prof. Edward A. Cook, Arizona State University, USA

Assoc. Prof. Mohamed Ali Elssaidi, Environmental Science Dept., Faculty of Engineering and Technology, Sebha University, Libya

Assoc. Prof. Dr. Ajay Raghunath Tembhurkar, Head, Department of Civil Engineering, Visvesvaraya National Institute of Technology, Maharashtra India

Assoc. Prof. Dr. Mohd Irwan Juki, Universiti Tun Hussein Onn Malaysia, Malaysia

Assoc. Prof. Dr. Norzila Othman, Universiti Tun Hussein Onn Malaysia, Malaysia

Asst. Prof. Dr. Ahmed Mohammed, Dept. of Plant Production, College of Agriculture and Veterinary Medicine, Ibb University, Yemen

Asst. Prof. Dr. Aynur DEMIR, Department of Environmental Protection Technologies, Aksaray University, Aksaray, TURKEY

Asst. Prof. Dr. Donludee Jaisut, Faculty of Agriculture, Kasetsart University, Bangkok, Thailand

Asst. Prof. Dr. Sherif El-Sayed El-Gamal, Civil and Architectural Engineering Department, Sultan Qaboos University, Sultanate of Oman

Asst. Prof. Ikbāl ERBAS, Department of Architect, Faculty of Architecture, Akdeniz University, Turkey

Dr. Elmira Shamshiry, Research Management Center (RMC), University Terengganu Malaysia (UMT), Malaysia

Dr. Osama El Gelani Elsaid, Faculty of Agricultural Technology and Fish Sciences, Al-Neelain University, Khartoum, Sudan

Kemal Reha Kavas, Department of Architecture, Akdeniz University, Turkey

## PREFACE

Dear Distinguished Delegates, Colleagues and Guests,

The URUAE Organizing Committee warmly welcomes our distinguished delegates and guests at Universal Researchers (UAE), ICCEME-16, ICFABS-2016 and ICNECE-2016 Int'l Conferences scheduled on *Dec. 25-26, 2016 Bangkok (Thailand)*. The main themes and tracks are Computer, Electronics, Manufacturing Engineering, Food, Agricultural, Biological Sciences, Nanotechnology, Environmental and Civil Engineering.

These conferences are managed and sponsored by Universal Researchers (UAE) and assisted by University of Johannesburg and University of Quebec. URUAE is striving hard to compile the research efforts of scientists, researchers and academicians across the broad spectrum of Science, Engineering and Technology. These conferences are aimed at discussing the wide range of problems encountered in present and future high technologies among the research fraternity.

The conferences are organized to bring together the members of our international community at a common platform, so that, the researchers from around the world can present their leading-edge work. This will help in expansion of our community's knowledge and provide an insight into the significant challenges currently being addressed in that research. The conference Program Committee is itself quite diverse and truly international, with membership from the America, Australia, Europe, Asia and Africa.

The conferences has solicited and gathered technical research submissions related to all aspects of major conference themes and tracks. This proceeding records the fully refereed papers presented at the conference.

All the submitted papers in the proceeding have been peer reviewed by the reviewers drawn from the scientific committee, external reviewers and editorial board depending on the subject matter of the paper. Reviewing and initial selection were undertaken electronically. After the rigorous peer-review process, the submitted papers were selected on the basis of originality, significance, and clarity for the purpose of the conference. The main goal of these events is to provide international scientific forums for exchange of new ideas in a number of fields that interact in-depth through discussions with their peers from around the world.

The program has been structured to favor interactions among attendees coming from many diverse horizons, scientifically, geographically, from academia and from industry. We would like to thank the program chairs, organization staff, and the members of the program committee for their work. We like to thank and show gratitude to Editors from URUAE. We are grateful to all those who have contributed to the success of URUAE Dec. 25-26, 2016 Bangkok (Thailand) Conference. We hope that all participants and other interested readers benefit scientifically from the proceedings and also find it stimulating in the Process in their quest of achieving greater heights. Finally, we would like to wish you success in your technical presentations and social networking.

We hope you have a unique, rewarding and enjoyable week at URUAE Conferences at vibrant Bangkok.

With our warmest regards,

*Organizing Committee*

*Dec. 25-26, 2016  
Bangkok (Thailand)*

# INDEX

**Dec. 25-26, 2016 Bangkok (Thailand)**

<b>Paper ID</b>	<b>Title of the Paper and Authors</b>	<b>Page No.</b>
-----------------	---------------------------------------	-----------------

## **5th International Conference on Computer, Electronics and Manufacturing Engineering (ICCEME-16)**

<b>AE1216101</b>	Enhanced Interactive Multimedia System as an Additional Learning Tool for Pre-Scholars Robyn N. Letsoalo, and Johnson O. Dehinbo	<b>1</b>
<b>AE1216103</b>	Traffic Accident Simulation System with Pedestrian View for Traffic Safety Education Suguru Tsujioka, Satoshi Togawa, Hirofumi Hirose, Tetsuo Naka, and Akihide Takeda	<b>9</b>
<b>AE1216106</b>	Modified K-Means for Document Clustering System Tin Thu Zar Win, and Moe Moe Aye	<b>13</b>
<b>AE1216110</b>	A simple Grounded Fractional-order Capacitance Multiplier employing only single CFTA T. Comedang, and P. Intani	<b>19</b>
<b>AE1216111</b>	Grounded Fractional-order Inductance Simulation Using Single CCCFTA T. Comedang, and P. Intani	<b>23</b>
<b>AE1216112</b>	Effect of Heat Treatment Parameters on Microstructure, Mechanical and Wear Properties of Al-7%Si-0.5%Mg reinforced with Dual Size SiCp Avinash L, Srikanth Bontha, Abhishek Nagendra, Nihal R Rao, H.N.Reddappa, and Nagaraja C Reddy	<b>27</b>
<b>AE1216109</b>	A Study on Detecting a Planing Vessel's Motion using a Fusion System with Low-Cost Sensors YoungHun Kim, HyeonKyu Yoon, and NamHyun Yoo	<b>32</b>

## **5th International Conference on Food, Agricultural and Biological Sciences (ICFABS-2016)**

<b>AE1216210</b>	The Management of Organic Wastes Produced Vermicompost Using Earthworm <i>Eudrilus eugeniae</i> and Effects of Vermicompost on Growth of <i>Helianthus annuus</i> Pitchya Tangsombatvichit, Sakuntep Chupong, Daojarus Ketrot, and Kitti Boonlerthirun	<b>33</b>
<b>AE1216214</b>	Screening on Lipase-producing Bacteria Isolated from Oil Mill Soil Khaing Phyo Wai, Weine Nway Nway Oo, and Moe Thazin Shwe	<b>37</b>
<b>AE1216215</b>	Development of Photoperiod-insensitive Mutants from Rice var. Ayarmin (Machando) by Using Gamma Radiation Nay Aung, Aye Aye Khai, and Myat Minn	<b>40</b>
<b>AE1216216</b>	Investigation of Heavy Metal Removal by <i>Sargassum</i> spp from Industrial Wastewater Hlaing Swe Win, Saw Sandar Maw, and Hnin Hnin Htun	<b>46</b>
<b>AE1216217</b>	Investigation of Toxic Heavy Metal Removal Activity of Activated Coconut Husk as Solid Support Medium for Isolated Bacterial Strains Sa Aung Ko Ko, Saw Sandar Maw, and Zaw Khaing Oo	<b>50</b>

Paper ID	Title of the Paper and Authors	Page No.
AE1216218	<i>Cytotoxic Effect of a Mycelial extracts on Different Cell lines and Experimental Animals</i> Osman, M.E., Wafaa Ahmed., Heba, E.E. and Om Hashim.M	54
AE1216219	<i>Chicken Eggshell Waste as a Suitable Catalyst for Transesterification of Palm Oil: Optimization for Biodiesel Production</i> Thi Thi Win, and May Myat Khine	60
AE1216231	<i>Graft Copolymerization of Acrylic Monomers onto Coir Dust by Using Gamma Irradiator of Co-60 Source for Superabsorbent Polymer</i> Min Naing, and Khin Khin Lay	66
AE1216234	<i>Biological Control and Characterization of Mycelium Forming Bacteria Causing Potato Diseases</i> Gholam Khodakaramian and Nasim Khodakaramian	72
AE1216236	<i>Ameliorative Role of Moringa oleifera Plant Extract Against Zinc Oxide Nanoparticles Induced Sperm and Sex Hormones Abnormalities in Male Albino Rats</i> Sabha Elsayed Elballat	76
AE1216237	<i>Some New Parameters Can Be Used In Diagnosis of Subclinical Mastitis in Buffaloes</i> Salama, E. M., and Saad A.H	81
AE1216238	<i>Effect of Some Natural Extracts On Shelf Life of Chilled Lucioperca Lucioperca Fillets</i> Mohamed A. Hussein, Elsaid A. Eldaly, and Huda K. Hussein	86
AE1216239	<i>Effect of Mirazid on the Micronuclei Polychromatic Erythrocyte Cells (MNPCEs) In the Bone Marrow Cells of Schistosoma Infected and Non Infected Mice</i> Fawzia Abd Elhady Zayed, Hasanat El-Sayed Metwally ALI, and Amani Abdelhamid Mahmoud	91
AE1216240	<i>Plasma Lipid Profiles, Milk Yield and Milk Fat Concentration of Lactating Holstein Fed Rumen-Protected Choline</i> Indra S. Hesti, D. Trizana, A. Wicaksono, P. Sambodho, Christina B. Soejono, Sri A. B. Santoso, A. Subrata, and Dian W. Harjanti	96
AE1216208	<i>Antioxidant activity of Sri Lankan traditional rice (Oryza sativa L.) and the effect of parboiling</i> Thiloka I. Kariyawasam, Pahan I. Godakumbura, M. A. B. Prashantha, and G. A. S. Premakumara	101
AE1216211	<i>Correlation between Total Phenolic and Flavonoid Contents with Antioxidant Activity of Three Different Propolis Derived from Trigona apicalis, Trigona itama and Trigona thoracica</i> N. Asem, N. Abd Hapit, N. A. Abdul Gapar, and E. A. Omar	102
AE1216222	<i>The Application of the Theory of Planned Behavior to Predict Blood Glucose Monitoring In Type 2 Diabetic Patients</i> Ali karimi, sirous shasavari, Arezoo Fallahi, and Tahereh Pashaei	103
AE1216224	<i>Exploring the Perspectives of Students at the Kurdistan University of Medical Sciences about Spiritual Vitality</i> Roya Ebrahimi, Pari Taymoori, Maryam khazaei pool, and Tahereh Pashaei	104
AE1216225	<i>Prevalence and Factors Associated With Self-Medication with Antibiotics in Sanandaj City</i> Tahereh Pashaei, Sharareh Baghari, and Roya Ebrahimi	105

# Plasma Lipid Profiles, Milk Yield and Milk Fat Concentration of Lactating Holstein Fed Rumen-Protected Choline

Indra S. Hesti, D. Trizana, A. Wicaksono, P. Sambodho, Christina B. Soejono, Sri A. B. Santoso, A. Subrata, and Dian W. Harjanti\*

**Abstract** — Choline plays an essential role in fat metabolism in the liver. The purpose of this study was to evaluate the effects of addition rumen-protected choline (RPC) on plasma lipid profiles (concentrations of plasma cholesterol, low density lipoprotein (LDL), high density lipoprotein (HDL) and triglycerides), milk yield and milk fat concentration in lactating Holstein. Eight of lactating Holstein (in the 2<sup>nd</sup> lactation period and  $456 \pm 31$  kg of body weight) used. The cows were fed two diets in a crossover experiment for 90 days. The first diet was total mixed ration (TMR) containing Napier grass and concentrate (40:60) as control diet, and the second diet was TMR added with RPC as choline chloride 30 g/head/d (choline chloride 60% corn-cob as equal to 0.02% BW<sup>0.75</sup>). The data was analyzed with one simple T-Test using SPSS ver.21. The results showed that addition RPC in the diet did not significantly influence ( $p > 0.05$ ) the concentrations of plasma lipid profiles. The inclusion of RPC in the diet increased ( $p < 0.05$ ) milk yields up to 0.38 kg/d and increased milk fat concentration up to 0.35%/d. Thus, choline chloride could be added in the diets of lactation cows to increase milk yield and milk fat.

**Keywords** — choline chloride, plasma lipid, milk yield, milk fat, dairy cows.

## I. INTRODUCTION

In the tropic region, milk yield of dairy cow is relatively low and could not meet the total consumption need. The quality of feeding is important to improve the quantity and quality of milk produced. Choline has been classified as one of the B-complex vitamins, but it's determine was not complied into the definition of vitamins whose had enzyme co-factors [24]. Choline is formed as lecithin on feed and part of phosphatidylcholine whose the main phospholipid in cell membrane, lipoproteins, and milk fat globule on dairy cows [15]. Choline plays an essential role in fat metabolism in the liver by increasing the utilization of fatty acids in the liver. Moreover, choline contributes in lipid metabolism that serve Acetyl CoA as the precursor of cholesterol synthesis [10], as

phosphatidylcholine (the main part of phospholipid that used in lipid transport) and as methyl group donor in metabolism (methionine, folic acids and vitamin B12) [24]. Methionine is used in very low density lipoprotein (VLDL) synthesis in liver then will be converted into LDL in blood. Methylation pathways form phosphatidylcholine, the main phospholipid of HDL. Vitamin B12 is needed as co-enzyme in methylation pathways and used to be precursor of gluconeogenesis from propionate. Highly energy efficiency implicated to increasing milk yield and highly lipid anabolism implicated to increasing milk fat. Because of that, choline become to be feed additive that need in dairy cows feed. Choline had been fully degraded in the rumen because of that choline have to be protected as Rumen Protected Choline (RPC) or choline chloride 60% corn-cob (contained 60% choline chloride mixed with 40% corn-cob as carrier) that choline can pass through reticulorumen and absorbed into small intestine [4,15]. Addition of choline chloride in feed is expected to optimize methyl group metabolism in order to prevent negative energy balance which can improve the increasing milk yield and to optimize lipid metabolism that the transport of fat milk precursor to the mammary gland will increase which the indicator is lipid plasma (cholesterol, triglycerides, LDL, and HDL in blood).

We hypothesized that addition choline chloride can improve the energy metabolism in order to improve the milk yield in dairy cows and increase plasma lipid within normal range in order to improve the milk fat of dairy cows. This research is expected that choline chloride can be incorporated into feed and could improve total milk production and milk fat.

## II. MATERIALS AND METHODS

### A. Animals diets

This research was used eight of Lactating Holstein cows (61 to 91 days in milk; 3<sup>th</sup> lactation period;  $13.66 \pm 1.26$  kg/head/d of milk; and  $456 \pm 31$  kg of BW =  $99 \pm 5$  kg BW<sup>0.75</sup>) and held in dairy farm on Central Java, Indonesia. The proximate analysis of diet was listed in Table 1. The diet was total mixed ration (TMR) with the ingredients were Napier grass and concentrate (40:60) which was assumed from the National Research Council [12], as shown in Table 2. Additive feed was RPC as choline chloride 60% corn-cob 30 g/head/d (as equal to 0.02% BW<sup>0.75</sup>) referred by research before about RPC [9,15,23]. The

Manuscript received December, 1, 2016. This work was supported in part by Faculty of Animal and Agricultural Sciences, Diponegoro University, for facility of study and World Class University of Diponegoro University, Indonesia, for funding the paper.

I. S. Hesti is with the Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University.

\*Corresponding Author: D. W. Harjanti was with the Department of Animal Science, Faculty of Agriculture, Iwate University, Morioka 020-8550, Japan. She is now with Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang 50275, Indonesia

<https://doi.org/10.17758/URUAE.AE1216240>

TABLE I: THE PROXIMATE ANALYSIS OF DIETS INGREDIENTS

Nutrient composition	Napier grass	Concentrate
DM <sup>a</sup> (%)	15.32	88.39
TDN (%)	57.33 <sup>b</sup>	69.54 <sup>c</sup>
CP <sup>d</sup> (%)	8.01	16.70
CF <sup>a</sup> (%)	33.06	19.92
EE <sup>a</sup> (%)	2.72	4.55

<sup>a</sup>Nutrient and Feed Science Laboratory, Faculty of Animal and Agricultural Sciences Diponegoro University, (2015).

<sup>b</sup>TDN count based on Hartadi et al., (1980).

<sup>c</sup>TDN count based on Sutardi, (1979).

<sup>d</sup>Ecology and Plant Production Laboratory, Faculty of Animal and Agricultural Sciences, Diponegoro University, (2015).

DM : dry matter

TDN : total digestible nutrient

CP : crude protein

CF : crude fiber

EE : ether extract

RPC was purchased from Feed Grade Vitamins (Hebei Kangdali Pharmaceutical Co., Ltd, Hebei, China). The cows were fed two diets in a crossover experiment for 90 days (throughout into 2 periods) that consisted of 30 days of dietary adaptation and 60 days of dietary treatment. The diet treatments consisted of control diet (TMR) and RPC diet (TMR with choline chloride). Feeding of Napier grass was given twice per day on 08.00 a.m. and 04.00 p.m. then concentrate was given 4 times on 04.00 a.m., 07.00 a.m., 01.00 p.m. and 03.00 p.m. Adding RPC was given in the morning after 2nd concentrate on 07.00 a.m. had done. The water available was ad-libitum with nipple cup.

#### B. Sample collection and analysis

Blood samples were collected every 30 days and were taken from jugular vein using non EDTA tubes then were stored in cooling box. Serum was gotten from centrifuges on 2060 × g for 15 min at 5°C. Lipid plasma was analyzed using hemodialysis method with standardization value of analyzer (Caretium Biochemistry Analyzer). Standard value for cholesterol was 138 mg/dl that ranged 96-179 mg/dl, triglyceride was 104 mg/dl that ranged 73-135mg/dl, and HDL was 79 mg/dl that ranged 55-102 mg/dl. Milk yield was collected every days in the morning milking at 04.00 a.m. and evening milking at 03.00 p.m. Milk fat was analyzed by lactoscan machine every 7 days of periods.

#### C. Calculations

The LDL was calculated by Friedewald equation which was known with cholesterol total level, HDL level and triglycerides level.  $LDL = \text{Cholesterol Total} - (\text{HDL} + \text{Triglycerides}/5)$

### III. STATISTICAL ANALYSIS

The mean values and standard errors of the mean (SEM) were calculated for all of the data. All of the data were analyzed with One Simple T-Test procedures in SPSS ver.21. Statistical significance was declared at  $P \leq 0.05$ , and a tendency to significance was declared at  $0.05 < P \leq 0.20$ .

TABLE II: THE INGREDIENT AND NUTRIENT COMPOSITION OF DIETS

Composition	Dietary Treatments			
	TMR diet		RPC diet	
	Napier grass	Concentrate	Napier grass	Concentrate
Ratio	40	60	40	60
DM <sup>a</sup> (%)	6.12	53.03	6.12	53.03
TDN (%)	22.93	41.72	22.93	41.72
CP <sup>d</sup> (%)	3.20	10.02	3.20	10.02
CF <sup>a</sup> (%)	13.22	11.95	13.22	11.95
EE <sup>a</sup> (%)	1.08	2.73	1.08	2.73
RPC (g/head/d)	0		30	

## IV. RESULT

#### A. Dry matter intake

The average of DMI was not affected significantly by RPC diet. DMI of TMR diet was  $13.50 \pm 2.18$  kg/head/d and  $2.91 \pm 0.42\%$  BW. The DMI of RPC diet was  $14.19 \pm 1.98$  kg/head/d and  $3.09 \pm 0.35\%$  BW (showed in Table III.). The DMI of two treatments were ranging between 2.25 – 4.32% BW. The sufficiency of DMI is used for performance and milk production. Based on National Research Council [12] that the DMI of dairy cows ranged 2.25 – 4.32% BW. The averages DMI showed that RPC diet tended to increase  $0.69 \pm 0.20$  kg/head/d ( $P = 0.2$ ) and  $0.18 \pm 0.07\%$  BW ( $P = 0.2$ ) than TMR diet. The role play of choline in increasing utilization efficiency of energy tended to cause the increasing energy metabolism. DMI was not affected by metabolism rate but was affected by type of feed and digestibility. Choline plays an important role in the regulation of carbohydrate conversion into CO<sub>2</sub> and ATP [13]. The liver glycogen concentration increased linearly while increasing intake of choline chloride that to assist the formation of glycogen from hydrolysis to be used as a source of glucose in the liver become faster that affected to increasing milk yield [4,14]. Addition 40 g/head/d RPC increased 0.3 kg/d DMI (Davidson, 2008) the addition 25 g/head/d RPC (50% choline chloride) has been significantly increased DMI of 2.2 kg/head/d and milk yield of 4.4 kg/d [6].

#### B. Ether extract intake

The average of EEI was not affected by RPC diet. EEI of TMR diet averaged  $512 \pm 83$  g/head/d and  $0.11 \pm 0.18\%$  BW. EEI of RPC diet averaged  $543 \pm 81$  g/head/d and  $0.12 \pm 0.17\%$  BW (has been shown in Table III.). It showed that the EEI with RPC diet increased numerically  $31.25 \pm 0.2$  g/head/d ( $P = 0.19$ ) and  $0.01 \pm 0.01\%$  BW ( $P = 0.18$ ). EEI has not affected significantly due to DMI which not significantly affected. The EEI affected increasing lipid metabolism rapidly so transfer of fatty acid into mammary gland will increase. Choline which provides a methyl group donor in the synthesis of carnitine is an essential component for the oxidation of fatty acids so that the mobilization of non esterified fatty acids (NEFA) from feed in blood to the liver increases and does not accumulate in the liver. EEI can increase cholesterol and LDL levels in blood [18]. Feed intake is a factor that determine of function, respond and nutrient feed using [2].

TABEL III: EFFECTS OF CHOLINE CHLORIDE TO DRY MATTER INTAKES AND ETHER EXTRACT INTAKES

Parameters	Treatments	
	TMR diet	RPC diet
DMI (kg/head/d)	13,50 ± 2,18	14,19 ± 1,98
DMI (% BW)	2,91 ± 0,42	3,09 ± 0,35
EEI (g/head/d)	512,38 ± 82,23	543,63 ± 79,81
EEI (% BW)	0,11 ± 0,18	0,12 ± 0,17

DMI = Dry Matter Intake; EEI = Ether Extract Intake.

TABEL IV: EFFECT OF CHOLINE CHLORIDE TO PLASMA LIPID

Parameters	Treatments	
	TMR diet	RPC diet
Cholesterol level (mg/dl)	159,37 ± 21,02	161,43 ± 21,52
Trygliceride (mg/dl)	70,14 ± 6,76	69,16 ± 3,53
LDL level (mg/dl)	74,77 ± 16,94	80,21 ± 24,67
HDL level (mg/dl)	70,52 ± 11,76	67,48 ± 15,62

LDL = Low density lipoprotein; HDL = High density lipoproteins.

TABEL V: EFFECT OF CHOLINE CHLORIDE TO MILK YIELD AND MILK FAT

Parameters	Treatments	
	TMR diet	RPC diet
Milk yield (kg/d)	13,84 ± 2,41 <sup>a</sup>	14,29 ± 2,90 <sup>b</sup>
Milk yield 4% FCM (kg/d)	13,36 ± 2,63 <sup>a</sup>	14,53 ± 3,12 <sup>b</sup>
Milk fat (%)	3,76 ± 0,27 <sup>a</sup>	4,11 ± 0,28 <sup>b</sup>

### C. Cholesterol

Cholesterol level in blood has not been affected by treatments. Cholesterol level of TMR diet averaged 159.37 ± 21 mg/dl and cholesterol level of RPC diet averaged 161.43 ± 22 mg/dl (as showed in Table IV.). Cholesterol level was in the normal value that ranged between 139-177 mg/dl [21]. Cholesterol level is influenced by the consumption of DMI and EEI that provides a source of energy for the precursor of cholesterol, named Acetyl CoA which is produced from glucose and fatty acids in the mitochondria and amino acid catabolism [10]. Cholesterol level with RPC diet was higher 1.06 ± 0.1 mg/dl (P = 0.6) in normal range than TMR diet although it was not affected significantly. The role play of choline was to improve supply acetyl-CoA which the main precursor for cholesterol synthesis and as a phospholipid (phosphatidylcholine) in lipoproteins that used in cholesterol transport through the blood flow. Cholesterol acts as the main compounds such as bile steroid body whose function as an emulsifier of fat so that the body can absorb it. Based on cholesterol was the main of all steroid compounds which had been synthesized in the body, one function of bile salts to emulsify fats into fatty acids that can be absorbed by the body [11]. Choline that formed phosphatidylcholine contained in bile function in maintaining the hydrophobic protective layer on the mucosa of the small intestine. Cholesterol was formed in the liver and used as the source of dietary fat which was absorbed by the small intestine of chylomicrons form. Fat absorption which occurs actively for triglycerides, cholesterol and phospholipids is formed in the gut, then the flow of blood to further join the protein (apoprotein) forming a lipoprotein that can circulate in the blood circulation [1].

### D. Triglycerides

The average of trygliceride level was not affected by traetments. Triglyceride of TMR diet averaged 70.14 ± 6.76 mg/dl and triglyceride of RPC diet averaged 69.16 ± 3.53 mg/dl (showed in Table IV.). Triglyceride with RPC diet decreased 0.98 ± 3.23 md/dl (P = 0.77) than TMR diet although it was not affected significantly. Triglyceride level of two treatments were in the normal value that ranged 63 -77 mg/dl [20]. Choline forms phospatidylcholine of lipoprotein as phospholipids form so that can help the mobilization and β oksidation in liver which made trygliceride level in the blood lowest because more of lipoprotein produced. Addition choline chloride in feed increased NEFA mobilization to make taking up fatty acids by liver for oxidazing into ketone bodies and carbondioxide or trygliceride increased [4]. Lipoprotein contained of trygliceride, fatty acids, and fosfolipids so that lipoprotein was produced more, then made the more higher of trygliceride which was brought by lipoprotein [5].

### E. Low density lipoprotein

The average of LDL level were not affected significantly (showed in Table IV.) and averaged 74.77 ± 12 mg/dl of TMR diet and 80.21 ± 16 mg/dl of RPC diet. LDL level with RPC diet was higher 5.44 ± 0.4 mg/dl (P = 0.35) than TMR diet including in the range of normal value eventhough it was not affected significantly. It indicated cause by increasing fat metabolism and role of choline as phosphatidylcholine that improve secretion of NEFA and triglycerides in VLDL then it will became to be LDL. LDL is a lipoprotein which has a lower density, serves a major carrier of essential fatty acids from the liver to the tissue for cells absorption and metabolism [19]. Normal levels of LDL in dairy cows ranged between 60 - 130 mg/dl and contained of triglycerides 10% and cholesterol

50% [21]. Addition 45 g/head/d of RPC (25% choline chloride) increased the de novo synthesis which was shown by the present of highly milk fatty acids [23]. Highly milk fatty acids due to the influence of choline caused by high level fatty acids in blood from being transported through VLDL. The addition of RPC 75 g/head/d (contained 37.5% choline chloride) increased the synthesis and secretion NEFA through VLDL from liver [14]. Methionine is supplied by choline that provides the amino acid profile of VLDL and as phosphatidylcholine (main phospholipids) surface layer of VLDL that plays a role in the lipid transport [8].

#### F. High density lipoprotein

The average of HDL level in blood were not affected significantly by treatments (has been shown in Table IV.). HDL level of TMR diet averaged  $70.52 \pm 17$  mg/dl and HDL level of RPC diet averaged  $67.48 \pm 25$  mg/dl. HDL level of two treatments was in the normal range of dairy cow's HDL level which was 40-90 mg/dl. HDL is a lipoprotein with low cholesterol levels and higher protein levels and it function plays in cholesterol transport from tissues to the liver. HDL is known as a molecule composed of fat (13% of cholesterol and triglycerides less than 5%) and 50% protein [21]. HDL level with RPC diet was lower  $3.04 \pm 0.8$  mg/dl ( $P = 0.37$ ) than TMR diet including in the normal value eventhough not affected significantly. The function of choline as phosphatidylcholine in bonding lipoprotein HDL mechanism has not been optimal because choline was more usable for chatabolism in order to produce energy than anabolism to form HDL. High of EEI caused fatty acid in blood higher so that made LDL level higher than HDL level. The nutrients from the food begins to be absorbed by the small intestine and then carried by the blood stream to the liver. HDL plays an important role in the mobilization of NEFA from the feed to be brought to the liver in lipoprotein formed, HDL. Choline is a precursor of phosphatidylcholine, a major phospholipid in high-density lipoprotein or HDL [24]. The function of lipoproteins is to transport cholesterol from peripheral tissues and from other lipoproteins to be oxidized in the liver so that HDL plays a role in to prevent of atherosclerosis in blood vessels [22].

#### G. Milk yield

Milk yield was significant affected by treatments (has been shown in Table 3.) Milk yield of TMR diet averaged  $13.84 \pm 2.41$  kg/d and the milk yield of RPC diet averaged  $14.29 \pm 2.90$  kg/d ( $P = 0.05$ ). The averages of milk yield in 4% FCM (Fat corrected milk) conversion of TMR diet was  $13.36 \pm 2.63$  kg/day and RPC diet was  $14.53 \pm 3.12$  kg/day ( $P = 0.04$ ) (Shown in Table V.). It showed that increasing milk yield was made by choline that used for energy metabolism especially in chatabolism to produce energy for milk yield. Choline contributes in increasing energy efficiency and play as methyl group donor in energy metabolism, especially in lipid metabolism. Choline improved chatabolism of fatty acids and amino acids to produce Asetil CoA so it was obtained not only from glicolysis but also gluconeogenesis. Deposit of fatty acids increased while increasing choline chloride addition in certain level, it made gluconeogenesis better that improved glycogen from glycolysis to use for glucose source [3]. Asetyl

CoA that produced by glycolysis and gluconeogenesis used to produced energy which used for milk yield. Glycogen increased while increasing of RPC consumption in dairy cow's liver. The slightly lower triglyceride concentrations for cows that received increasing amounts of RPC may have allowed for greater rates of gluconeogenesis, sparing glycogen from hydrolysis for use as a glucose source or perhaps replenishing glycogen in liver more quickly than cows not receiving RPC [14]. Addition RPC 12 g/head/d increased milk yield 3.6 kg [9], addition RPC 45 g/head/d which consists of 20 g choline chloride) in feed increased milk yield 2.9 kg [16] and addition RPC 30 g/head/d which consists of 37.5% choline chloride increased milk yield 0.3 kg than without addition choline chloride [23].

#### H. Milk fat

The average of milk fat was significant affected by RPC diet (showed in Table V.) Milk fat of TMR diet averaged  $3.76 \pm 0.27\%$  and RPC diet averaged  $4.11 \pm 0.28\%$ . RPC diet increased significantly  $0.35\% \pm 0.01\%$  of milk fat ( $P = 0.03$ ). The averages of milk fat include in normal value of dairy cow's milk fat that ranged 3.24-4.49%. Milk fat increased because lipid metabolism in liver increased. LDL which was precursor of milk fat synthesis was higher with RPC diet than TMR diet. Mobilization of NEFA to take fatty acids by liver for oxidize into ketone and carbondioxide or tryglyseride increased [4]. Addition RPC increased cholesterol level 30% so lipid storage and lipid metabolism in balance by liver reached. Choline play role as methyl group donor in carnitine synthesis those use for lipid oxidation and lipotropic effect (Forms NEFA into tryglyserides and secretion tyglycerides) [17]. Milk fat increased because LDL which was as a precursor of milk fat tented to increase so lipid was carried into mammary gland quite a lot. Choline as phosphatidylcholine in lipoprotein bond made lipoprotein was more produced so that lipid esterification and tryglycerides from liver increased then carried as LDL into mammary gland. It implicated to increase mechanism in mammary gland such as milk yield and milk fat. Choline also play role as fosfolipid component in fat globule of milk fat so fluidity and rigidity from fat globule of milk depended by its [8].

## IV. CONCLUSION

Choline Chloride as Rumen Protected Choline could be added in the diets of lactation cows to increase milk yield and milk fat.

## ACKNOWLEDGMENT

Authors acknowledge to Faculty of Animal and Agricultural Sciences, Diponegoro University, for facility of study and World Class University of Diponegoro University, Indonesia, for funding the paper.

## REFERENCES

- [1] S. Agustina, "Utilization of shrimp shell chitosan as reduce blood cholesterol levels in rat Sprague dawley," M.S. thesis, Dept. Applied Chem. Stud., Udayana Univ., Denpasar, Indonesia, 2015.
- [2] A. Astuti. (June 2009). The effect of high quality feed supplement addition on the nutrient consumption and digestibility of early lactating dairy cow. *Buletin Peternakan*. [Online]. 33(2). pp. 81-87. Available:

- <http://journal.ugm.ac.id/index.php/buletinpeternakan/article/download/120/476>
- [3] H. M. Awwad, S. H. Kirsch, J. Geisel, and R. Obeid. (February 2014). Measurement of concentrations of whole blood levels of choline, betaine, and dimethylglycine and their relations to plasma levels. *Journal of Chromatogr B*. [online]. 975 (2014). pp. 41-45. Available: [www.elsevier.com/locate/chromb](http://www.elsevier.com/locate/chromb)  
<https://doi.org/10.1016/j.jchromb.2014.02.030>
- [4] A. Baldi and L. Pinotti, "Choline metabolism in high-producing dairy cows: metabolic and nutritional basis," in *the Canadian Society of Animal Science Symposium. 2005* .pp. 207-212, vol. 86, USA : Canadian Journal of Anim. Sci., 2006.  
<https://doi.org/10.4141/a05-061>
- [5] W. W. Christie, *Lipid Metabolism in Ruminant Animals*, 1st ed. Headington Hill Hall, U.K.: Oxford, 1981, ch. 2, pp. 48-50.
- [6] Y. H. Chung, N. E. Brown, C. M. Martinez, T. W. Cassidy, and G. A. Varga. (February 2009). Effects of rumen protected choline and dry propylene glycol on feed intake and blood parameters for Holstein dairy cows in early lactation. *J. Dairy Sci.* [Online]. 92(6). pp. 2729-2736. Available: [http://www.journalofdairyscience.org/article/S0022-0302\(09\)70590-9/fulltext?mobileUi=0](http://www.journalofdairyscience.org/article/S0022-0302(09)70590-9/fulltext?mobileUi=0)
- [7] S. Davidson, B. Hopkins, J. Odle, C. Brownie, V. Fellner, and L. Whitlow. (2008). Supplementing limited methionine diets with rumen-protected methionine, betaine, and choline in early lactation Holstein cows. *J. Dairy Sci* [Online]. 91(4). pp.1552-1559. Available: [http://agri.lu.ac.ir/uploads/PIIS0022030208712839\\_3751.pdf](http://agri.lu.ac.ir/uploads/PIIS0022030208712839_3751.pdf)  
<https://doi.org/10.3168/jds.2007-0721>
- [8] V. M. A. Etcheverry, "Absorbtion and utilization of choline and vitamin b12 in lactating dairy cows using different delivery methods," Ph.D. dissertation, Dept. Agric., Tennessee Univ., Knoxville, USA, 2014.
- [9] J. R. Hartwell, M. J. Cecava, and S. S. Donkin. (October 2000). Impact of dietary rumen undegradable protein and rumen-protected choline on intake, peripartum liver triacylglyceride, plasma metabolites, and milk production in transition dairy cows. *J. Dairy Sci.* [Online]. 83(12). pp. 2907-2917. Available: <http://www.sciencedirect.com/science/article/pii/S0022030200751915>
- [10] D. B. Marks, A. D. Marks, and C. M. Smith. *Basic medical biochemistry: a clinical approach*, 1st ed. Lippincot Williams and Wilkins, US: Philadelphia, 1996.
- [11] R. K. Murray, D. K. Granner, P. A. Mayes, and V. W. Rodwell, *Harper Biochemistr*. 25th ed. Stamford, CT: Appleton and Lange; 2003.
- [12] National Research Council, *Nutrient Requirements of Dairy Cattle Seventh Revised Edition*, Rev. ed. National Academy Press, USA: Washington D.C.
- [13] M. Parakkasi, *Ruminant Nutrition and Feed Science*, Indonesia University Press, Indonesia : Jakarta, 1999.
- [14] M. Piepenbrink and T. Overton. (November 2003). Liver metabolism and production of cows fed increasing amounts of rumen-protected choline during the periparturient period. *J. Dairy Sci.* [Online]. 86(5). Pp. 1722-1733. Available: <https://www.ncbi.nlm.nih.gov/pubmed/12778583>
- [15] L. Pinotti, A. Baldi, and V. Dell'Orto. (October 2002). Comparative memmalian choline metabolism with emphasis on the high-yielding dairy cow. *Nut. Reserc. Rev.*[Online]. 047(15). pp.315-331.  
<https://doi.org/10.1079/NRR200247>
- [16] L. Pinotti, A. Baldi, I. Politis, R. Rebucci, L. Sangalli, and V. Dell'Orto. (2003). Rumen-protected choline administration to transition cows: effects on milk production and vitamin e status. *J. Veterinary Med. series A*. [Online]. 50(1). pp.18-21.  
<https://doi.org/10.1046/j.1439-0442.2003.00502.x>
- [17] L. Pinotti, A. Campagnoli, L. Sangalli, R. Rebucci, V. Dell'Orto, and A. Baldi. (2004). Metabolism of periparturient dairy cows fed rumen-protected choline. *J. Anim. Feed Sci.* [Online]. 13(1). pp.551-554.
- [18] K. Sejrsen, T. Hvelplund, and M. O. Nielsen. *Ruminant Physiology*. Wageningen Academic Publishers, UK: Netherland, 2008.
- [19] R. Soegih. (1995). Gangguan Penyakit Jantung dan Pembuluh Darah. Sadar Pangan dan Gizi. *Bulletin*. [Online]. 4(3). pp.2-3.
- [20] M. M., Tighe and M. Brown. *Comprehensive Review for Veterinary Tehnicians*. Mosby Elsevier, USA: St. Louis, 2008.
- [21] D. Weatherby and S. Ferguson. *Blood Chemistry and CBC Analysis-Clinical Laboratory Testing from a Functional Perspective*. Bear Mountain Publishing, USA: Ashland, 2002.
- [22] Widiyanto, M. Soejono, H. Hartadi, dan Z. Bachrudin. (2007). Effect of protected kapok seed oil supplementation on in vitro ruminal lipid status. *Anim. Prod.* [Online]. 11(2). pp.122-128
- [23] G. Xu, J. An Ye, L. Jianxin and Y. Yu. (March 2005). Effect of rumen-protected choline addition on milk performance and blood metabolic parameters in transition dairy cows. *J. Anim. Sci.* [Online]. 19(1). pp.390-395.
- [24] S. H. Zeisel and K. A. da Costa. (2009). Choline: An Essensial Nutrient for Public Health. *Nutrition Review*. [Online] 67(11). pp.615-623.  
<https://doi.org/10.1111/j.1753-4887.2009.00246.x>