

Effects of Replacing Concentrate with Soybean Curd Residue Silage on Ruminal Characteristics, Plasma Leucine and Glucose Turnover Rates in Sheep

by D. W Harjanti

Submission date: 13-Jan-2020 10:05AM (UTC+0700)

Submission ID: 1241272597

File name: rnal_of_Animal_Sciences_Dian_W_Harjanti_2012_JASA-24-361-374.pdf (791.38K)

Word count: 7299

Character count: 36776

**Effects of Replacing Concentrate with Soybean Curd
Residue Silage on Ruminal Characteristics, Plasma Leucine
and Glucose Turnover Rates in Sheep**

Harjanti D. W., Sugawara Y., Al-Mamun M. and Sano H.

J Anim Sci Adv 2012, 2(4): 361-374



Online version is available on: www.grjournals.com

Effects of Replacing Concentrate with Soybean Curd Residue Silage on Ruminal Characteristics, Plasma Leucine and Glucose Turnover Rates in Sheep

Harjanti D. W., Sugawara Y., Al-Mamun M. and *Sano H.

Department of Animal Sciences, Faculty of Agriculture, Iwate University, Morioka 020-8550, Japan

Abstract

The effects of replacing concentrate with soybean curd residue silage on the plasma leucine (Leu) and glucose turnover rates (LeuTR and GluTR) of sheep were evaluated using an isotope dilution method of [1-¹³C]Leu and [U-¹³C]glucose. The ruminal characteristics and nitrogen (N) balance of the sheep were also determined. The sheep were fed two diets in a crossover experiment for 21 days. The first diet consisted of 80% mixed hay and 20% commercial concentrate (CON diet), and the second diet consisted of 80% mixed hay and 20% soybean curd residue silage (SCRS diet), on a dry matter basis for both diets. The primed continuous infusion method of [1-¹³C]Leu and [U-¹³C]glucose was performed on day 21. The N absorption, N retention, and N digestibility did not differ between the diets. Furthermore, the ruminal total volatile fatty acid concentration did not differ, but the molar concentration of propionate was significantly higher ($P=0.02$) in the sheep fed the SCRS diet than the CON diet. The plasma LeuTR and GluTR were comparable between the diets. The results showed that soybean curd residue silage could be comparable with commercial concentrate on ruminal fermentation as well as on the plasma Leu and glucose kinetics. Thus, soybean curd residue silage could be used to replace commercial concentrate in the diet of sheep.

Key words: food by-product, glucose kinetic, isotope dilution, protein kinetic, wether

*Corresponding Author: Department of Animal Sciences, Faculty of Agriculture, Iwate University, Ueda 3-18-8, Morioka 020-8550, Japan; sano@iwate-u.ac.jp

Received on: 26 Feb 2012

Revised on: 13 Mar 2012

Accepted on: 25 Mar 2012

Online Published on: 14 Apr 2012

Introduction

Economic and environmental concerns prompt investigation into the possibility of using the organic resources generated by the food industry as alternative animal feeds. Soybean curd residue (SCR), a by-product of tofu, soy milk and soy protein manufacturing, is high in protein content; however, because SCR is considered an industrial waste, it is often incinerated due to its rapid rate of deterioration. Ensiling SCR (soybean curd residue silage; SCRS) could prolong its shelf life and increase its nutrient composition and digestibility (Cho *et al.*, 2007). An *in vitro* study on the nutritional properties of SCRS showed that the fermentation process also enhanced antioxidant activity (Rashad *et al.*, 2001), and previous studies (Oltjen & Putnam, 1966; Clifford & Tillman, 1968; Prior, 1976; Knaus *et al.*, 2002) have reported increased nitrogen (N) absorption and N retention when soy protein-based diets were fed to steers and sheep. Inclusion of SCRS in the diet of sheep has been shown to increase the glucogenic propionate concentration in the rumen without any negative effects on ruminal fermentation (Xu *et al.*, 2001). Nevertheless, research regarding the use of SCRS as ruminant feed on the N balance and the kinetics of plasma amino acid (AA) and glucose has been limited. Thus, additional studies should be needed to promote the utilization of this by-product as ruminant feed.

We hypothesized that SCRS can replace commercial concentrate in the diet of sheep without any adverse effects on whole-body N and glucose metabolism due to its high soluble N and fermentable carbohydrate contents. Improved utilization of residue generated by the food industry such as SCRS could reduce feed cost and environmental problem. The objective of this study was to determine the effects of replacing a commercial concentrate with SCRS on the plasma leucine (Leu) and glucose turnover rates (LeuTR and GluTR), using an isotope-dilution method with [$1\text{-}^{13}\text{C}$]Leu and [$\text{U-}^{13}\text{C}$]glucose. The N balance, ruminal characteristics and blood metabolites were also analyzed.

47

Materials and Methods

Animals and diets

All of the animal handling, including blood sampling, was conducted according to the guidelines established by the Animal Care Committee of Iwate University. Four crossbred (Suffolk x Corriedale) wether, all of approximately 4 years of age and initially of 54 ± 4 kg of body weight were used. This experiment used a crossover design: a 21-day period that consisted of 14 days of dietary adaptation and 7 days of sample collection. Throughout the adaptation period, the sheep were kept in individual pens and fed two dietary treatments. The first diet consisted of 80% mixed hay (orchardgrass and reed canarygrass) and 20% commercial concentrate (CON diet), and the second diet consisted of 80% mixed hay and 20% soybean curd residue silage (SCRS diet), on a dry matter basis. The SCRS contained 15% beet pulp. The chemical composition of mixed hay, SCRS and commercial concentrate are listed in Table 1. The diets were estimated to be isonitrogenous and the metabolizable energy intakes assumed from the National Research Council (1985) were slightly above the maintenance level for both diets, as shown in Table 2. The sheep were fed twice daily at 08.30 and 20.30 hours and had *ad libitum* access to water. The SCRS was purchased from a food company (Hirakawa Food Co. Ltd., Iwate, Japan), placed into 90 L polyvinyl bags, compacted, and sealed tightly. After 15 days of ensiling, the container was opened and the silage was fed to the animals as reported previously (Amaha *et al.*, 1995).

On day 15, the sheep were moved to individual metabolic cages in a controlled-environment room, with an air temperature of 23°C, a relative humidity of 70%, and lighting from 08.00 to 22.00 hours.

Sample collection

Before starting the feeding experiment, samples of the feed were collected and dried in a forced-air oven at 60°C for 48 hours to determine the chemical composition and to formulate the dietary treatments. Offered feed and refused feed (if any) were sampled and dried using the same procedure for the chemical analysis.

From day 16 to day 20, urine and feces samples were collected at 24-hour intervals using a

3 mm plastic screen separator. The samples were collected once daily at 10.00 hours. The urine was collected from each sheep in a bucket containing 50 mL of 6 N H₂SO₄, and the volume was recorded; 50 mL of the urine was stored at -30°C

until analysis. Feces samples were dried in a forced-oven at 60°C for 48 hours and were then passed through a 15 mm mesh and stored at room temperature until analysis.

Table 1: Chemical compositions of the mixed hay, soybean curd residue silage and commercial concentrate

	Mixed hay	Soybean curd residue silage ^a	Commercial concentrate ^b
DM (g/kg)	910	309	880
CP (g/kg DM)	118	183	127
EE (g/kg DM)	30	110	38
NDF (g/kg DM)	680	150	207
ME (Mcal/kg DM)	1.97	3.32 ^c	2.68

^aSoybean curd residue silage contained 15% beet pulp (Hirakawa Food Co. Ltd., Japan)

^bCommercial concentrate: formula feed ("α-Beef" made by Chubu Feed Co. Ltd., Japan)

^cCalculated as the proportion of soybean curd residue and beet pulp (85:15) in the silage according to AFF (1995).

DM: dry matter

CP: crude protein

EE: ether extract

NDF: neutral detergent fiber

ME: metabolizable energy

Table 2: Diet formulation and intakes of crude protein (CP) and metabolizable energy (ME) of the dietary treatments

	Treatment ^a	
	CON diet	SCRS diet
Mixed hay (g · kg BW ^{-0.75} · d ⁻¹)	52.6	48.5
Commercial concentrate ^b (g · kg BW ^{-0.75} · d ⁻¹)	13.2	0
Soybean curd residue silage (g · kg BW ^{-0.75} · d ⁻¹)	0	12.1
CP intake (g · kg BW ^{-0.75} · d ⁻¹)	7.9	7.9
ME intake (Mcal · kg BW ^{-0.75} · d ⁻¹)	139	135

^aTreatment: mixed hay (orchardgrass and reed canarygrass) as the basal diet which is supplemented with either commercial concentrate (CON diet) or soybean curd residue silage (SCRS diet) at a ratio of 80:20 on dry matter basis.

Ruminal fluid (50 mL) was collected on day 20 at 0 (before feeding), 3 and 6 hours after the morning feeding via a stomach tube. The pH of the ruminal fluid was immediately measured using a pH meter (HM-10P, Toa Electronics Ltd., Japan) after collecting each sample. The liquid fraction the ruminal fluid was separated by centrifuging at 8,000 x g for 10 minutes at 4°C. A 5 mL aliquot of the liquid fraction was used to measure the ruminal VFA content, and 1 mL of the liquid fraction was mixed with 1 mL of a 0.1 mol/L HCl for the

ruminal ammonia N (NH₃-N) analysis. All of the samples were stored at -30°C until further analysis.

Measurements of amino acid and glucose kinetics

On day 21 of each dietary treatment, a primed-continuous infusion [1-¹³C]Leu and [U-¹³C]glucose method was conducted to determine the plasma Leu and glucose kinetics simultaneously over a period of 4 hours, between 3 and 7 hours after the morning feeding. Polyvinyl catheters were placed in both jugular veins in the morning and filled with

a sterile solution of 3.8% (w/v) trisodium citrate. A mixed saline solution of 7.2 $\mu\text{mol/kg BW}^{0.75}$ [^{13}C]Leu (L-leucine $1\text{-}^{13}\text{C}$, 99 atom% excess ^{13}C ; Cambridge Isotope Laboratories, USA) and 3.0 $\mu\text{mol/kg BW}^{0.75}$ [^{13}C]glucose (D-glucose- $^{13}\text{C}_6$, 99 atom% excess ^{13}C ; Cambridge Isotope Laboratories, USA) was injected into the right jugular catheter as a priming dose. This solution was then continuously infused through the same catheter using a multichannel peristaltic pump (AC-2120, Atto, Japan) at rates of 7.2 and 3.0 $\mu\text{mol} \cdot \text{kg BW}^{-0.75} \cdot \text{h}^{-1}$ for [^{13}C]Leu and [^{13}C]glucose, respectively. The infusion rate of the tracer solution was recorded every 30 minutes throughout the infusion period. Blood samples (6 mL) were collected from the left jugular catheter immediately before the priming injection and at 30-minute intervals during the last 2 hours of isotope infusion (5 to 7 hours after feeding). The blood samples were placed in heparinized tubes and stored in ice. The plasma from the blood sample was separated by centrifugation at 8,000 $\times g$ for 10 minutes at 4°C and then stored at -30°C until further analysis.

Sample analysis

The chemical components of the diets were analyzed according to the Association of Official Analytical Chemists (AOAC, 1990). The N contents in the diets, feces and urine were analyzed using the Kjeldahl method (Kjeltec 2100, Foss, Sweden). The fermentation products of the silage were analyzed using cold-water extracts (Cai *et al.*, 1999). The pH values of the silage and ruminal fluid were measured using a pH meter (HM-10P, Toa Electronics Ltd., Japan). The lactic acid concentration of the silage was measured by the colorimetric method (Taylor, 1996), and the absorbance was measured using a spectrophotometer (U-1000, Hitachi, Japan) at 570 nm. The molar VFA concentrations of the silage and ruminal fluid were determined using gas chromatography (HP-5890, Hewlett Packard, USA) after steam distillation. The volatile basic N (VBN) content of the silage was determined based on the steam distillation method described by Dhaouadi *et al.* (2007). To assess the quality of silage, we calculated the Flieg point from the DM and pH values (Yilmaz *et al.*, 2009) and the V-

score from the VBN/total N and VFA concentrations (Takahashi *et al.*, 2005).

The concentrations of plasma Leu and α -ketoisocaproic acid (α -KIC) and enrichments of the plasma [^{13}C]Leu and α -[^{13}C]KIC were determined by gas chromatography mass spectrometry (QP-2010, Shimadzu, Japan) with selected ion monitoring according to the procedures of Rocchiccioli *et al.* (1981) and Alder & Smith (1988). The plasma [^{13}C]glucose enrichment was measured using the methods of Tserng & Kalhan (1983) with slight modifications described by Fujita *et al.* (2006). The plasma glucose concentration was measured by the glucose oxidase method (Huggett & Nixon, 1957). The plasma free AA and urea concentrations were determined using an automatic AA analyzer (JLC-500/V, JEOL, Japan). The plasma non-esterified fatty acid (NEFA) concentrations were determined enzymatically using a commercial diagnostic kit (NEFA-C test, Wako, Japan).

Calculations

The turnover rates of plasma Leu and glucose (LeuTR and GluTR) were calculated according to the equation described by Wolfe (1984):

$$\text{TR} (\text{mmol} \cdot \text{kg BW}^{-0.75} \cdot \text{h}^{-1}) = I \times (1/E - 1)$$

where I represents the infusion rates of [^{13}C]Leu and [^{13}C]glucose and E represents the isotope enrichments of the plasma α -[^{13}C]KIC, metabolite of [^{13}C]Leu, and plasma [^{13}C]glucose at the steady state.

Statistical Analysis

The mean values and standard errors of the mean (SEM) were calculated for all of the data. All of the data were analyzed using the MIXED procedure in SAS (1996). The fixed effects in the model were the period, diet, and the period \times diet interaction, and the random effect was the sheep. The results were considered significant for $P < 0.05$, and the tendency was defined as $0.05 \leq P < 0.10$. Repeated statements were used for the time course of the ruminal characteristics, and the difference of the least square means using Tukey's adjustment was used ($P < 0.05$).

Results

Silage quality and balance

The fermentative characteristics of the SCRS are shown in Table 3. The pH value was 4.14, and organic acids, such as lactic acid, acetate, propionate and butyrate, were detected. The effects of replacing commercial concentrate with SCRS on the N balance are presented in Table 4. Although both diets were estimated to be isonitrogenous, the

N intake determined at the end of the experiment tended ($P=0.05$) to be higher for the SCRS diet than the CON diet. The fecal N excretion did not differ ($P=0.17$), but the urinary N excretion tended to be higher ($P=0.05$) for sheep fed the SCRS diet than those fed the CON diet. Both groups of sheep retained similar amounts of N ($P=0.42$), and the N absorption and digestibility did not differ ($P=0.11$ and $P=0.14$, respectively) between the two diets.

Table 3: Fermentative characteristics of the soybean curd residue silage

Item	Amount
Moisture (g/kg)	690
pH	4.14
Lactic acid (g/kg FM ^a)	40
Acetate (g/kg FM)	1.1
Propionate (g/kg FM)	0.03
Butyrate (g/kg FM)	0.003
VBN ^b (g/kg total N)	68
Flieg point	100
V-score	98

^aFM: fresh matter

^bVBN: volatile basic nitrogen

Table 4: Effects of feeding soybean curd residue silage on the nitrogen (N) balance in sheep^a

Items	Treatment ^c		SEM ^b	P value
	CON diet	SCRS diet		
N intake (g · kg BW ^{-0.75} · d ⁻¹)	1.267	1.278	0.005	0.05
Fecal N (g · kg BW ^{-0.75} · d ⁻¹)	0.48	0.45	0.03	0.17
Urinary N (g · kg BW ^{-0.75} · d ⁻¹)	0.48	0.55	0.07	0.05
N absorption (g · kg BW ^{-0.75} · d ⁻¹)	0.79	0.83	0.01	0.11
N retention (g · kg BW ^{-0.75} · d ⁻¹)	0.31	0.28	0.02	0.42
N digestibility (%)	62.5	65.0	1.0	0.14

^aValues represent the means for n = 4.

^bSEM: standard error of the mean.

^cTreatment: mixed hay (orchardgrass and reed canarygrass) as the basal diet which is supplemented with either commercial concentrate (CON diet) or soybean curd residue silage (SCRS diet), at a ratio of 80:20 on dry matter basis.

Ruminal characteristics

The observed changes in the ruminal characteristics during the sampling period are provided in Figure 1. The effects of the two diets on the ruminal pH, NH₃-N and VFA concentrations, represented as the mean value of four sheep in three sampling periods (0, 3, and 6 hours after feeding), are shown in Table 5. No

interaction was observed between the dietary treatment and the sampling time for the ruminal characteristic ($P>0.10$). The sheep that consumed the SCRS tended ($P=0.05$) to have a higher ruminal pH than those fed the CON diet. The pH values under both treatments were above 6.8 prior to feeding, and the values dropped to approximately 6.7 at 3 hours after feeding.

EFFECTS OF REPLACING CONCENTRATE WITH SOYBEAN CURD RESIDUE SILAGE ON RUMINAL ...

57 Table 5: Effects of feeding soybean curd residue silage on the ruminal characteristics of sheep^a

Item	Treatment ^c		SEM ^b	P value	
	CON diet	SCRS diet		Treatment	Time
pH	6.69	6.87	0.03	0.05	0.008
NH ₃ -N (mmol/L)	11.2	11.9	0.5	0.22	0.002
Total VFA (mmol/L)	87.9	93.4	1.5	0.13	0.002
Individual VFA concentrations (mmol/L)					
Acetate	61.7	64.9	0.4	0.26	0.007
Propionate	15.9	19.2	0.8	0.02	<0.001
Isobutyrate	0.7	0.8	0.1	0.05	0.01
Butyrate	8.3	6.8	0.4	0.02	0.07
56 Valerate	0.8	1.1	0.1	0.06	0.007
Valerate	0.5	0.6	0.04	0.02	0.001
Acetate:propionate ratio	3.9	3.4	0.2	0.03	0.008

^aValues represent the means for n = 4.

^bSEM: standard error of the mean.

^cTreatment: mixed hay (orchardgrass and reed canarygrass) as the basal diet which is supplemented with either commercial concentrate (CON diet) or soybean curd residue silage (SCRS diet), at a ratio of 80:20 on dry matter basis.

NH₃-N: ammonia nitrogen; VFA: volatile fatty acid

Comparable NH₃-N concentrations were found for both groups ($P=0.22$); the NH₃-N concentrations were increased at 3 hours after feeding and eventually returned to the previous values under both dietary treatments.

The total VFA concentration in the rumen did not differ significantly ($P=0.13$) between the two diets. The acetate concentrations were similar for the two diets, but the propionate concentration was higher ($P=0.02$) in the sheep fed the SCRS than the CON diet, resulting in a lower acetate:propionate ratio ($P=0.03$) for the SCRS than CON diet. The butyrate concentration was lower ($P=0.02$) for the SCRS diet compared to the CON diet, whereas the valerate concentration was higher ($P=0.02$) and

other branched-chain VFAs (isobutyrate and isovalerate) concentrations tended to be higher ($P=0.05$ and $P=0.05$, respectively) under the SCRS treatment. The total VFA concentration and the concentrations of acetate, propionate, butyrate, and valerate were increased at 3 hours after feeding and eventually returned toward previous values. The isobutyrate and isovalerate concentrations were gradually decreased at 6 hours after feeding.

Pre-isotope infusion plasma concentration of blood metabolites

The plasma AA, urea and NEFA concentrations determined at the pre-isotope infusion period at 3 hours after feeding are presented in Table 6.

Table 6: Effects of feeding soybean curd residue silage on the plasma amino acid, urea and non-esterified fatty acid concentrations during pre-isotope infusion in sheep^a

Item	Treatment ^c		SEM ^b	P value
	CON diet	SCRS diet		
Essential AA (μmol/L)				
Threonine	232	235	13	0.83
Valine	290	259	13	0.14
Methionine	36	34	1.3	0.07
Isoleucine	105	89	19	0.02
Leucine	158	130	7	0.03
Phenylalanine	69	60	2	0.03
Histidine	65	57	2	0.05
Lysine	118	100	7	0.13
Non-essential AA (μmol/L)				
Serine	151	131	16	0.34
Asparagine	58	48	5	0.19
Glutamic acid	52	58	4	0.28
Glutamine	390	336	54	0.11
Glycine	584	523	40	0.27
Alanine	197	173	15	0.25
Tyrosine	98	86	3	0.04
Arginine	133	118	4	0.07
Proline	129	107	8	0.09
Urea, mmol/L	6.2	7.8	0.2	0.02
NEFA, mEq/L	0.10	0.13	0.03	0.38

^aValues represent the means for n = 4.

^bSEM: standard error of the mean.

^cTreatment: mixed hay (orchardgrass and reed canarygrass) as the basal diet which is supplemented with either commercial concentrate (CON diet) or soybean curd residue silage (SCRS diet), at a ratio of 80:20 on dry matter basis.

AA: amino acid

NEFA: non-esterified fatty acid

The plasma isoleucine, Leu, phenylalanine, and tyrosine concentrations were lower ($P<0.05$)

and the plasma methionine, histidine, arginine, and proline concentrations tended to be lower ($P<0.10$)

in the sheep fed the SCRS than CON diet; the concentrations of other AA were comparable between the diets. The plasma urea concentration was higher ($P=0.02$) in the sheep fed the SCRS than CON diet, and the plasma NEFA concentration did not differ ($P=0.38$) between the diets.

Amino acid and glucose kinetics

The plasma Leu and α -KIC concentrations and plasma α -[1- 13 C]KIC enrichment were in a steady state during the last 2 hours of isotope infusion

(Figure 2). The plasma Leu and α -KIC concentrations did not differ ($P=0.18$ and $P=0.13$, respectively) between the diets (Table 7). The sheep fed the SCRS had similar ($P=0.74$) plasma LeuTR with those of sheep fed the CON diet. The plasma glucose concentration and [U- 13 C]glucose enrichment were essentially constant during the last 2 hours of the isotope infusion (Figure 7). The plasma glucose concentration and GluTR did not differ ($P=0.68$ and $P=0.27$, respectively) between the two diets.

Table 7: Effects of feeding soybean curd residue silage on the plasma leucine (Leu) and glucose kinetics in sheep^a

Item	Treatment ^c		SEM ^b	P value
	CON diet	SCRS diet		
Leu concentration ($\mu\text{mol/L}$)	135	107	13	0.18
α -KIC concentration ($\mu\text{mol/L}$)	12.8	10.3	0.5	0.13
LeuTR ($\text{mmol} \cdot \text{kg BW}^{-0.75} \cdot \text{h}^{-1}$)	0.52	0.51	0.05	0.74
Glucose concentration (mmol/L)	3.34	3.40	0.18	0.68
GluTR ($\text{mmol} \cdot \text{kg BW}^{-0.75} \cdot \text{h}^{-1}$)	2.24	2.14	0.09	0.27

^aValues represent the means for $n = 4$.

^bSEM: standard error of the mean.

^cTreatment: mixed hay (orchardgrass and reed canarygrass) as the basal diet which is supplemented with either commercial concentrate (CON diet) or soybean curd residue silage (SCRS diet) at a ratio of 80:20 on dry matter basis.

α -KIC: α -ketoisocaproic acid

LeuTR: leucine turnover rate

GluTR: glucose turnover rate

Discussion

Soybean curd residue silage quality

The SCRS was well preserved, as indicated by the low pH, high lactic acid content, high Flieg point and V-score. The lactic acid concentration in our SCRS was similar to that found by Amaha *et al.* (1996), but it was higher than that found by Xu *et al.* (2001). During the fermentation process, some hemicellulose and cellulose were broken down into various simple sugars and smaller end products, such as lactic acid, acetate and butyrate (Cao *et al.*, 2009). The NDF content in our SCRS (150 g/kg DM) was comparable to the NDF content reported from raw SCR (145 g/kg DM) by O'Toole (1999).

Nitrogen balance

Previous studies (Oltjen & Putnam, 1966; Clifford & Tillman, 1968; Prior, 1976; Knaus *et*

al., 2002) have shown that inclusion soy protein in the diet of ruminants enhances the N intake, urinary N excretion, N retention and N absorption. Sano *et al.* (2009) studied the effect of different N sources supplementation (urea and soybean meal) in the isonitrogenous design and found a large urinary N excretion difference between treatments, indicating a change in N retention. In the present study, replacing the commercial concentrate with SCRS did not affect the amount of N absorbed and retained, whereas it did slightly affect the urinary N excretion. Hence, our findings demonstrate the feasibility of replacing commercial concentrate with soybean curd residue silage to maintain a positive N balance.

Ruminal characteristics

The ruminal pH values of both groups of sheep were above 6.5 for all of the sampling times, and this is the optimal pH for normal ruminal fermentation.




Fig. 1: Time-course changes of ruminal characteristics in sheep fed the SCRS diet (■) and the CON diet (Δ). SCRS diet, mixed hay supplemented with soybean curd residue silage; CON diet, mixed hay supplemented with 53 commercial concentrate. A ratio of 80:20 (dry matter basis) was used for both diets. Ruminal fluid samples were collected at 0 (before feeding), 3 and 6 hours after morning feeding via a stomach tube. Values are expressed as mean ± SEM for n = 4. Different letters (a, b, c) indicate significant different (P<0.05) between time after feeding within each treatment.

Thus, replacing the commercial concentrate with the SCRS did not impair ruminal fermentation; a similar response observed by Xu *et al.* (2001). The SCRS and CON diets in the present study were approximately isonitrogenous and isoenergetic; thus, similar N compounds should be available for $\text{NH}_3\text{-N}$ production in the rumen. This finding is consistent with the results of Ipharraguerre *et al.* (2005) who reported that the




Fig. 2: Time course changes of plasma leucine (Leu) and α -ketoisocaproic acid (α -KIC) concentrations, plasma $[1\text{-}^{13}\text{C}]\text{Leu}$ and $\alpha\text{-}[1\text{-}^{13}\text{C}]\text{KIC}$ enrichments during the last 2 hours of isotope infusion in sheep fed the SCRS diet (■) and the CON diet (Δ). SCRS diet, mixed hay supplemented with soybean curd residue silage; CON diet, mixed hay supplemented with commercial concentrate. A ratio of 80:20 (dry matter basis) was used for both diets. Values are expressed as mean \pm SEM for $n = 4$.

concentration of $\text{NH}_3\text{-N}$ in the rumen fluid only affected by the level of dietary CP intakes.




Fig. 2: Time course changes of plasma leucine (Leu) and α -ketoisocaproic acid (α -KIC) concentrations, plasma $[1\text{-}^{13}\text{C}]\text{Leu}$ and $\alpha\text{-}[1\text{-}^{13}\text{C}]\text{KIC}$ enrichments during the last 2 hours of isotope infusion in sheep fed the SCRS diet (■) and the CON diet (Δ). SCRS diet, mixed hay supplemented with soybean curd residue silage; CON diet, mixed hay supplemented with commercial concentrate. A ratio of 80:20 (dry matter basis) was used for both diets. Values are expressed as mean \pm SEM for $n = 4$.

16 Volatile fatty acids are produced in the rumen as the end product of microbial fermentation, and their production largely depends on the type of carbohydrate ingested (Dijkstra, 1994). The large increment in the ruminal propionate concentration in the sheep fed the SCRS diet could be attributed to the highly soluble carbohydrate and starch contents (Yang, 2005). Although the NDF concentration in a ruminant diet is positively correlated with the amount of VFA produced in the rumen (Beauchemin, 1993), the lower NDF content for the SCRS (150 g/kg DM) compared to the commercial concentrate (207 g/kg DM) did not alter the total VFA concentration in the rumen. Replacing the commercial concentrate with the SCRS did not significantly influence the ruminal acetate concentration, indicating similar efficiencies of energy-source production for both diets.

Pre-isotope infusion plasma concentration of blood metabolites

The primary source of dietary protein (Schelling *et al.*, 1967) affects the plasma AA concentrations in sheep. For the different protein sources in the present study, the lower concentrations of certain plasma AAs in the sheep fed the SCRS diet might not be indicative of the total AA supply because the protein supply in both treatments was similar and sufficient. Thus, the observed lower AA levels were most likely due to the limiting AAs in the diet (Lapierre & Lobley, 2001), which, in soybean-based diets, are sulfur-containing AAs (methionine and cysteine) (Ma *et al.*, 1996). Moreover, Lapierre and Lobley (2001) suggested that the concentration of plasma AA after feeding varied depend on the rate of digestion and AA catabolism of the dietary protein source. Furthermore, these authors noted that approximately 46% of AA from a highly degradable type of protein such as silage did not appear in the peripheral vein, whereas for a concentrate the amount varied between 23% and 24%. Although the sheep fed the SCRS diet in our study exhibited lower concentrations of certain plasma AAs, replacing the commercial concentrate with the SCRS did not induce an AA imbalance,

based on the positive N balance and similar N retention for both of the diets.

The use of different protein sources affected the plasma urea concentration, which is a metabolic end product of protein catabolism or the product of soluble protein fermentation in the rumen. Although the ruminal NH₃-N concentrations within the sampling periods (0, 3, and 6 hours after feeding) did not differ significantly between the two diets, the high plasma urea concentration for the SCRS diet, reflected by the rapid degradation of N compounds and the rapid absorption of NH₃-N through the rumen, resulted in an increased level of plasma urea. Furthermore, the N excretion via the urine tended to increase over time. Similarly, Leibholz and Cook (1967) showed that elevated plasma urea concentrations increased the N excretion rate via the urine and the rate of return of N to the rumen as urea in the saliva in sheep fed alfalfa hay and concentrate.

The circulating plasma NEFA concentration was markedly related to the nutritional status. The plasma NEFA concentration increases with high rates of lipolysis, such as occurs with fasting, negative energy balance or stress (Trenkle & Kuhlemeier, 1966). Hence, the comparable plasma NEFA concentrations indicated comparable nutritional status between sheep fed the SCRS and CON diet.

Amino acid and glucose kinetics

In the present study, both plasma [1-¹³C]Leu and α-[1-¹³C]KIC enrichments were determined, but only plasma α-[1-¹³C]KIC enrichment is discussed because it represents an accurate index of the intra- and extra-cellular amino acid exchange (Magni *et al.*, 1994). The plasma LeuTR observed in our study were comparable to previous studies using single-isotope infusion of [1-¹³C]Leu to determine protein kinetics in sheep (Sano *et al.*, 2004; 2009); similarly, the plasma GluTR observed in our study were comparable to those from previous study on glucose kinetic in sheep determined using single-isotope infusion of [U-¹³C] glucose (Sano *et al.*, 2000). These results indicate that the double isotope used in our experiment did not alter either the plasma Leu or

glucose kinetics of the sheep. The use of different protein sources resulted in varying plasma AA concentrations, which could influence the turnover rate of proteins (Nielsen *et al.* 1993). Although reports on the AA kinetics of soybean-based diets in ruminants are limited, the results from other studies in pigs (Deutz *et al.*, 1998) and humans (Bos *et al.*, 2003) showed that the AAs from soybean-based diets are rapidly digested and absorbed, indicating an enhanced AA catabolism. The low concentration of plasma AAs in the sheep fed the SCRS diet could have been caused by the utilization of AAs as an energy substrate to support protein turnover within the tissues; thus, the plasma LeuTR measured were comparable between the two diets. In addition, the dietary treatments were approximately isoenergetic and isonitrogenous, which could have contributed to the comparable plasma LeuTR for the two diets because the plasma LeuTR was positively correlated with the dietary protein and metabolizable energy intake in sheep (Liu *et al.*, 1995) and cows (Lapierre *et al.*, 2002). Similarly, Sano *et al.* (2009) reported comparable plasma LeuTR for different protein sources (urea and soybean meal) in sheep fed the isoenergetic and isonitrogenous diets.

The GluTR is influenced by several factors, including the energy intake and supply of gluconeogenic substrate to the liver (Ortiqes-Marty *et al.*, 2003). The propionate produced in the rumen is a major glucogenic substrate and precursor of de novo glucose synthesis (Herbein *et al.*, 1978). In the present study, although the glucogenic propionate concentration in the rumen was higher for the SCRS than CON diet, the plasma GluTR were similar for both of the diets. In accordance with this result, an increasing supply of glucogenic propionate did not systematically increase plasma GluTR in steers (Seal & Parker, 1994) and sheep (Linington *et al.*, 1998). Herbein *et al.* (1978) and Seal *et al.* (1992) pointed out that similar energy intake resulted in similar plasma GluTR in sheep and growing steers. Thus, the isoenergetic diets in the present study did not significantly influence the plasma GluTR. The comparable results for the plasma Leu and glucose kinetics indicated similar nutritive value for the two diets. Similar plasma Leu and glucose kinetic

responses could be achieved by replacing commercial concentrate with SCRS in the same amount of metabolizable energy intake.

In conclusion, replacing commercial concentrate with SCRS in the diet of sheep did not impair ruminal fermentation, and enhanced the glucogenic propionate concentration. The inclusion of SCRS in the diets of sheep could ensure a positive N balance and resulted in similar plasma AA and glucose kinetics with commercial concentrate when formulated in the same energy intake. Thus, feed cost can be reduced by replacing commercial concentrate with soybean curd residue. Furthermore, technological developments in the use of residue and by-products will help to improve the feed self-sufficiency rates and make animal agriculture more sustainable.

Acknowledgements

The authors are grateful to Hirakawa Food Co, Ltd. for providing the soybean curd residue for the experiments.

References

- AOAC. 1990: *Official Methods of Analysis*. 15th ed. Assoc. Off. Anal. Chem., Arlington, VA.
- Amaha K, Sasaki Y, Segawa T (1996). Utilization of tofu (soybean curd) by-product as feed for cattle. *Food and Fertilizer Tech.* Accessed May 27, 2010. [http://www.agnet.org/library/eb/419\[1\].pdf](http://www.agnet.org/library/eb/419[1].pdf).
- Beauchemin KA (1991). Effects of dietary neutral detergent fiber concentration and alfalfa hay quality on chewing, rumen function, and milk production of dairy cows. *J. Dairy Sci.*, 74: 3140-3151.
- Bequette BJ, Metcalf JA, Wray-Cahen D, Backwell FRC, Sutton JD, Lomax MA, MacRae JC, Lobley GE (1996). Leucine and protein metabolism in the lactating dairy cow mammary gland: responses to supplemental dietary crude protein intake. *J. Dairy Res.*, 63: 209-222.
- Bergen WG, Henneman HA, Magee WT (1973). Effect of dietary protein level and protein source on plasma and tissue free amino acids in growing sheep. *J. Nutr.*, 103: 575-585.
- Bos C, Metges CC, Gaudichon C, Petzke KJ, Pueyo ME, Morens, C, Everwand J, Benamouzig R, Tome D (2003). Postprandial kinetics of dietary amino acids are the main determinant of their metabolism after soy or milk protein ingestion in humans. *J. Nutr.*, 133: 1308-1315.
- Cai Y, Benno Y, Ogawa M, Kumai S (1999). Effect of applying lactic acid bacteria isolated from forage crops on fermentation characteristics and aerobic deterioration

- of silage. *J. Dairy Sci.*, 82: 520-526.
- Calder AG, Smith A (1988). Stable isotope ratio analysis of leucine and ketoisocaproic acid in blood plasma by gas chromatography/mass spectrometry. Use of tertiary butyldimethylsilyl derivatives. *Rapid Comm. Mass Spectrom.*, 2: 4-16.
- Cao Y, Takahashi T, Horiguchi K (2009). Effects of addition of food by-products on the fermentation quality of a total mixed ration with whole crop rice and its digestibility, preference, and rumen fermentation in sheep. *Anim. Feed Sci. Technol.*, 151: 1-11.
- Cho JH, Min BJ, Chen YJ, Yoo JS, Wang Q, Kim JD, Kim IH (2007). Evaluation of FSP (fermented soy protein) to replace soybean meal in weaned pigs: growth performance, blood urea nitrogen and total protein concentrations in serum and nutrient digestibility. *Asian-Aust. J. Anim. Sci.*, 20: 1874-1879.
- Clifford AJ, Tillman AD (1968). Urea and isolated soybean protein in sheep purified diets. *J. Anim. Sci.*, 27: 484-489.
- Deutz NEP, Bruins MJ, Soeters PB (1998). Infusion of soy and casein protein meals affects interorgan amino acid metabolism and urea kinetics differentially in pigs. *J. Nutr.*, 128: 2435-2445.
- Dhaouadi A, Monser L, Sadok S, Adhoum N (2007). Validation of flow-injection-gas diffusion method for total volatile basic nitrogen determination in seafood products. *J. Food Chem.*, 103: 1049-1053.
- Dijkstra J (1994). Production and absorption of volatile fatty acids in the rumen. *Livest. Prod. Sci.*, 39: 61-69.
- Fujita T, Kajita M, Sano H (2006). Responses of whole body protein synthesis, nitrogen retention and glucose kinetics to supplemental starch in goats. *Comp. Biochem. Physiol.*, 144B:180-187.
- Given DI, Rulquin H (2004) Utilisation by ruminants of nitrogen compounds in silage-based diets. *Anim. Feed Sci. Technol.*, 114: 1-18.
- Harris PM, Skene PA, Buchan V, Milne E, Calder AG, Anderson SE, Cornell A, Lobley GE (1992). Effect of food intake on hind-limb and whole body protein metabolism in young growing sheep: chronic studies based on arterio-venous techniques. *Br. J. Nutr.*, 68: 389-407.
- Herbein JM, Van Maaren RW, McGilliard AD, Young JW (1978). Rumen propionate and blood glucose kinetics in growing cattle fed isoenergetic diets. *J. Nutr.*, 108: 994-1001.
- Huggett AG, Nixon DA (1957). Enzymatic determination of blood glucose. *J. Biochem.*, 66: 12.
- Ipharraguerre IR, Clark JH, Freeman DE (2005). Varying protein and starch in the diet of dairy cows. 1. Effects on ruminal fermentation and intestinal supply of nutrients. *J. Dairy Sci.*, 88: 2537-2555.
- Knaus WF, Beermann DH, Tedeschi LO, Czajkowski M, Fox DG, Russel JB (2002). Effects of urea, isolated soybean protein and blood meal on growing steers fed a corn-based diet. *Anim. Feed Sci. Technol.*, 102: 3-14.
- Lapierre H, Lobley GE (2001). Nitrogen recycling in the ruminant: A Review. *J. Dairy Sci.*, 84: E223-E236.
- Lapierre H, Blouin JP, Bernier JF, Reynolds CK, Dubreuil P, Lobley GE (2002). Effect of supply of metabolizable protein on whole body and splanchnic leucine metabolism in lactating cows. *J. Dairy Sci.*, 85: 2631-2641.
- Leibholz J, Cook CF (1967). Free amino acids, ammonia and urea concentrations in the blood plasma of starved lambs. *J. Nutr.*, 93: 561-567.
- Linington MJ, Meyer JHF, van der Walt JG (1998). Ruminant VFA production rates, whole body metabolite kinetics and blood hormone concentrations in sheep fed high and low fibre diets. *S. Afr. J. Anim. Sci.*, 28: 83-98.
- Liu SM, Lobley GE, MacLeod NA, Kyle DJ, Chen XB, Ørskov ER (1995). Effects of long-term protein excess or deficiency on whole body protein turnover in sheep nourished by intragastric infusion of nutrients. *Br. J. Nutr.*, 73: 829-839.
- Ma CY, Liu WS, Kwok KC, Kwok F (1997). Isolation and characterization of protein from soymilk residue (okara). *Food Res. Int.*, 29: 799-805.
- MAFF. Ministry of Agriculture, Forestry and Fisheries Research Council (1995). Standard tables of feed composition in Japan. Central association of livestock industry. Japan.
- Magni F, Arnoldi L, Galati G, Kienle M (1994). Simultaneous determination of plasma levels of α -ketoisocaproic acid and leucine and evaluation of α -[1-¹³C]ketoisocaproic acid and [1-¹³C]leucine enrichment by gas chromatography-mass-spectrometry. *Anal. Biochem.*, 220: 308-331.
- Mlay PS, Pereka AE, Weisbjerg MR, Hvelplund T, Madsen J (2003). Digestion and passage kinetics of fibre in mature dairy heifers maintained on poor quality hay as affected by the source and level of nitrogen supplementation. *Anim. Feed Sci. Technol.*, 109: 19-33.
- Nielsen K, Kondrup J, Elsner P, Juul A, Jensen ES (1994). Casein and soybean protein have different effects on whole body protein turnover at the same nitrogen balance. *Br. J. Nutr.*, 72: 69-81.
- NRC (1985). Nutrient Requirements of Sheep, 6th revised edition. National Academy Press, Washington, DC.
- Ortigue-Marty I, Vernet J, Majdoub L (2003). Whole body glucose turnover in growing and non-productive adults ruminants: meta-analysis and review. *Reprod. Nutr. Dev.*, 43: 371-383.
- Oltjen RR, Putnam PA (1966). Plasma amino acid and nitrogen retention by steers fed purified diets containing urea or isolated soy protein. *J. Nutr.*, 89: 385-390.
- O'Toole D (1999). Review: Characteristics and use of okara, the soybean residue from soy milk production. *J. Agric. Food Chem.*, 47: 363-371.
- Prior RL (1976). Effects of dietary soy or urea nitrogen and feeding frequency on nitrogen metabolism, glucose metabolism and urinary metabolite excretion in sheep. *J. Anim. Sci.*, 42: 160-167.

EFFECTS OF REPLACING CONCENTRATE WITH SOYBEAN CURD RESIDUE SILAGE ON RUMINAL ...

- Rocchiccioli F, Leroux JP, Cartier P (1981). Quantitation of 2-ketoacids in biological fluids by gas chromatography chemical ionization mass spectrometry of *o*-trimethylsilyl-quinoxalinol derivatives. *Biomed. Mass Spectrom.*, 8: 160-164.
- Sano H, Konno S, Shiga A (2000). Chromium supplementation does not influence glucose metabolism or insulin action in response to cold exposure in mature sheep. *J. Anim. Sci.*, 78: 2950-2956.
- Sano H, Kajita M, Fujita T (2004). Effect of dietary protein intake on plasma leucine flux, protein synthesis, and degradation in sheep. *Comp. Biochem. Physiol.*, 139B: 163-164.
- Sano H, Shibasaki S, Sawada H (2009). The effect of the source of nitrogen supplementation on nitrogen balance, rates of plasma leucine turnover, protein synthesis and degradation in sheep. *Arch. Anim. Nutr.*, 63: 401-412.
- SAS. 1996. SAS/STAT® software: changes and enhancement through Release 6.11. SAS Institute Inc. Cary, NC, USA.
- Schelling GT, Hinds FC, Hatfield EE (1967). Effect of dietary protein levels, amino acid supplementation and nitrogen source upon the plasma free amino acid concentrations in growing lambs. *J. Nutr.*, 92: 339-347.
- Seal CJ, Parker DS, Avery PJ (1992). The effect of forage and forage-concentrate diets on rumen fermentation and metabolism of nutrients by mesenteric and portal-drained viscera in growing steers. *Br. J. Nutr.*, 67: 335-370.
- Seal CJ, Parker DS (1994). Effect of intraruminal propionic infusion on metabolism of mesenteric and portal drained-viscera in growing steers fed a forage diet: I. Volatile fatty acids, glucose and lactate. *J. Anim. Sci.*, 72: 1325-1334.
- Takahashi T, Horiguchi K, Goto M (2005). Effect of crushing unhulled rice and the addition of fermented juice epiphytic lactic bacteria on the fermentation quality of whole crop rice silage, and its digestibility and rumen fermentation status in sheep. *Anim. Sci. J.*, 76: 353-358.
- Taylor KACC (1996). A simple colorimetric assay for muramic acid and lactic acid. *Appl. Biochem. Biotechnol.*, 56: 49-58.
- Trenkle A, Kuhlemeier KV (1996). Relationship of rumen volatile acids, blood glucose, and plasma nonesterified fatty acids in sheep. *J. Anim. Sci.*, 25: 111-1115.
- Tserng KY, Kalhan SC (1983). Estimation of glucose carbon recycling and glucose turnover with [U-¹³C]glucose. *Am. J. Physiol.*, 245: E476-E482.
- Wolfe RR (1984). Tracers in Metabolic Research. Radioisotope and stable isotope/mass spectrometry methods. Alan R. Liss, Inc, New York.
- Xu CC, Suzuki H, Toyokawa K (2001). Characteristics of ruminal fermentation of wethers fed tofu cake silage with ethanol. *Anim. Sci. J.*, 72: 299-305.
- Yang CMJ (2005). Soybean milk residue ensiled with peanut hulls: fermentation acids, cell wall composition, and silage utilization by mixed ruminal microorganisms. *Biores. Technol.*, 96: 1419-1424.
- Yilmaz A, Altinok S, Kocabas Z (2009). An investigation on quality parameters of the silages made by corn and soybean grown in different seeding rates. *J. Anim. Vet. Adv.*, 8: 1856-1859.

Effects of Replacing Concentrate with Soybean Curd Residue Silage on Ruminal Characteristics, Plasma Leucine and Glucose Turnover Rates in Sheep

ORIGINALITY REPORT

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

11%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** archimer.ifremer.fr 1%

Internet Source
- 2** Jeong, Jinsuk, and In Ho Kim. "Comparative efficacy of up to 50% partial fish meal replacement with fermented soybean meal or enzymatically prepared soybean meal on growth performance, nutrient digestibility and fecal microflora in weaned pigs : Fermented Soybean Meal in Pigs", *Animal Science Journal*, 2015. 1%

Publication
- 3** iwate-u.repo.nii.ac.jp 1%

Internet Source
- 4** Gaafar, Khalid (Universität Leipzig, Veterinärmedizinische Fakultät). "Effect of fat in comparison to starch in an isoenergetic diet on the metabolism of high yielding dairy cows", *Universitätsbibliothek Leipzig*, 2004. 1%

Publication

5	<p>Mohammad AL-MAMUN. "Effect of ensiled hop (Humulus lupulus L.) residues on plasma acetate turnover rate in sheep : ACETATE METABOLISM IN SHEEP", Animal Science Journal, 06/2011</p> <p>Publication</p>	1%
6	<p>Tadahisa Fujita, Takahiro Itoh, Hiroya Majima, Hiroaki Sano. "Influence of Dietary Salinomycin on Feeding-induced Variations of Glucose Kinetics and Blood Volatile Fatty Acids and Insulin Concentrations in Sheep Fed a High-roughage Diet", Asian-Australasian Journal of Animal Sciences, 2007</p> <p>Publication</p>	<1%
7	<p>arno.unimaas.nl</p> <p>Internet Source</p>	<1%
8	<p>hal.archives-ouvertes.fr</p> <p>Internet Source</p>	<1%
9	<p>shareok.org</p> <p>Internet Source</p>	<1%
10	<p>mro.massey.ac.nz</p> <p>Internet Source</p>	<1%
11	<p>pt.slideshare.net</p> <p>Internet Source</p>	<1%
12	<p>M. Al-Mamun, Y. Sako, H. Sano. "Effect of cold</p>	<1%

exposure on plasma glucose and acetate
turnover rates in sheep", Animal Production
Science, 2014

Publication

13

Md. Mostafizar Rahman, Paramintra
Vinitchaikul, Arvinda Panthee, Xue Bi, Hiroaki
Sano. "Effect of feeding whole-crop corn silage
as dietary roughage on physiological and
digestive response of sheep under heat
exposure", Animal Production Science, 2017

Publication

<1%

14

e-sciencecentral.org

Internet Source

<1%

15

academic.oup.com

Internet Source

<1%

16

pt.scribd.com

Internet Source

<1%

17

www.lib.ncsu.edu

Internet Source

<1%

18

ajas.info

Internet Source

<1%

19

edepot.wur.nl

Internet Source

<1%

20

Kaori Yasuda, Masayuki Kitagawa, Kazato
Oishi, Hiroyuki Hirooka, Takemi Tamura, Hajime

<1%

Kumagai. "Growth performance, carcass traits, physiochemical characteristics and intramuscular fatty acid composition of finishing Japanese black steers fed soybean curd residue and soy sauce cake", Animal Science Journal, 2015

Publication

21 www.tdx.cat <1 %
Internet Source

22 www.scribd.com <1 %
Internet Source

23 Hoang Lam Dang, Renlong Lv, Taketo Obitsu, Toshihisa Sugino. "Effect of replacing alfalfa hay with a mixture of cassava foliage silage and sweet potato vine silage on ruminal and intestinal digestion in sheep", Animal Science Journal, 2018 <1 %
Publication

24 tsukuba.repo.nii.ac.jp <1 %
Internet Source

25 www.fibrogenesis.com <1 %
Internet Source

26 www.uady.mx <1 %
Internet Source

27 sasas.co.za <1 %
Internet Source

28	adsa.asas.org Internet Source	<1%
29	"Rumenology", Springer Nature, 2016 Publication	<1%
30	Kashif ur Rehman, Abdul Rehman, Minmin Cai, Longyu Zheng et al. "Conversion of mixtures of dairy manure and soybean curd residue by black soldier fly larvae (<i>Hermetia illucens</i> L.)", <i>Journal of Cleaner Production</i> , 2017 Publication	<1%
31	libres.uncg.edu Internet Source	<1%
32	ar.scribd.com Internet Source	<1%
33	"Sheep Production Adapting to Climate Change", Springer Nature, 2017 Publication	<1%
34	hss.ulb.uni-bonn.de Internet Source	<1%
35	Kiran Doranalli, Timothy Mutsvangwa. "Feeding sunflower oil to partially defaunate the rumen increases nitrogen retention, urea-nitrogen recycling to the gastrointestinal tract and the anabolic use of recycled urea-nitrogen in growing lambs", <i>British Journal of Nutrition</i> ,	<1%

2011

Publication

36

Tadahisa Fujita, Masahiro Kajita, Hiroaki Sano, Akio Shiga. "Effects of Non-protein Energy Intake on the Concentrations of Plasma Metabolites and Insulin, and Tissue Responsiveness and Sensitivity to Insulin in Goats", Asian-Australasian Journal of Animal Sciences, 2006

Publication

<1%

37

www.biochemj.org

Internet Source

<1%

38

www.ijpab.com

Internet Source

<1%

39

Sano, H.. "Effect of Mild Cold Exposure on Whole-Body and Net Hindquarters Glucose Metabolism in Sheep", Comparative Biochemistry and Physiology -- Part B: Biochemistry and Molecular Biology, 199701

Publication

<1%

40

jn.nutrition.org

Internet Source

<1%

41

R. García-Martínez, M. J. Ranilla, M. L. Tejido, M. D. Carro. " Effects of disodium fumarate on rumen microbial growth, methane production and fermentation of diets differing in their

<1%

forage:concentrate ratio ", British Journal of Nutrition, 2007

Publication

42

H. SANO, M. KAJITA, M. ITO, T. FUJITA, A. TAKAHASHI. "Effect of metabolizable protein intake on rates of plasma leucine turnover and protein synthesis in heifers", The Journal of Agricultural Science, 2007

Publication

<1%

43

repository.up.ac.za

Internet Source

<1%

44

B. J. Min, J. H. Cho, Y. J. Chen, H. J. Kim, J. S. Yoo, C. Y. Lee, B. C. Park, J. H. Lee, I. H. Kim. "Effects of Fermented Soy Protein on Growth Performance and Blood Protein Contents in Nursery Pigs", Asian-Australasian Journal of Animal Sciences, 2009

Publication

<1%

45

uhra.herts.ac.uk

Internet Source

<1%

46

cgspace.cgiar.org

Internet Source

<1%

47

www.tandfonline.com

Internet Source

<1%

48

macau.uni-kiel.de

Internet Source

<1%

49 Bampidis, V.A.. "Citrus by-products as ruminant feeds: A review", *Animal Feed Science and Technology*, 20060628 $<1\%$

50 Savary-Auzeloux. "Effect of intake on whole body plasma amino acid kinetics in sheep", *Reproduction Nutrition Development*, 2003 $<1\%$

51 Chuncheng XU. "The effect of replacing brewers' grains with barley tea grounds in total mixed ration silage on feed intake, digestibility and ruminal fermentation in wethers", *Animal Science Journal*, 10/2008 $<1\%$

52 B. E. A. Borhami. "Nitrogen Utilization in Sheep when Feeding Either Sodium Hydroxide Treated Straw Plus Urea or Ammonia Treated Straw", *Acta Agriculturae Scandinavica*, 1983 $<1\%$

53 Jiachen Fang, Guangjun Xia, Yang Cao. "Effects of replacing commercial material with apple pomace on the fermentation quality of total mixed ration silage and its digestibility, nitrogen balance and rumen fermentation in wethers", *Grassland Science*, 2019 $<1\%$

54 Y. Jiang, P.-F. Zhao, S.-M. Lin, R.-J. Tang, Y.-J.

Chen, L. Luo. " Partial substitution of soybean meal with fermented soybean residue in diets for juvenile largemouth bass, ", Aquaculture Nutrition, 2018

Publication

<1%

55

Correddu, F., A. Nudda, G. Battacone, R. Boe, A.H.D. Francesconi, and G. Pulina. "Effects of grape seed supplementation, alone or associated with linseed, on ruminal metabolism in Sarda dairy sheep", Animal Feed Science and Technology, 2015.

Publication

<1%

56

Tao Ma, Dandan Chen, Yan Tu, Naifeng Zhang, Bingwen Si, Kaidong Deng, Qiyu Diao. "Effect of supplementation of allicin on methanogenesis and ruminal microbial flora in Dorper crossbred ewes", Journal of Animal Science and Biotechnology, 2016

Publication

<1%

57

Lei Chen, Xianjun Yuan, Junfeng Li, Zhihao Dong, Tao Shao. "Effects of applying oil-extracted microalgae on the fermentation quality, feed-nutritive value and aerobic stability of ensiled sweet sorghum", Journal of the Science of Food and Agriculture, 2018

Publication

<1%

58

www.science.gov

Internet Source

<1%

59

www.journalofanimalscience.org

Internet Source

<1%

60

www.ncbi.nlm.nih.gov

Internet Source

<1%

61

R. L. Ludwick, J. P. Fontenot, R. E. Tucker. "Studies of the Adaptation Phenomenon by Lambs Fed Urea as the Sole Nitrogen Source, Chemical Alterations in Ruminal and Blood Parameters", Journal of Animal Science, 1972

Publication

<1%

62

Thongsuk Jetana, Cheerapath Vongpipatana, Sirima Thongruay, Sungworn Usawang, Sunpeth Sophon. "Apparent Digestibility, Nitrogen Balance, Ruminal Microbial Nitrogen Production and Blood Metabolites in Thai Brahman Cattle Fed a Basal Diet of Rice Straw and Supplemented with Some Tropical Protein-rich Trees", Asian-Australasian Journal of Animal Sciences, 2010

Publication

<1%

63

C. S. Ali, A. N. Sorensen. "Effect of Level of Urea-N on its Utilization in Alkali-treated Wheat Straw in Pelleted Rations by Sheep", Acta Agriculturae Scandinavica, 2009

Publication

<1%

64

link.springer.com

Internet Source

<1%

65

www.animalsciencepublications.org

Internet Source

<1%

66

Liufeng Zheng, Fangrui Zuo, Shengjun Zhao, Pingli He, Hongkui Wei, Quanhong Xiang, Jiaman Pang, Jian Peng. "Dietary supplementation of branched-chain amino acids increases muscle net amino acid fluxes through elevating their substrate availability and intramuscular catabolism in young pigs", *British Journal of Nutrition*, 2017

Publication

<1%

67

Saman Lashkari, Mahmood Habibian, Søren Krogh Jensen. "A Review on the Role of Chromium Supplementation in Ruminant Nutrition—Effects on Productive Performance, Blood Metabolites, Antioxidant Status, and Immunocompetence", *Biological Trace Element Research*, 2018

Publication

<1%

68

Tadahisa Fujita. "Combined effect of salinomycin and feeding on whole body glucose kinetics in sheep fed a high-concentrate diet", *Reproduction Nutrition Development*, 09/2006

Publication

<1%

Arvinda Panthee, Ayana Matsuno, Mohammad

69

Al-Mamun, Hiroaki Sano. "Effect of feeding garlic leaves on rumen fermentation, methane emission, plasma glucose kinetics, and nitrogen utilization in sheep", Journal of Animal Science and Technology, 2017

Publication

<1%

70

Cao, Yang, Yanqing Zang, Renlong Lv, Toshiyoshi Takahashi, Norio Yoshida, and Huanmin Yang. "Effects of adding urea on fermentation quality of pruned persimmon branch silage and its digestibility, preference, nitrogen balance and rumen fermentation in beef cattle : Persimmon Branches and Feeding Value", Animal Science Journal, 2014.

Publication

<1%

71

"Theatre Presentations", Advances in Animal Biosciences, 2011

Publication

<1%

72

Tatsuya Morita. "Rice Protein Isolates Produced by the Two Different Methods Lower Serum Cholesterol Concentration in Rats Compared with Casein", Journal of the Science of Food and Agriculture, 08/1996

Publication

<1%

73

Lu, Qi, Jian Wu, Min Wang, Chuanshe Zhou, Xuefeng Han, Edwin Nicholas Odongo, Zhiliang Tan, and Shaoxun Tang. "Effects of dietary

<1%

addition of cellulase and a *Saccharomyces cerevisiae* fermentation product on nutrient digestibility, rumen fermentation and enteric methane emissions in growing goats", *Archives of Animal Nutrition*, 2016.

Publication

74

Thongsuk JETANA. "The comparative study digestion and metabolism of nitrogen and purine derivatives in male, Thai, Swamp buffalo and Thai, Brahman cattle", *Animal Science Journal*, 04/2009

<1%

Publication

75

Jenny S Jennings, Beverly E Meyer, Pablo J Guiroy, N Andy Cole. "Energy costs of feeding excess protein from corn-based by-products to finishing cattle", *Journal of Animal Science*, 2018

<1%

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Effects of Replacing Concentrate with Soybean Curd Residue Silage on Ruminal Characteristics, Plasma Leucine and Glucose Turnover Rates in Sheep

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15
