

Calcium deposition in egg due to substitution of limestone by eggshell flour in feed of laying hens

by Sri Kismiyati

Submission date: 11-Mar-2020 02:49PM (UTC+0700)

Submission ID: 1273528446

File name: Calcium_deposition_in_egg_due_to_substitution_of_limestone.pdf (151.99K)

Word count: 5139

Character count: 25581

Calcium deposition in egg due to substitution of limestone by eggshell flour in feed of laying hens

S. Kismiati¹, T. Yuwanta², Zuprizal², Supadmo² and U. Atmomarsono¹

¹Faculty of Animal and Agricultural Sciences, Diponegoro University,
Tembalang Campus, Semarang, 50275 - Indonesia.

² Faculty of Animal Science, Universitas Gadjah Mada,
Jl. Fauna 3 Bulaksumur 55281 Yogyakarta - Indonesia
Corresponding E-mail : kismiati59@gmail.com

Received February 02, 2018; Accepted April 09, 2018

ABSTRAK

Penelitian bertujuan untuk mengevaluasi deposisi kalsium pada telur ayam yang menggunakan tepung kerabang telur sebagai pengganti batu kapur dalam pakan. Dua ratus ekor ayam petelur strain Isa Brown umur 25 minggu digunakan pada penelitian ini. Perlakuan yang dicobakan adalah : penggunaan batu kapur 7,5% dalam pakan sebagai kontrol (T0), penggantian 2,5% batu kapur dengan tepung kerabang telur (T1), penggantian 5% batu kapur dengan tepung kerabang telur (T2) dan penggantian batu kapur dengan tepung kerabang telur (T3). Rancangan percobaan yang digunakan adalah Rancangan Acak Lengkap dengan 4 perlakuan dan 5 ulangan. Tiap unit percobaan terdiri 10 ekor ayam petelur. Parameter yang diukur meliputi berat telur, berat kuning telur, berat albumen, berat kerabang telur dan kalsium telur (kuning telur, albumen dan kerabang telur), panjang, berat dan Ca tulang tibia. Hasil penelitian menunjukkan bahwa penggantian batu kapur dengan tepung kerabang telur berpengaruh nyata ($P < 0,05$) berat kerabang telur, deposisi Ca pada kuning telur, putih telur dan tulang tibia, tetapi tidak berpengaruh nyata terhadap berat telur, berat dan persentase kuning telur, berat dan persentase putih telur, persentase dan Ca kerabang telur, panjang dan berat tulang tibia. Simpulan bahwa deposisi Ca pada kuning telur tertinggi pada penggunaan 7,5% tepung kerabang telur sebagai pengganti batu kapur tetapi diperoleh Ca tulang terendah, sedangkan deposisi Ca pada putih telur tertinggi pada penggunaan 2,5% tepung kerabang telur.

Kata kunci : batu kapur, kerabang telur, deposisi kalsium

1 ABSTRACT

The aim of this research was to evaluate calcium deposition in egg using eggshell flour as a limestone substitute in feed. Two hundreds laying hen of Isa Brown strain of 25 weeks were used in this study. Treatments were diet with 7.5% limestone as control (T0), 2.5% limestone is substituted with eggshell flour (T1), 5% limestone is substituted with eggshell flour (T2) and limestone is substituted with eggshell flour (T3). A completely randomized design were used to allocated the treatments with 5 replications of each. Each experimental unit consists of 10 laying hen. Parameters measured were egg weight, yolk weight, albumen weight, eggshell weight, calcium of egg (yolk, albumen and eggshell), length, weight and Ca of tibia bone. The results showed that substitution of limestone with eggshell flour had significantly effect ($P < 0.05$) eggshell weight, Ca deposition on yolk, albumen, and Ca of tibia bone but non significantly effect on egg weight, weight and percentage of yolk, weight and percentage

of albumen, percentage and Ca of eggshell, length and weight of tibia bone. In conclusion, calcium deposition in yolk was the highest in the use of 7.5% eggshell flour to substitute limestone but obtained the lowest Ca of bone, while calcium deposition in albumen was the highest in the use of 2.5% eggshell flour.

Keywords: limestone, eggshell, calcium deposition

INTRODUCTION

Poultry industry development may give positive and negative impacts. The positive impact was nutrition sufficiency and job opportunities, the negative impact was waste product of excreta and eggshell. Excreta can be used as organic fertilizer and source of energy, while eggshell if not handled properly will effect on environmental pollution.

Egg production of Indonesia was 1401079.2 tons/year (Direktorat Jenderal Peternakan dan Kesehatan Hewan, 2016), it is estimated that there are 140107.92 tons/year eggshell waste and not utilized. The eggshell is covered with cuticles. The cuticle has a compositional with proteins, sulfated polysaccharides and phosphates (Rodríguez-Navarro *et al.*, 2013). The eggshell membranes consists of highly collagens, glycoproteins and cysteine-rich, eggshell membrane proteins as used in the development of the embryo and protect it against pathogen invasion (Cordeiro and Hinckley, 2016).

According to Panheleux *et al.* (2000), Mine *et al.* (2003), Rodríguez-Navarro *et al.* (2015), Marie *et al.* (2015) and Zang *et al.* (2016) eggshell contains minerals and a little protein. Proteins involved in mineral supply during eggshell calcification (Brionne *et al.*, 2014). Yasothai and Kavithaa (2014) stated that eggshell is highly rich in calcium (90%) with little of phosphorus. Siuplapwa *et al.* (2014) reported that the eggshell contains various minerals ie Ca, P, Fe, Mg, Cu, Co, K, Na, Zn and C. Mineral composition of eggshell are Ca (35080 mg/100 g), P (150.2 mg/100 g), Mg (262mg/100g), Fe (13.06 mg/100 g), Zn (145.1 mg/g), K (50 mg/100 g), Cu (4.1 mg/100 g) and Mn (149.9 mg/100 g) (Hassan, 2015).

Calcium is one of minerals essential for poultry. Requirement of Ca of poultry for laying period was around 4 - 5 times compared with brooding and growing period. Calcium content of the feed will affected the eggshell and bone quality of laying hens (An *et al.*, 2016). Neijat *et al.* (2011) stated that Ca consumption was deposite on the bone, meat, eggshell, albumen, yolk and some

other will excrete through excreta. Medullary bone acts as Ca storage for eggs formation especially eggshell (Kim *et al.*, 2012) and reserve of eggshell formation when Ca deficiency in feed (Olgun and Aygun, 2016). Ca consumption of laying hen was 4000 mg and deposit to the egg 3100 mg, faeces 500 mg and urine 400 mg (Leeson dan Summers, 2005). Ca of egg is needed for embryo development (Uni *et al.*, 2012, Tahara and Obara, 2014; Scanes, 2015).

Ca source is widely used as poultry feed material is limestone. Limestone contains many minerals but no protein, while eggshell flour contains mineral and a little protein. Noviyanti *et al.* (2015) stated that limestone composed by Ca, K, Na, dan Si. Limestone was composed by macro mineral Ca, P, Mg, Na, K, Cl, S and micromineral Mn, Zn, Fe, Cu, Co, and Se (Khalil and Anwar, 2009). Limestone was composed 35.7 - 36.4% of Ca (Pelicia *et al.*, 2009).

Alu *et al* (2013) stated that eggshell flour may used for substitution of bone flour on broiler diet without blood hematology and nutrient digestibility changing. Olgun *et al.* (2015) stated that differences of Ca source showed significance effected on egg production and eggshell quality. Research of Cufadar *et al.* (2014) showed that used of eggshell flour, limestone, oyster shell, mixture of limestone with eggshell, limestone with oyster shell and eggshell with oyster shell showed non significance effect on the egg weight and eggshell weight. Research of Scheideler (1998) showed that Ca of eggshell has higher digestibility then of limestone. According to Dolinska *et al.* (2016), calcium carbonat (CaCO₃) of eggshell has a higher solubility than of synthetic CaCO₃. Szeleszczuk *et al.* (2015) stated that the *in vitro* solubility Ca of eggshell is higher than CaCO₃. The aim of this research was to evaluate a substitution of limestone with eggshell flour on the Ca deposition in egg of laying hens.

MATERIALS AND METHODS

Two hundred laying hens (Isa Brown) were used in this study. The experimental period was 12 weeks. The laying hens were distributed in a

Completely Randomized Design with 4 treatments and 5 replications. The treatment were limestone of 7.5% in feed (T0), substitution 2.5% lime limestone with eggshell flour (T1), substitution of 5% limestone with eggshell flour (T2) and the replacement of limestone with eggshell flour (T3). The composition of feedstuff and nutrient content of dietary treatment is presented in Table 1. Before use, the eggshell was immersed in water temperature 80 °C for 15 minutes, drained and then immersed into 5% phosphoric acid, dried by the sun for 3 days and grounded with 3 mm particle size.

Parameter measured were weight of egg, weight and percentage of yolk, weight and percentage of albumen, weight and percentage of eggshell, Ca of yolk, Ca of albumen, Ca of eggshell and length, weight and Ca content of tibia bone. Analysis of calcium yolk, calcium albumen and calcium eggshell using AOAC method (Horwitz and Latimer, 2005). Data was analyzed by Analysis of Variance (ANOVA) and continued with Duncan's Multiple Range Test (DMRT) when there was significantly difference.

RESULTS AND DISCUSSIONS

Calcium Deposition on Egg

Calcium deposition on egg with feed supplementation eggshell flour as a substitution of limestones is presented in Table 2. Used of eggshell flour substitution of limestone showed an effect ($P < 0.05$) eggshell weight, Ca of yolk, Ca of albumen but there was no effect on egg weight, weight and percentage of yolk, weight and percentage of albumen, percentage of eggshell weight and Ca of eggshell .

Egg Weight

Results of this research showed that used of eggshell flour up to 7.5% as substitution of limestone did not affect on egg weight. Eggshell flour contains Ca almost the same as limestone, so that the use of eggshell flour as a substitute for limestone had no effect on egg weight. Gongruttananun (2011) stated that Ca content of eggshell is 34.89%, while a content of limestone is 36.4 – 35.7% (Pelicia *et al.*, 2009). The use of eggshell flour as a substitute of limestone in diet

Table 1. Composition of Feed Stuff and Nutrient Content of Dietary Treatment

Feedstuffs	T0	T1	T2	T3
%			
Corn	70.00	70.00	70.00	70.00
Soybean meal	10.00	10.00	10.00	10.00
Poultry meat meal	11.00	11.00	11.00	11.00
Limestone	7.50	5.00	2.50	0.00
Eggshell flour	0.00	2.5	5.0	7.5
DCP	1.00	1.00	1.00	1.00
Salt	0.25	0.25	0.25	0.25
Calculated Nutritional Level				
Crude protein (%)	16.17	16.31	16.45	16.59
EM (kcal/kg)	2892.50	2892.50	2892.50	2892.50
Ca (%)	3.44	3.44	3.44	3.44
P (%)	0.46	0.50	0.55	0.59
Crude fiber (%)	2.70	2.70	2.70	2.70
Lysine (%)	1.10	1.10	1.10	1.10
Methionine (%)	0.44	0.44	0.44	0.44

T0: feed using 7.5% of limestone, T1: 2.5% of limestone replaced by eggshell flour, T2: 5% limestone replaced by eggshell flour, T3: 7.5% limestone replaced by eggshell flour

Table 2. Calcium Deposition in Egg on using Eggshell Flour as Substitution of Limestone in the Feed of Laying Hens

Parameters	T0	T1	T2	T3
Egg weight(g)	51.418	50.567	53.027	52.101
Yolk weight (g)	12.536	12.317	13.292	13.681
Yolk percentage (%)	24.406	24.330	25.106	26.246
Albumen Weight (g)	33.943	32.898	34.452	33.116
Albumen percentage (%)	65.972	65.061	64.919	63.549
Eggshell weight (g)	4.938 ^b	5.352 ^{ab}	5.283 ^a	5.304 ^a
Eggshell percentage (%)	9.53	10.58	9.8	10.17
Ca Yolk (mg)	6.086 ^d	6.865 ^c	8.433 ^b	11.359 ^a
Ca Albumen (mg)	1.618 ^c	2.293 ^a	1.922 ^b	1.249 ^d
Ca Eggshell (g)	1.48	1.49	1.55	1.41

T0: feed using 7.5% of limestone, T1: 2.5% of limestone replaced by eggshell flour, T2: 5% limestone replaced by eggshell flour, T3: 7.5% limestone replaced by eggshell flour

of laying hen was not effect on the egg weight with the egg weight is 47.47 – 47-81 g (Gongruttananun, 2011). Result of the research of Cufadar (2014) showed that different source of Ca was not affected on the egg weight. According to Sultana *et al.* (2007) and Safaa *et al.* (2008) differences of Ca source and Ca content in diet did not effect on egg weight. An *et al.* (2016) confirmed that was not the factor affecting to the egg weight. Hanusová *et al.* (2015) stated that egg weight was affected by genetic factor. Leeson and Summers (2005) and King'ori (2012) stated that factor affecting the egg weight were age, body weight and nutrition consumption. According to Adeyemo *et al.* (2012), protein content in diet was effect on the egg weight.

Weight and Percentage of Yolk

The used of eggshell flour as substitution of limestone did not affect the yolk weight and percentage of yolk. The results of the study was in agree to the results of previous studies. Saldanha *et al.* (2009) stated that percentage of yolk was not affected by source of Ca. The research of Safaa *et al.* (2008) showed that the percentage of yolk was not different between used limestone or oyster shell powder as a source of Ca in the feed. Gongruttananun (2011) stated that substitution of limestone with eggshell flour did not effect the yolk percentage. Pelicia *et al.* (2009) reported that

Ca in the diet was not affected on percentage of yolk. Factor affected the percentage of yolk is genetic (Hanusová *et al.*, 2015), protein level of the feed (Gunawardana *et al.*, 2008), eggs storage (Scott dan Silversides, 2000) and age of laying hens (Silversides and Budgell, 2004). Longer time of storage will increase weight of yolk (Khan *et al.*, 2014). Yolk percentage was positive correlated to weight of egg and age of laying hens (Ulmer-Franco *et al.*, 2012).

Weight and Percentage of Albumen

Result of the research showed that used of eggshell flour as substitution of limestone did not affect the weight and percentage of albumen. Saldanha *et al.* (2009) and Safaa *et al.* (2009) stated that Ca source feed did not affect percentage of albumen. The results of the study were in agree with the research of Gongruttananun (2011) that the limestone substitution with eggshell flour in feed of Rhode Island Red strain did not significantly affect the percentage of albumen. Pelicia *et al.* (2007) state that difference source of Ca (limestone or marine calcium) did not affect the weight of albumen. Ca level in the feed did not affect the albumen percentage (Pelicia *et al.*, 2009). The factor influencing albumen weight was genetic (Suk and Park, 2001; Hanusová *et al.*, 2015). Silversides and Budge (2004) stated that factors affecting the

weight of albumen were age of laying hens and a long of storage.

Weight and Percentage of Eggshell

Table 2 showed that substitution of 5% and 7.5% limestone with eggshell flour will increase weight of eggshell ($P < 0.05$). Increasing of eggshell weight was affected by protein content and minerals (Fe, 41g, Zn, Cu, and Se) of eggshell flour. Panheleux *et al.* (2000), Mine *et al.* (2003), Rodríguez-Navarro *et al.* (2015), Marie *et al.* (2015) and Zang *et al.* (2016) stated that eggshell contains many minerals and very low of protein. Schaafsma *et al.* (2000) stated that eggshell contained high Ca and other minerals such as P, Mg, Fe, Zn, Se, Cr, Na, F, Cr, V and slightly protein. Khalil and Anwar (2009) and Noviyanti *et al.* (2015) stated that limestone contains minerals but without protein. Supplementation of Mn, Zn, and Cu increased eggshell quality (Stefanello *et al.* 2014).

Eggshell percentage showed 37 different. Khalil and Anwar (2009) reported that eggshell percentage was not affected by different of Ca source of feed. Research of Gongruttananun (2011) showed that used of eggshell flour as substitution of limestone showed non significance effect the eggshell percentage. Olgun *et al.* (2015) stated that used of eggshell flour as substitution of limestone (66.7%) did not affect the percentage of eggshell.

Ca of Yolk

Substitution of limestone by eggshell flour affected ($P < 0.05$) the Ca deposition in yolk. Ca deposition on yolk increased along with used of eggshell flour as substitution for limestone in feed. The result showed that Ca utilization from eggshell flour was higher compared to limestone. This result was supported by statement of Scheideler (1998) in which digestibility of eggshell flour was higher than limestone and Szeleszczuk *et al.* (2015) stated that bioavailability Ca of eggshell was high. Leeson dan Summers (2005) stated that calcium of diet was storage in bone, deposited in eggs and excreted on the excreta. Differences of Ca source was not affected the Ca of yolk (Vargas-Rodríguez *et al.*, 2016).

Ca of Albumen

There was significantly effect ($P < 0.05$) of substitution of limestone with eggshell flour on Ca deposition in albumen. Substitution of

limestone with eggshell flour would increase Ca albumen. The highest of Ca deposition was found on T1 (2.5% or 33.33% eggshell flour). Kiczorowska *et al.* (2015) stated that Ca of yolk positively correlates with Ca albumen, but the results of this study Ca of albumen would increase on T1 and followed by decreasing in T2 and T3, but T2 and T3 were still higher than T0. Calcium of albumen decreased along with increased calcium of yolk. This 36 result was different from the result of Pastore *et al.* (2012) and Vargas-Rodríguez *et al.* (2016) that Ca of the feed did not affect the deposition Ca in albumen.

Ca of Eggshell

Ca deposition in eggshell was not significantly different, whereas Ca deposition in yolk and albumen was higher ($P < 0.05$) on the use of eggshell flour than limestone (Table 2). Tunç and Cufadar (228) stated that different Ca sources of feed had no significantly effect on eggshell weight, but significant on Ca deposition in eggshell. The research of Olgun *et al.* (2015) showed that the different calcium sources had significantly effect on eggshell weight and eggshell calcium content.

Length, Weight and Calcium (Ca) Content of Tibia Bone

Result of this study showed that substitution of limestone by eggshell flour significantly effect on Ca content 27 tibia bone but no significantly effect on the length and weight of tibia bone (Table 3).

Length and Weight of Tibia Bone

The length and weight of tibia bone were not significantly different. The results of this study were in accordance with the study of Gongruttananun (2011) that the use of eggshell flour had no significant effect on the length and weight of tibia bone.

Calcium Content of Tibia Bone

Table 3 showed that the replacement of 5% limestone with eggshell flour increased the calcium percentage and calcium content of tibia bone, but substitution of 7.5% limestone with eggshell flour decreased the percentage and weight calcium of bone. This is due to increase Ca deposition in yolks (Table 2). The results showed that the use of 7.5% eggshell flour has occurred Ca mobilization of tibia bone for egg formation, especially egg yolks. Tunç *et al.* (2015) stated

Table 3. Length, Weight and Ca Content of the Tibia Bone on Using Eggshell Flour as Substitution of Limestone in the Feed of Laying Hens

Parameters of Tibia Bone	T0	T1	T2	T3
Length (cm)	11.90	11.62	11.80	11.67
Weight (cm)	10.27	11.05	10.89	10.19
Ca (%)	12.80 ^{ab}	12.60 ^{ab}	13.61 ^a	12.03 ^b
Ca (g)	1.31 ^{ab}	1.39 ^{ab}	1.480 ^a	1.22 ^b

26 that the Ca content tibia bone is affected by the calcium source, but the results of this study show not significant effect.

The result 25 of this study was different from research of Saunders-Blades *et al.* (2009) in which the weight and calcium of tibia were not affected by calcium sources. Tümová *et al.* (2016) and Swiatkiewicz *et al.* (2015) suggested that tibia bone Ca is affected by Ca levels of feed.

CONCLUSION

Calcium deposition in yolk was the highest in the use of 7.5% eggshell flour to substitute limestone, while calcium deposition in albumen was the highest in the use of 2.5% eggshell flour. The use of 7.5% eggshell flour as a substitute for limestone resulted in the mobilization of Ca from the tibia bone so that the Ca content of the tibia bone decreased.

REFERENCES

- Adeyemo, G.O., S.A. Abioye and F. A. Aderemi. 2012. The effect of varied dietary crude protein levels with balanced amino acids on performance and egg quality characteristics of layers at first laying phase. *Food and Nutr. Sci.* 3: 526-529.
- An, S. H., D. W. Kim and B. K. An. 2016. Effects of dietary calcium levels on productive performance, eggshell quality and overall Calcium status in aged laying hens. *Asian-Aust. J. Anim. Sci.* 29 (10):1477-1482.
- Brionne, A., Y. Nys, C. Hennequet-Antier and J. Gautron. 2014. Hen uterine gene expression profiling during eggshell formation reveals putative proteins involved in the supply of minerals or in the shell mineralization process. *BMC Genomics* 15:1 - 17.
- Cufadar, Y. 2014. Effect of alternative Calcium sources on performance and eggshell quality in laying hens. *J. Macro. Trends Appl. Sci.* 2 (1):10-15.
- Cordeiro, C.M. And M.T. Hincke. 2016. Quantitative proteomics analysis of eggshell membrane proteins during chick embryonic development. *J. Proteomics* 130:11 -13.
- Direktorat Jenderal Peternakan dan Kesehatan Hewan. 2016. Statistik Peternakan dan Kesehatan Hewan. Kementerian Pertanian Republik Indonesia.
- Dolińska, B., M. Jelińska, B. Szulc-Musiół and F. Ryszka. 2016. Use of eggshells as a raw material for production of calcium preparations. *Czech J. Food Sci.* 34(4):313-317.
- Gunawardana, P., D. A. Roland Sr. and M. M. Bryant. 2008. Effect of energy and protein on performance, egg components, egg solids, egg quality, and profits in molted Hy-Line W-36 hens. *Appl. Poult. Res.* 17:432 - 439.
- Gongruttanun, N. 2011. Effects of eggshell calcium on productive performance, plasma calcium, bone mineralization, and gonadal characteristics in laying hens. *Poult. Sci.* 90:524 - 529.
- Hall, R.E., R.B. Shirly, R.I. Bakalli, S.E. Aggrey, S.E., G.M. Pesti and H.M. Edwards Jr. 2003. Power of two methods for the estimation of bone ash of broiler. *Poult. Sci.* 82: 414 - 418.
- Hanusová, E., C. Hrnčár, A. Hanus and M. Oravcová. 2015. Effect of breed on some parameters of egg quality in laying hens. *Acta Fytotechn. Zootechn.*, 18 (1): 20-24.
- Hassan, N.M.M. 2015. Chicken eggshell powder as dietary calcium source in biscuits. *World J. Dairy Food Sci.* 10 (2): 199 - 206.
- Horwitz, W. and Latimer, G.W. 2005. Official

- Methods of Analysis of AOAC International 18th Edition. AOAC International. Gaithersburg, USA.
- Khan, M.J.A., S. H. Khan, A. Bukhsh, and M. Amin. 2014. The effect of storage time on egg quality and hatchability the effect of storage time on egg quality and hatchability characteristics of Rhode Island Red (RIR) hens characteristics of Rhode Island Red (RIR) hens. *Vet. Arhive.* 84 (3), 291-303.
- Khalil and S. Anwar. 2009. Limestone of Bukit Kemang as a calcium source for laying hens. *J. Ind. Trop. Anim. Agric.* 34(3):174-180.
- Kiczorowska, B., W. Samolińska, M. Kwiecień, A. Winiarska-Mieczan, E. Rusinek-Prystupa and A. R. M. Al-Yasiry. 2015. Nutritional value and the content of minerals in eggs produced in large-scale, courtyard and organic systems. *J. Elem.* 20(4): 887-895.
- Kim, W.K., S.A. Bloomfield, T. Sugiyama and S.C. Rick. 2012. Concepts and methods for understanding bone metabolism in laying hens. *World Poult. Sci. J.* 68:71-82.
- King'ori, A.M. 2012. Poultry egg external Characteristics : egg weight, shape and shell colour. *Res. J. Poult. Sci.* 5 (2):17-14.
- Leeson, S. and J.D. Summers. 2005. *Commercial Poultry Nutrition 3rd Edition.* University Books, Guelph, Ontario, Canada.
- Marie, P.P., V. Labas, A. Brionne, g. Harichaux, C. Hennequetntier, Y. Nys and J. Gautron. 2015. Quantitative proteomics and bioinformatic analysis provide new insight into protein function during avian eggshell biomineralization. *J. Proteomics* 113:178-193.
- Mine, Y., C. Oberle and Z. Kassaiy. 2003. Eggshell matrix proteins as defense mechanism of avian eggs. *J. Agric. Food Chem.* 51(1):249-253.
- Neijat, M, J. D. House, W. Guenter, and E. Kebreab. 2011. Calcium and phosphorus dynamics in commercial laying hens housed in conventional or enriched cage systems. *Poult. Sci.* 90:2383-2396.
- Noviyanti, Jasruddin and E. H. Sujiono. 2015. Karakterisasi kalsium karbonat (CaCO₃) batu kapur kelurahan Tellu Limpoe kecamatan Suppa. *Jurnal Sains dan Pendidikan Fisika* 11(2):169-172.
- Olgun, O., A.O. Yildiz and Y Cufadar. 2015. The effects of eggshell and oyster shell supplemental as calcium sources on performance, eggshell quality and mineral excretion in laying hens. *Indian J. Anim. Res.* 49(2):205-209.
- Olgun, O. and A. Aygun. 2016. Nutritional factors affecting the breaking strength of bone in laying hens. *World Poult. Sci. J.* 72(4):821-832.
- Panheleux, Y. Nys, J. Williams, J. Gautron, T. Boldicke, and M. T. Hinck. 2000. Extraction and quantification by ELISA of eggshell organic matrix proteins (ovocleidin-17, ovalbumin, ovotransferrin) in shell from young and old hens. *Poult.Sci.* 79(4):580-588.
- Pastore, S.M. , P. C. Gomes, H. S. Rostagno, L. F. T. Albino, A.A. Calderano, C. R. Vellasco, G. da Silva Viana and R. L. de Almeida. 2012. Calcium levels and calcium : available phosphorus ratios in diets for white egg layers from 42 to 58 weeks of age. *R. Bras. Zootec.* 41(12):2424-2432.
- Pelicia, K., E. Garcia, C. Móri, A.B.G. Faitarone, A.P. Silva, A.B. Molino, F. Vercese and D.A. Berto. 2009. Calcium levels and limestone particle size in the diet of commercial layers at the end of the first production cycle. *Braz. J. Poult. Sci.* 11(2):87- 94.
- Pelícia, K., E.A. Garcia, M.R.S. Scherer, C. Móri, J.A. Dalanezi, A.B.G. Faitarone, A.B. Molino and D.A. Berto. 2007. Alternative Calcium source effects on commercial egg production and quality. *Braz. J. Poul. Sci.* 9 (2): 105-109.
- Rodríguez-Navarro, A.B., N. Domínguez-Gasca, N. A. Muñoz and M. Ortega-Huertas. 2013. Change in the chicken eggshell cuticle with hen age and egg freshness. *Poult. Sci.* 2: 3026-3035.
- Rodríguez-Navarro, A.B., P. Marie, Y. Nys, M.T. Hincke and J. Gautron. 2015. Amorphous calcium carbonate controls avian eggshell mineralization: A new paradigm for understanding rapid eggshell calcification. *J. Struct. Biol.* 190 (3):291-303.
- Safaa, H. M. , M. P. Serrano, D. G. Valencia, M. Frikha, E. Jiménez-Moreno, and G. G. Mateos. 2008. Productive performance and egg quality of brown egg-laying hens in the late phase of production as influenced by level and source of calcium in the diet. *Poult. Sci.* 87:2043-2051.
- Saldanha, E.S.P.B. , EA Garcia, C.C. Pizzolante, A.B.G.Faitarone, A. Sechinato A da, A.B.

- Molino and C. Laganá. 2009. Effect of organic mineral supplementation on the egg quality of semi-heavy layers in their second cycle of lay. *Braz. J. Poult. Sci.* 11(4):215-222.
- Saunders-Blades, J. L., J. L. MacIsaac, D. R. Korver and D. M. Anderson. 2009. The effect of calcium source and particle size on the production performance and bone quality of laying hens. *Poult. Sci.* 88(2):338-353.
- Schaafsma, A., I. Pakan., G.J.H. Hpfstede, F.A.J. Muskiet, E. Van Der Veer and P.J.F. De Vries. 2000. Mineral, amino acid, and hormonal composition of chicken eggshell powder and the evaluation of its use in human nutrition. *Poult. Sci.* 79(12):1833-1838.
- Scanes, C.G. 2015. *Sturky's Avian Physiology*. Sixth Edition. Academic Press, Elsevier Inc.
- Scheideler, S.E. 1998. Eggshell calcium effects on egg first – or third cycle laying hens quality and Ca digestibility. *J. App. Poult. Res.* 7:69-74.
- Scott, T.A. and F. G. Silversides. 2000. The effect of storage and strain of hen on egg quality. *Poult. Sci.* 79(12):1725-1729.
- Silversides, F.G. and K. Budgell. 2004. The relationships among measures of egg albumen height, pH, and whipping volume. *Poult. Sci.* 83(10):1619-1623.
- Siuplapwa, A. Mwambungu and L. Mubbunu. 2014. Comparison of mineral composition of commercial hen egg shell to fresh water crocodile egg shell. *Int. J. Res. Agric. Food Sci.* 2(7):16-18.
- Stefanello, C., T. C. Santos, A. E. Murakami, E. N. Martins, and T. C. Carneiro. 2014. Productive performance, eggshell quality, and eggshell ultrastructure of laying hens fed diets supplemented with organic trace minerals. *Poult. Sci.* 93(1):104-113.
- Suk, Y.O. and C. Park. 2001. Effect of breed and age of hens on the yolk to albumen ratio in two different genetic stocks. *Poult. Sci.* 80(7):855-858.
- Swiatkiewicz, S., A. Arczewska-Wlosek and D. Jozefiak. 2015. Bone quality, selected blood variables and mineral retention in laying hens fed with different dietary concentrations and sources of calcium. *Livest. Sci.* 181:194-199.
- Szeleszczuk, L., D. M. Pisklak, M. Kuras and I. Wawer. 2015. *In vitro* dissolution of calcium carbonate from the chicken eggshell: A study of calcium bioavailability. *Int. J. Food Properties* 18(12):2791-2799.
- Tahara, Y. and K. Obara. 2014. A novel shell-less culture system for chick embryos using a plastic film as culture vessels. *Japan Poult. Sci.* 51(3):307-312.
- Tůmová, E., J. Vlčková, V. Charvátová, O. Drábek, V. Tejnecký, M. Ketta and D. Chodová. 2016. Interactions of genotype, housing and dietary calcium in layer performance, eggshell quality and tibia characteristics. *South African J. Anim. Sci.* 46(3):285-293.
- Tunç, A.E., Y. Cufadar. 2015. Effect of Calcium sources and particle size on performance and eggshell quality in laying hens. *Turkish J. Agric.- Food Sci. and Technol.* 3(4):205-209
- Ulmer-Franco, A.M., G. Cherian, N. Quezada, G. M. Fasenko and L. M. McMullen. 2012. Hatching egg and newly hatched chick yolk sac total IgY content at 3 broiler breeder flock ages. *Poult. Sci.* 91(3):758-764.
- Uni, Z., L. Yadgary and R. Yair. 2012. Nutritional limitations during poultry embryonic development. *J. Appl. Poult. Res.* 21(1):175-184
- Vargas-Rodríguez, L.M., J. E. Morales-Barrera, J. G. Herrera-Haro, J. Antonio-Bautista, R. López-Pozos and D. Hernández-Sánchez. 2016. Effect of citric acid, phytase and Calcium levels on the Calcium and Phosphorus content in egg: yolk-albumen and shell, yolk color and egg quality in diets of laying hens. *Food and Nutr. Sci.* 7(14):1364-1374.
- Yasothei, R. and N.V. Kavithaa. 2014. Chemical characterization of egg shell meal. *Int. J. Sci. Environment and Technol.* 3(4):1436-1439
- Zang, M., N. Wang, Q. Xu, P.W. Harlina and M. Ma. 2016. An efficient method for co-purification of eggshell matrix proteins OC-17, OC-116, and OCX-36. *Korean J. Food Sci. Anim. Resour.* 36(6):769-778.

Calcium deposition in egg due to substitution of limestone by eggshell flour in feed of laying hens

ORIGINALITY REPORT

13%

SIMILARITY INDEX

8%

INTERNET SOURCES

10%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

- 1 worldwidescience.org 1%

Internet Source
- 2 John N. Maina. "Chapter 9 Structure and Function of the Shell and the Chorioallantoic Membrane of the Avian Egg: Embryonic Respiration", Springer Science and Business Media LLC, 2017 1%

Publication
- 3 www.cambridge.org 1%

Internet Source
- 4 fr.scribd.com 1%

Internet Source
- 5 Astorga Espana, M.S.. "Comparison of mineral and trace element concentrations in two molluscs from the Strait of Magellan (Chile)", Journal of Food Composition and Analysis, 200705 <1%

Publication

6

"Karakteristik Karkas Sapi Peranakan Simmental Jantan Di Rumah Potong Hewan Kota Semarang Carcass Characteristics Of Simmental Crossbreed Cattle Slaughtered At Semarang City Slaughterhouse", Jurnal Pengembangan Penyuluhan Pertanian, 2018

Publication

<1%

7

www.locus.ufv.br

Internet Source

<1%

8

www.i-scholar.in

Internet Source

<1%

9

peternakan.unpad.ac.id

Internet Source

<1%

10

Ruiqiao Li, Xuefeng Qi, Xueying Han, Caihong Liu, Jing Wang, Ruichun Wang, Jingyu Wang, Jinhai Huang. "Deterioration of eggshell quality is related to calbindin in laying hens infected with velogenic genotype V/Id Newcastle disease virus", Theriogenology, 2017

Publication

<1%

11

Traineau, M., I. Bouvarel, C. Mulsant, L. Roffidal, C. Launay, and P. Lescoat. "Effects on performance of ground wheat with or without insoluble fiber or whole wheat in sequential feeding for laying hens", Poultry Science, 2013.

Publication

<1%

12

Submitted to Fakultas Ekonomi dan Bisnis
Universitas Gadjah Mada

Student Paper

<1%

13

bmcgenomics.biomedcentral.com

Internet Source

<1%

14

www.dlg.dk

Internet Source

<1%

15

Submitted to Higher Education Commission
Pakistan

Student Paper

<1%

16

Muhammad Waqas, Hammad Qamar, Jialu Zhang, Wangyuan Yao et al. "Puerarin enhance vascular proliferation and halt apoptosis in thiram-induced avian tibial dyschondroplasia by regulating HIF-1 α , TIMP-3 and BCL-2 expressions", *Ecotoxicology and Environmental Safety*, 2020

Publication

<1%

17

www.scielo.br

Internet Source

<1%

18

S. Swiatkiewicz, A. Arczewska-Wlosek, D. Jozefiak. "Bone quality, selected blood variables and mineral retention in laying hens fed with different dietary concentrations and sources of calcium", *Livestock Science*, 2015

Publication

<1%

19

Y. H. Jin, K. T. Lee, W. I. Lee, Y. K. Han.
"Effects of Storage Temperature and Time on
the Quality of Eggs from Laying Hens at Peak
Production", Asian-Australasian Journal of
Animal Sciences, 2010

Publication

<1%

20

mgavm.ru

Internet Source

<1%

21

K.I Ereifej, S.G Haddad. "Chemical composition
of selected Jordanian cereals and legumes as
compared with the FAO, Moroccan, East Asian
and Latin American tables for use in the Middle
East", Trends in Food Science & Technology,
2000

Publication

<1%

22

ijset.net

Internet Source

<1%

23

eprints.ulm.ac.id

Internet Source

<1%

24

fapet.ub.ac.id

Internet Source

<1%

25

animalsciencepublications.org

Internet Source

<1%

26

Submitted to Selçuk Üniversitesi

Student Paper

<1%

27 Muhammad Waqas, Yaping Wang, Aoyun Li, Hammad Qamar et al. "Osthole: A Coumarin Derivative Assuage Thiram-Induced Tibial Dyschondroplasia by Regulating BMP-2 and RUNX-2 Expressions in Chickens", *Antioxidants*, 2019
Publication

28 www.tandfonline.com
Internet Source

29 Mehran Torki, Ahmad Mohebbifar, Hossein Ali Ghasemi, Afshin Zardast. "Response of laying hens to feeding low-protein amino acid-supplemented diets under high ambient temperature: performance, egg quality, leukocyte profile, blood lipids, and excreta pH", *International Journal of Biometeorology*, 2014
Publication

30 Sakine Yalçın. "Effects of dietary black cumin seed (*Nigella sativa* L.) on performance, egg traits, egg cholesterol content and egg yolk fatty acid composition in laying hens", *Journal of the Science of Food and Agriculture*, 2009
Publication

31 researcherslinks.com
Internet Source

32 repositorio.ufsm.br
Internet Source

-
- 33 Ricke, S. C., C. S. Dunkley, and J. A. Durant. "A review on development of novel strategies for controlling Salmonella Enteritidis colonization in laying hens: Fiber-based molt diets", Poultry Science, 2013.
Publication <1%
-
- 34 docplayer.net
Internet Source <1%
-
- 35 garuda.ristekdikti.go.id
Internet Source <1%
-
- 36 Submitted to Mansoura University
Student Paper <1%
-
- 37 Y. Cufadar, O. Olgun, A.Ö. Yildiz. "The effect of dietary calcium concentration and particle size on performance, eggshell quality, bone mechanical properties and tibia mineral contents in moulted laying hens", British Poultry Science, 2011
Publication <1%
-
- 38 Antonio González Ariza, Francisco Navas González, Ander Arando Arbulu, José León Jurado et al. "Non-Parametrical Canonical Analysis of Quality-Related Characteristics of Eggs of Different Varieties of Native Hens Compared to Laying Lineage", Animals, 2019
Publication <1%
-

39

N. SAH, B. MISHRA. "Regulation of egg formation in the oviduct of laying hen", World's Poultry Science Journal, 2018

Publication

<1%

40

Kim, Young Jik, Gee Dong Lee, and In Hag Choi. "Effects of dietary red ginseng marc on egg production, egg quality and blood characteristics of laying hens", Journal of Applied Animal Research, 2015.

Publication

<1%

41

Vojtěch Anderle, Martina Lichovnicková, Alena Przywarova, Eliška Dračková. "Egg quality of gene reserve Czech golden spotted hens", Acta fytotechnica et zootechnica, 2014

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Calcium deposition in egg due to substitution of limestone by eggshell flour in feed of laying hens

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
