

Effect of H₃PO₄ Concentration and Particle Size of the Eggshell Used in Laying Hens Fed on Bone and Blood

by Sri Kismiyati

Submission date: 11-Mar-2020 02:51PM (UTC+0700)

Submission ID: 1273528792

File name: ect_of_H₃PO₄_Concentration_and_Particle_Size_of_the_Eggshell.pdf (204.81K)

Word count: 3775

Character count: 18084



Effect of H_3PO_4 Concentration and Particle Size of the Eggshell Used in Laying Hens Fed on Bone and Blood

S. Kismiati^{1,2}, T. Yuwanta², Zuprizal² and Supadmo²

¹Faculty of Animal and Agricultural Sciences, Diponegoro University Tembalang Campus, Semarang, 50275 - Indonesia

²Faculty of Animal Science, Gadjah Mada University Jl. Fauna 3 Bulaksumur 55281 Yogyakarta - Indonesia

Email: kismiati59@gmail.com

Abstract - The objective of this research was to evaluate the effect of H_3PO_4 concentrations and particle size of eggshell used in the feed of laying hens on bone and blood profiles. Ninety-six laying hens (Isa Brown strain) age 25 weeks were reared in individual battery cage and divided into 8 groups randomly. Group 1 was fed using eggshell without H_3PO_4 and particle size of ≤ 1 mm (feed 1), group 2 was fed using eggshell that has been soaked in H_3PO_4 3% and particle size of ≤ 1 mm (feed 2), group 3 were fed using eggshell that has been soaked in H_3PO_4 4% and particle size of ≤ 1 mm (feed 3), group 4 was fed using eggshell that has been soaked in H_3PO_4 5% and particle size of ≤ 1 mm (feed 4), group 5 was fed using eggshell that without H_3PO_4 and particle size of ≤ 3 mm (feed 5), group of 6 was fed using eggshell that has been soaked in H_3PO_4 3% and particle size of ≤ 3 mm (feed 6), group 7 was fed using eggshell that has been soaked in H_3PO_4 4% and particle size of ≤ 3 mm (feed 7) and a group of 8 was fed using eggshell that has been soaked in H_3PO_4 5% and particle size of ≤ 3 mm (feed 8). A Completely Randomized Design patterns factorial $4 \times 2 \times 3$ was used in this research. Result of this research showed that had interaction effect ($P > 0.05$) between the H_3PO_4 concentration and particle size of eggshell on weight, diameter of tibia bone and calcium and phosphorus content of the blood. The concentration of H_3PO_4 or particles size also had no effect ($P > 0.05$) on all variables.

Keywords — Egg shell, H_3PO_4 , Particle Size, Bone and Blood

Submission: April 25, 2013

Corrected : June 3, 2013

Accepted: June 10, 2013

Doi: [10.12777/ijse.5.1.12-14](https://doi.org/10.12777/ijse.5.1.12-14)

[How to cite this article: Kismiati, S., Yuwanta, T., Zuprizal, and Supadmo. (2013). Effect of H_3PO_4 Concentration and Particle Size of the Eggshell Used in Laying Hens Fed on Bone and Blood. *International Journal of Science and Engineering*, 5(1),6-11. Doi: [10.12777/ijse.5.1.12-14](https://doi.org/10.12777/ijse.5.1.12-14)]

I. INTRODUCTION

Development of the poultry industry have produced waste that can cause environmental pollution. Eggshell is one of the poultry industry waste but contain high calcium and little phosphorus. Besides that eggshell also contains micro mineral, protein, hormone and essential amino acid. Chicken eggshell powder contains Ca, P, Sr, Pb, Al, Cd, Hg, V, B, Fe, Zn, P, Mg, N, F, Se, Cu, and Cr (Schaafsma, et al. 2000). King'ori (2011) stated that eggshells can be utilized for various purposes that minimize their effect on environmental pollution. Eggshell powder has been reported to increase bone mineral density in people and animals with osteoporosis. According to Gongruttananun (2011) eggshell contains 34.89% calcium, 0.001% phosphorus and 5.35% protein. The use of coarse eggshell as substitute of fine limestone in the diet did not affect on plasma calcium and phosphorus content of the blood and tibia bone quality of RIR strain laying hens aged 38 weeks. Rovensky et al. (2003) stated that clinical and

experimental studies indicate that eggshell powder has a positive effect on bone and cartilage that can be used for the prevention and treatment of osteoporosis.

The use of eggshell should be sterilized because it contains bacteria. The research of Kismiati et al. (2012) proved that soaking eggshell in the H_3PO_4 decrease the amount of bacteria and increase the phosphorus content. Meddleton and Ferket (2001) reported that the use of H_3PO_4 can kill bacteria and increase the phosphorus content of broiler carcass meal used as animal feed. Animal feed industry also uses H_3PO_4 to make monocalcium phosphate and dicalcium phosphate.

Phosphorus is the third most expensive nutrient after protein and energy. Phosphorus can be obtained from vegetable or animal feedstuffs. Phosphorus from animal and its products have higher availabilities than phosphorus from plant. Availability of phosphorus derived from plants approximately 30-40% (NRC, 1994) or 50% (North, 1978) and 30% (Larbier and Leclercq, 1994). This

is caused phosphorus of plants bound with phytic acid that is not easily digested by birds, because birds do not have the phytase enzyme.

The research objective was to evaluate the effect of H₃PO₄ concentration and particle size of eggshell used in the feed on the bone and blood profile of laying hens. Utilization of eggshell in poultry feed expected contribute to minimize environmental pollution.

II. MATERIALS AND METHODS

The study begins with collecting the eggshell, soaking in hot water (80° C) for 15 minutes and then divided into 4 part. Part 1 without H3PO4, part 2 was soaked in H3PO4 3%, part 3 was soaked in H3PO4 4% and part 4 was soaked in H3PO4 5% during 15 minutes. Each piece was milled with a filter size of 1 and 3 mm in order to obtain a particle size of <1 mm and <3 mm and then used as a mineral source of hens.

A total of 96 laying hens (Isa Brown strains) 25 weeks of age were used in this research and reared in individual battery cages for 12 weeks. Laying hens were divided into 8 groups randomly, so that each group consisted of 12 laying hens (4 hens/replication). Group 1 was fed using eggshell without H3PO4 and particle size of < 1 mm (Feed 1), group 2 was fed using eggshell that has been soaked in H3PO4 3% and particles size of < 1 mm (feed 2), group 3 was fed using eggshell that has been soaked in H3PO4 4% and particles size of < 1 mm (feed 3), group 4 was fed using eggshell that has been soaked in H3PO4 5% and particles size of < 1 mm (feed 4), group 5 was fed using eggshell without H3PO4 and particles size of < 3 mm (feed 5), group 6 was fed using eggshell that has been soaked in H3PO4 3% and particles size of < 3 mm (feed 6), group 7 was fed using eggshell that has been soaked in H3PO4 4% and particles size of < 3 mm (feed 7), group 8 was fed using eggshell that has been soaked in H3PO4 5% and particles size of < 3 mm (feed 8).

The composition of feed ingredients and nutrient content of feed research can be seen in Table 1. Proximate analysis of feed ingredients was used Wende methods (Tillman et al., 1986). Analysis of calcium and phosphorus contents of eggshell were determined by AAS method according to AOAC (1995) were cited in Hall et al. (2003).

Variables measured was the tibia bone profile (weight, volume, diameter, calcium and phosphorus content) and a blood profile (calcium and phosphorus content). Tibia profile measurements performed at the end of the study, while analysis of calcium and phosphorus in the blood done every 4 weeks. Completely randomized factorial design 4 x 2 x 3 was used in this research. Analysis of data used an analysis of variance and processed by Statistical Analysis Program (SAS).

III. RESULT AND DISCUSSION

Tibia Bone Profile

The effect of H₃PO₄ concentrations and particle size of eggshell on tibia profile (weight, volume, and diameter) were presented in Table 2. The results of various analyzes indicate that there is no interaction effect between H₃ PO₄

concentrations and particle size of eggshell on the tibia weight, volume, diameter, calcium and phosphorus content of the tibia bone.

Table 1. The Ingredients and Calculated Composition of the Experimental Feed

Feedstuff	Treatment feed							
	Feed 1	Feed 2	Feed 3	Feed 4	Feed 5	Feed 6	Feed 7	Feed 8
Corn	70.00	70.00	70.00	70.00	70.00	70.00	70.00	70.00
Soybean extract	10.00	10.00	10.00	10.00	10.00	10.00	10.00	10.00
Poultry Meat Meal	11.00	11.00	11.00	11.00	11.00	11.00	11.00	11.00
Eggshell waste	5.00*	5.00*	5.00*	5.00*	5.00*	5.00*	5.00*	5.00*
Dicalcium Phosphate	1.10	1.10	0.80	0.75	1.10	1.00	0.80	0.75
Ca CO ₃	2.40	2.50	2.70	2.75	2.40	2.50	2.70	2.75
Tyramin ^a	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Na Cl	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
ME (kcal/kg)	2892.5	2892.5	2892.5	2892.5	2892.5	2892.5	2892.5	2892.5
Crude Protein (%)	16.45	16.45	16.45	16.45	16.45	16.45	16.45	16.45
Extract Ether (%)	3.30	3.30	3.30	3.30	3.30	3.30	3.30	3.30
Crude fiber (%)	2.70	2.70	2.70	2.70	2.70	2.70	2.70	2.70
Ca (%)	3.50	3.50	3.50	3.50	3.50	3.50	3.50	3.50
P available (%)	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50
Lysine (%)	1.10	1.10	1.10	1.10	1.10	1.10	1.10	1.10
Methionine (%)	0.44	0.44	0.44	0.44	0.44	0.44	0.44	0.44

*eggshell was sterilized with water of 80° C with a particle size of <1 mm, ^aeggshell was sterilized with water of 80° C and then soaked in 3% H₃PO₄ with particle size of <1 mm, ^beggshell was sterilized with water of 80° C and then soaked in 4% H₃PO₄ with particle size <1 mm, ^ceggshell was sterilized with water of 80° C and then soaked in 5% H₃PO₄ and particle size of <1 mm, ^deggshell was sterilized with water of 80° C and particle size <3 mm, ^eeggshell was sterilized with water of 80° C and then soaked in 3% H₃PO₄ and particle size <3 mm, ^feggshell was sterilized with water of 80° C and then soaked in 4% H₃PO₄ and particle size <3 mm, ^geggshell was sterilized with water of 80° C and then soaked in 5% H₃PO₄ and particle size <3 mm.
^aMetionin, lisin, vitamin A, D₃, E, K, B₁, B₂, B₆, B₁₂, C, Ca-pantothenat, Niacin,Cholin Chloride Mn, Fe,I,Z.Co,Cu,Santoquin dan Zinc Bacitracin.

Profile of tibia bone (weight, volume, diameter, calcium and phosphorus content) had no significant different on different concentrations of H₃PO₄ (3-5%) and control. Onyango et al. (2003) and Fleming (2008) stated that bone mineral content correlated with mineral intake. Calcium and phosphorus are two most abundant mineral elements in bone (McDonald et al., 2002). Fard et al. (2010) stated that the use of calcium for bone formation. Bone calcium is mainly composed of calcium phosphate and some calcium carbonate. Feed of this studies contain a same of calcium and phosphorus despite concentrations of H₃ PO₄ was different so weight, volume, diameter, calcium and phosphorus content of the tibia were not significantly different.

Soaking eggshell in H3PO4 decreases the amount of bacteria and increase the phosphorus content and concentration of H₃PO₄ (3 – 5%) had no effected on feed intake, calcium intake and phosphorus intake (Kismiati et al., 2012). Table 2 showed that the higher concentration of H₃PO₄ caused DCP in feed decreases. Results of this research indicated that the phosphorus of H₃PO₄ can replace phosphorus from DCP.

Table 2. Tibia bone profile of laying hens fed using eggshell soaked in H₃ PO₄ and different particle sizes.

Variable	Particle size (mm)	Concentration of H ₃ PO ₄ (%)				Means ^{ns}
		0	3	4	5	
Weight (g)	≤1	9.47	9.33	9.57	9.34	9.43
	≤3	9.65	9.66	9.45	9.92	9.67
Means ^{ns}		9.56	9.49	9.51	9.63	
Volume (cc)	≤1	6.70	6.73	7.10	6.93	6.86
	≤3	7.16	7.17	7.20	7.13	7.14
Means ^{ns}		6.93	6.95	7.10	7.03	
Diameter (cm)	≤1	0.58	0.60	0.59	0.56	0.58
	≤3	0.59	0.58	0.60	0.58	0.59

^{ns} Non significant (P>0.05)

Particle size of eggshell had no significant effect (P>0.5) on the bone profile (bone weight, bone volume and bone diameter). Profile tibia bone did not differ due to

differences in particle size is very small. Pizzolante *et al.* (2009) stated that the use of limestone particle size of 2-5 mm and 11-5mm increasing calcium tibia bone. Saunders-Blades *et al.* (2009) examined the effect of particle size of oyster shell, white limestone, red limestone and gray limestone on bone quality and result that a higher weight on the use of a mixture of 1/3 of small particles + 2/3 larger than the size of small particles, but the calcium content was not significantly different. The results of this research correspond with the study of de Witt *et al.* (2009) that the particle size did not effect on weight and diameter of tibia bone of laying hens.

The average weight of tibia bone 7.41 to 9.92 g, the volume of 6.93 to 7.10 cc, while the particle size based, the weight of tibia 7.41 to 7, 47 g and volume 6.86 to 7.14 cc. Tibia weight lower than the study of Rath *et al.*, (2000), while the volume in the normal range. Volume of tibia bone is 7 to 7.17 cc (Zang and Coon, 1997) and tibia weight is 18.56 g (Rath *et al.*, 2000).

Calcium and Phosphorus of Blood

Calcium and phosphorus content of the blood can be seen in Table 3. Blood calcium and phosphorus did not influenced by the interaction of H₃PO₄ concentration and particle size. Concentration of H₃PO₄ or particle size did not significant effect on blood calcium and phosphorus.

Table 3. Blood calcium and phosphorus of laying hens fed using eggshell soaked in H₃PO₄ and different particle sizes.

Variable	Particle size (mm)	Concentration of H ₃ PO ₄ (%)				Means ^{ms}
		0	3	4	5	
		mg/dl				
Calcium	1	24,38	21,91	25,18	24,03	23,87
	3	24,76	27,21	24,75	24,59	25,32
Means ^{ms}		24,57	24,56	24,96	24,31	
Phosphorus	1	7,36	7,41	7,55	7,30	7,41
	3	7,39	7,56	7,35	7,56	7,46
Means ^{ms}		7,37	7,48	7,45	7,43	

^{ms} Non significant (P>0.05)

Calcium and phosphorus of the blood were not significantly different at the different H₃PO₄ concentrations. Research using the feed with the same content of calcium and phosphorus although concentrations of H₃PO₄ differently so the blood calcium and phosphorus were not significantly different. According to Celebi *et al.* (2005) blood phosphorus correlated with phosphorus of the feed. Pelicia *et al.* (2009) stated that calcium blood concentration is affected by calcium level of the feed. The study of Fard *et al.* (2010) showed that the decrease and increase in calcium and phosphorus feed 10% of 0.37% does not increase or decrease blood calcium and phosphorus of the blood.

Eggshell particle size used in the feed had no effect on calcium and phosphorus of blood caused by differences particle size can be tolerated in the mechanism of absorption of calcium and phosphorus. According to McNab and Boorman (2002) Calcium and phosphorus from feed digested in the proventriculus and ventrikulus then absorbed by the blood in the intestinal/intestines. Absorption was determined to be most efficient from the duodenum to the upper jejunum. Elble *et al.* (2009) stated that the absorption of calcium from the small particle size

of CaCO₃ is lower than the large particle size. According Pizzolante *et al.* (2006), the smaller particle size the more easily absorbed then larger. Pelicia *et al.* (2009) reported that the particle size of limestone 0.18 and 3.13 mm did not significantly affect on blood calcium concentrations of laying hens aged 58 weeks. Pelicia *et al.* (2011) examined the use of limestone particle size of 0.18 and 3.90 mm at strain Isa Brown hens aged 23 weeks. The results showed that the particle size of limestone 0.18 and 3.90 mm had no effect on blood calcium although the solubility *in vitro* particle size 0.18 mm higher than 3.90 mm. Azar *et al.* (2011) investigated the use of rock/perlite particle size of 1.5 and 3 mm as a mineral resource of broilers aged 42 days and obtained the result that the concentration of blood serum calcium and phosphorus did not differ.

IV. CONCLUSIONS

The conclusion of this study is H₃PO₄ concentrations used for soaking the eggshell did not effect on the tibia bone profile and blood profile, as well as particle size.

ACKNOWLEDGMENT

The authors wish to thank the Directorate of Higher Education, Ministry of Education and Culture of Indonesia, Dean of Faculty of Animal Science and Rector of the Gadjah Mada University.

REFERENCES

- [1]. Azar, H.R.E., Adl, K.N., Nezhad, Y.E. and Moghaddam, M., 2011. Effect of different levels and particle sizes of perlite on serum biochemical factors of broiler chicks. *Afr. J. Biotechnol.* 10: 3232-3236.
- [2]. Celebi, S., Bolukbasi, C. dan Utlu, N., 2005. The influence of dietary phosphorus level on plasma calcium and phosphorus, eggshell calcium and phosphorus. *Int. J. Poult. Sci.* 4(7): 497- 499.
- [3]. De Witt, F.H., Kuleile, N.P. van der Merwe, H.J. and Fair, M.D., 2009. Effect of limestone particle size on bone quality characteristics of hens at end-of-lay. *South Africa J. of Anim. Sci. Supplement 1*: 41-44.
- [4]. Elble A. E, Hill, K.M, Park, C.Y, Martin, B.R, Peacock, M. and Weaver, C.M. 2009. Effect of calcium carbonate particle size on calcium absorption and retention in adolescent girls. *J. Am. Coll. Nutr.* 30(3): 171-177.
- [5]. Fard, M.K, Moghadam, H. N. and Saki, A. A. 2010. Effect of different levels of calcium, phosphorus and vitamin D₃ on the calcium, phosphorus and magnesium of plasma, hatchability and performance on the broiler breeder hens. *Research Journal of Biological Sciences* 5(2): 223-227.
- [6]. Fleming, R.H. 2008. Nutritional Factor Affecting Poultry Bone Health. In *Proceedings of The Nutrition. Society*, 67: 177 - 183. Symposium on Diet and Bone Health. doi: 10.1017/S0029665108007015.
- [7]. Gongruttananun, N. 2011. Effects of eggshell calcium on productive performance, plasma calcium, bone mineralization, and gonadal characteristics in laying hens. *Poult. Sci.* 90: 524-529.
- [8]. Hall, R.E., Shirly, R.B, Bakallii, R.I, Aggrey, S.E., Pesti, G.M. and Edwards Jr H.M.2003. Power of two methods for the estimation of bone ash of broiler. *Poult. Sci.* 82: 414-418.
- [9]. Herzig, I., Strakova, E. and Sucy, P. 2008. Long-term Application of Clinoptilolite Via the Feed of Layer and Its Impact on the Chemical Composition a Long Bones of Pelvic Limb (*femur* and *tibiotarsus*) and Eggshell. *Veterinarni Medicina* 53 (10): 550-554.
- [10]. King'ori, A.M. 2011. A review of the uses of poultry eggshells and shell Membranes. *Int. J. Poult. Sci.* 10 (11): 908-912.
- [11]. Kismiati, S., Yuwanta, T., Zuprizal dan Supadmo. 2012. The Physical and Chemical Characteristics of Eggshell Waste as Phosphorus Fortification: Its Effect on Egg Production and Eggshell Quality of Laying Hens. *Proceedings of the The 1st Poult. Int. Seminar*: 136 - 143.
- [12]. Larbier, M. and H. Leclercq. 1994. *Nutrition and Feeding of Poultry*. Translated and Edited by J. Wiseman. University Press. Nottingham.

- [13]. McDonald, P., Edwards, R.A., Greenhalgh, J.F.D. and Morgan, C.A. 2002. *Animal Nutrition*. Six edition. Ashford Colour Press, Ltd, Gosport.
- [14]. McNab and Boorman, K.N. 2002. *Poultry Feedstuffs Supply, Composition and Nutritive Value*. Poultry Science Symposium Series 28. CABI Publishing.
- [15]. Middleton, T. F. and Ferket, P. R. 2001. Effect of level of acidification by phosphoric acid, storage temperature, and length of storage on the chemical and biological stability of ground poultry mortality carcasses I. *Poult. Sci.* 80: 1144-1153.
- [16]. N.R.C. 1994. *Nutrient Requirement of Poultry*. Ninth revised edition. National Academy Press. Washington, D.C.
- [17]. North, M.O. 1984. *Commercial Poultry Production*. Second Edition. Avi Publishing Company, Inc. West Port, Connecticut.
- [18]. Onyango, E.M., Hester, P.Y., Strohshine, R. and Adeola, O. 2003. Bone densitometry as an indicator of percentage tibia ash in broiler chicks fed varying dietary calcium and phosphorus levels¹. *Poult. Sci.* 82: 1787- 1791.
- [19]. Pelicia, K., Garcia, E.A., Faitarone, A.B.G., Silva, A.P., Berto, D.A., Molino, A.B. and Vercese, F. 2009. Calcium and available phosphorus levels for laying hens in second production cycle. *Brazilian J. Poult. Sci.* 11(1): 39-49.
- [20]. Pelicia, K., Maurao, J.L.M., Gracia, E.A., Viheiro, V.M.C., Berto, D.A., Molino, A.B., Faitarone, A.B.G., Vercese, F., Santos, G.C. and Silvia, A.P. 2011. Effect of dietary calcium levels and limestone particle size on the performance, tibia and blood of laying hens. *Brazilian J. Poult. Sci.* 1(13): 29-34.
- [21]. Pizzolante C.C., Gracia, E.A., Legane, C., Saldanha, E.S.P.B, Deotato, A.P., Faitarone, A.B.G., Scherer, M.R. and Batista, L. 2006. Effects of calcium levels and limestone particle size on the performance of semi-heavy layers in their second production cycle. *Brazilian J. Poult. Sci.* 8 (3): 173 - 176.
- [22]. Rath, N.C., Huff, G. R. and Balog, J.M. 2000. Factor regulating maturity and strength in poultry. *Poult. Sci.* 79: 1024 - 1032.
- [23]. Rovenský J, Stancíková, M, Masaryk, P, Svík, K. and Istok, R. 2003. Eggshell calcium in the prevention and treatment of osteoporosis. *Int. J. Clin. Pharmacol. Res.* 23 (2-3): 83-92.
- [24]. Saunders-Blades, J. L., MacIsaac, J. L., Korver, D. R. and Anderson, D. M. 2009. The effect of calcium source and particle size on the production performance and bone quality of laying hens. *Poult. Sci.* 88 :338-353.
- [25]. Schaafsma, A., Pakan, I., Hofstede, G.J.H. Muskiet, F.A.J., Van Der Veer, E. and De Vries, P.J.F. 2000. Mineral, amino acid, and hormonal composition of chicken eggshell powder and the evaluation of its use in human nutrition. *Poult. Sci.* 79: 1833-1838.

Effect of H3PO4 Concentration and Particle Size of the Eggshell Used in Laying Hens Fed on Bone and Blood

ORIGINALITY REPORT

9%

SIMILARITY INDEX

5%

INTERNET SOURCES

7%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	Orr, J.B.. "STUDIES ON NUTRITION IN RELATION TO IMMUNITY.", The Lancet, 19310530 Publication	1%
2	czytanki.net Internet Source	1%
3	Submitted to University of Fort Hare Student Paper	1%
4	Submitted to Pembroke Student Paper	1%
5	www.kutuu.co.uk Internet Source	<1%
6	www.arch-anim-breed.net Internet Source	<1%
7	www.ncbi.nlm.nih.gov Internet Source	<1%
8	Submitted to Nottingham Trent University Student Paper	<1%

9	www.dncustoms.gov.vn Internet Source	<1%
10	acervodigital.ufpr.br Internet Source	<1%
11	Submitted to Wageningen University Student Paper	<1%
12	Submitted to University of Edinburgh Student Paper	<1%
13	stud.epsilon.slu.se Internet Source	<1%
14	academic.oup.com Internet Source	<1%
15	fvhe.vfu.cz Internet Source	<1%
16	www.myanmarveterinaryassociation.org Internet Source	<1%
17	istocar.bg.ac.rs Internet Source	<1%
18	Ahmed, Nisrin, khadiga Atti, Khalid Elamin, Kamal Dafalla, Huwaida Malik, and M Dousa. "Effect of Dietary Calcium Sources on Laying Hens Performance and Egg Quality.", Journal of Animal Production Advances, 2013. Publication	<1%

19

Submitted to Universiti Putra Malaysia

Student Paper

<1%

20

K Pelicia, E Garcia, C Móri, ABG Faitarone, AP Silva, AB Molino, F Vercese, DA Berto.

"Calcium levels and limestone particle size in the diet of commercial layers at the end of the first production cycle", Revista Brasileira de Ciência Avícola, 2009

Publication

<1%

21

José Anchieta de Araujo, José Humberto Vilar da Silva, Fernando Guilherme Perazzo Costa, Janaína Maria Batista de Sousa et al. "Effect of the levels of calcium and particle size of limestone on laying hens", Revista Brasileira de Zootecnia, 2011

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Effect of H₃PO₄ Concentration and Particle Size of the Eggshell Used in Laying Hens Fed on Bone and Blood

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
