

The Microstructure of Egg Shell Waste Treated with H₃PO₄, In vitro Solubility in Different Particle Size and the Using Effect on the Egg Shell Quality of Laying Hens

by Sri Kismiyati

Submission date: 27-Apr-2020 01:03PM (UTC+0700)

Submission ID: 1308853192

File name: The_Microstructure_of_Egg_Shell_Waste_Treated_with_H3PO4.pdf (76.32K)

Word count: 3314

Character count: 15691

The Microstructure of Egg Shell Waste Treated with H₃PO₄, *In vitro* Solubility in Different Particle Size and the Using Effect on the Egg Shell Quality of Laying Hens

Sri Kismiati¹, Tri Yuwanta², Zupriza² and Supadmo²

¹Faculty of Animal Agriculture, Diponegoro University, Semarang, Central Java, Indonesia

²Faculty of Animal Science, Gadjah Mada University, Yogyakarta, Indonesia

Abstract: The aim of the research was to study the microstructure changes of egg shell waste treated with H₃PO₄ and its effect on *in vitro* solubility and egg shell quality of laying hens. Egg shell waste was soaked in water with the temperature of 80°C, drained and divided into 4 parts. Part 1 was sun dried (control), part 2 was soaked in H₃PO₄ 3%, part 3 was soaked in H₃PO₄ 4% and part 4 was soaked in H₃PO₄ 5%. Furthermore, egg shells were drained, sun dried and then grinded to a particle size of 1 and 3 mm and used in the feed of laying hens. A total of 96 twenty-five weeks old of laying hens (ISA Brown strain) were divided into 8 experimental dietary treatment in 4 x 2 factorial arrangements with 3 replication. Results of the research showed that the concentration of H₃PO₄ effected on egg shell microstructure changes and interaction of H₃PO₄ 3 and 4% and particle size 1 mm was increased *in vitro* solubility. Concentration of H₃PO₄ and particle size had not significantly effect (P>0.05) on calcium intake, egg shell weight and egg shell percentage but significantly effect (P<0.05) on phosphorus intake and egg shell thickness. Egg shell thickness was increased at H₃PO₄ 5% and particle size 3 mm. The research concluded that H₃PO₄ resulted in microstructure changes and *in vitro* solubility but did not change the quality of the egg shell, except egg shell thickness.

Key words: Egg shell waste, H₃PO₄ concentrations, particle size, microstructure, *in vitro* solubility and egg shell quality

INTRODUCTION

Egg shells are rich in minerals and contain protein essential and non-essential amino acids. The minerals contained in the egg shell are Ca, P, Fe, Mg, I, Zn, K and Na. The content of calcium (Ca): 35-37%, while phosphorus (P): 0.03 to 0.14% and Mg: 1.07 to 1.10%. Egg shell phosphorus is an inorganic phosphorus so that it has high availability. Excess calcium resulting in poor soil, while phosphorus cause damage to aquatic ecosystems/eutrophication. Egg shell is contaminated bacteria originating from *cloaca*. Davis *et al.* (2008) found the bacterium of *Salmonella enteritidis* and *Salmonella heidelberg*, then used water (80°C) to kill the bacteria. According to Guinotte *et al.* (1991), H₃PO₄ can be used as an anti-bacterial sea shells and loosen calcium bonds so that it has high solubility.

Ca and P are highly needed by the poultry for egg shell formation and egg shell quality is important for eggs producers. Improving of the egg shell quality has a positive impact on economic efficiency. According to the NRC (1994), the need for calcium and phosphorus for laying hens in laying period is 3.5% (Ca): 0.45% (P). The need for phosphorus is less than calcium but it is the most expensive mineral. The lack of calcium and phosphorus decreases egg shell quality. The

decreasing quality of egg shell would adverse manufacturers because the egg will break easily.

The effectiveness of mineral resources is associated with particle size. The research results of Pizzolante *et al.* (2009), showed that limestone with a large particle size has lower *in vitro* solubility than the smaller particle size but in reverse, *in vivo* solubility showed that the bigger particles have better absorption.

The objective of this study was to investigate the microstructure changes of egg shell waste treated with H₃PO₄, the *in vitro* solubility on the particle size of 1 and 3 mm and the effect in the feed to egg shell quality of laying hens. The results are expected to be used in reference to the use of egg shell waste in poultry feed.

MATERIALS AND METHODS

The research used egg shell of laying hens obtained from the bakery industries and the *lunpia* industries (the typical food of Semarang). The egg shell waste was soaked in water with the temperature of 80°C for 15 minutes, drained and divided into 4 parts. Part 1, without the treatment of H₃PO₄ (control), part 2 soaked in H₃PO₄ 3%, part 3 soaked in H₃PO₄ 4%, part 4 soaked in H₃PO₄ 5%. The soaking was conducted for 15 minutes and dried under the sun for 3 days. The egg shell waste was

Table 1: The ingredients and calculated composition of diet treatment

Ingredients (%)	Dietary treatment (Feed)			
	1 and 5	2 and 6	3 and 7	4 and 8
Egg shell waste	5.00 ^a	5.00 ^b	5.00 ^c	5.00 ^d
Corn	70.00	70.00	70.00	70.00
Soybean extract	10.00	10.00	10.00	10.00
Poultry Meat Meal	11.00	11.00	11.00	11.00
Topmix*	0.25	0.25	0.25	0.25
Dicalcium Phosphate	1.10	1.00	0.80	0.75
Ca CO ₃	2.40	2.50	2.70	2.75
Salt	0.25	0.25	0.25	0.25
Calculated composition	100.00	100.00	100.00	100.00
ME (kcal/kg)	2892.50	2892.50	2892.50	2892.50
Crude Protein (%)	16.50	16.50	16.50	16.50
Ca (%)	3.50	3.50	3.50	3.50
P available (%)	0.50	0.50	0.50	0.50
Lycine (%)	1.10	1.10	1.10	1.10
Methionine (%)	0.44	0.44	0.44	0.44

^aEgg shell waste is not soaked in water 80°C (control).

^bEgg shell waste is soaked in H₃PO₄ 3%.

^cEgg shell waste is soaked in H₃PO₄ 4%.

^dEgg shell waste is soaked in H₃PO₄ 5%.

1,2,3,4 Egg shell particle size 1 mm

5,6,7,8 Egg shell particle size 3 mm

*Methionine, lycine, vitamin A, D3, E, K, B1, B2, B6, B12, C, Ca-pantothenate, Niacin, Choline Chloride Mn, Fe, I, Zn, Co, Cu, Santoquin and Zinc Bacitracin.

then evaluated on the microstructure changes by Scanning Electron Microscope (SEM) according to Zamani *et al.* (2005). After that egg shell waste was grinded into 2 particle sizes 1 and 3 mm (the sieve size 1 and 3 mm) and then *in vitro* solubility test was performed using the method of Keshavarz and Scott (1993). Furthermore the egg shells were used in the feed of laying hens. A total of 96 twenty-five weeks old of laying hens (Isa Brown strain) were divided to 8 experimental dietary treatment with 3 replication. The composition of feed ingredients and nutrient content of feed used is shown in Table 1.

The variables measured were microstructure of egg shell waste, *in vitro* solubility, calcium intake, phosphorus intake, egg shell weight, egg shell percentage and egg shell thickness. Calcium and phosphorus are the average consumption per day during the research. The egg shell weight and egg shell thickness were obtained for 3 days every 4 weeks and 3 times during the research. The experimental design was Completely Randomized Design factorial pattern. The data were analyzed by analysis of variance. Duncan's Multiple Range Test was used in means comparison when found effect on treatment.

RESULTS AND DISCUSSION

The effect of H₃PO₄ concentration on microstructure of egg shells waste: The microstructure changes of egg shell waste which was soaked in water 80°C (control), soaked in H₃PO₄ 3%, soaked in H₃PO₄ 4% and soaked in H₃PO₄ 5% can be seen in Fig. 1. The microstructure of egg shell which was not soaked in H₃PO₄, looks

Table 2: The *in vitro* solubility of egg shell waste soaked in H₃PO₄ 3%, 4%, 5% and control on particle size 1 and 3mm

Particle size (mm)	Concentration of H ₃ PO ₄				
	Control	3%	4%	5%	Average
1	19.84 ^b	20.49 ^a	20.50 ^a	19.87 ^b	20.18
3	13.42 ^d	17.76 ^c	17.19 ^d	16.60 ^e	16.24
Average	16.56	19.13	18.84	18.24	

Different superscript at the same raw and column indicate significantly different (P<0.05).

cohesive and massive. The higher the concentration of H₃PO₄, it causes the microstructure to be more hollow/loose. Guinotte *et al.* (1991) stated that H₃PO₄ induce microcracking of calcium mineral binding. According to Rodriguez da-Navarro *et al.* (2007) who was cited by Arpasova *et al.* (2010), egg shell microstructure changes can be caused by chemical processes.

The effect of H₃PO₄ concentration and particle size on *in vitro* solubility of egg shell waste: *In vitro* solubility of egg shell waste soaked in H₃PO₄ 3%, 4%, 5% and control in the particle size of 1 and 3 mm can be seen in Table 2. The results of statistical analysis show that there is an interaction effect (P<0.05) between H₃PO₄ concentration and particle size on *in vitro* solubility of egg shell.

The *in vitro* solubility of egg shell waste increased significantly in H₃PO₄ 3-4% and the particle size of 1 mm but then decreased at a concentration of 5%. The solubility in 3 mm particle size increased along with the increase of H₃PO₄ concentrations (3-5%) but lower than the particle size 1 mm. The result agree with the observation of Guinotte *et al.* (1991) that the H₃PO₄ induce microcracking of sea shells so that it increases the solubility. Pizzolante *et al.* (2009) reported that *in vitro* solubility of limestone with the particle size 2.83 mm (10.85%) was lower than 0.185 mm (18.7%). The solubility of limestone calcium with the particle size 3.90 mm was lower than the particle size 0.18 mm. *In vitro* solubility of the particle size 3.90 mm was 22.37% and 0.18 mm was 25.56% (Pelicia *et al.*, 2011). Saunders-Blades *et al.* (2009) stated that *in vitro* solubility of large limestone particles was significantly lower than the smaller particle size.

Calcium and phosphorus intake: The effect of H₃PO₄ concentration and particle size on calcium and phosphorus intake presented in Table 3. The results showed that calcium intake is not significantly different to the use differences of H₃PO₄ concentrations and particle size but phosphorus intake is effected by interaction of H₃PO₄ concentrations and particle size. Calcium intake is not significant caused the H₃PO₄ concentrations and particle size are not change the nutrition of the feed (Table 1). However, the phosphorus intake was higher in use H₃PO₄ 3% particle size of 1




Fig. 1(a-d): Macrostructure of egg shells waste were soaked in water 80°C, H₃PO₄ 3%, 4% and 5%. A: cuticula, B: vertical, C: palisade layer, D: mamillary layer, E: inner shell membrane

Table 3: The effect of H₃PO₄ concentration and particle size of egg shell waste on calcium and phosphorus intake

Variables	Particle size (mm)	Concentration of H ₃ PO ₄				Average
		Control	3%	4%	5%	
Calcium intake (g/hen/day)	1	4.06	4.12	3.89	3.90	3.99
	3	3.90	4.01	4.11	4.05	
Average		3.98	4.06	4.00	3.97	
Phosphorus intake (g/hen/day)	1	0.58 ^a	0.59 ^a	0.57 ^{ab}	0.55 ^c	0.57
	3	0.56 ^{bc}	0.57 ^{ab}	0.59 ^a	0.59 ^a	

Different superscript at the same raw and column indicate significantly different (P<0.05).

mm, H₃PO₄ 4% particle size of 3 mm, H₃PO₄ 5% and particle size of 3 mm. Calcium and phosphorus intake was in the normal range. Calcium intake of laying hens aged 25-35 weeks is 3.98 to 5.21 g/hen/day (Pelicia *et al.*, 2011) and phosphorus intake is 4.0 to 4.1 g/hen/day (Pelicia *et al.*, 2009).

The quality of egg shell: Table 4 shows the egg shell quality. Concentrations of H₃PO₄ had no significant effect on egg shell weight and egg shell percentage but significant effect on egg shell thickness. It was due to the calcium intake and phosphorus intake (Table 3).

Squires (2003), Ahmadi and Rahimi (2011) stated that calcium have the role in the eggs formation especially egg shell. According to Ogawa *et al.* (2004) and Schaafsma *et al.* (2000) egg shell contains high calcium and little phosphorus. Phosphorus intake is important in maintaining egg shell quality (Leeson and Summers, 2005). Egg shell thickness is influenced by phosphorus intake (McDonald *et al.*, 2002).

The particle size of the egg shell waste had no significant effect on egg shell weight and egg shell percentage but significant effect on egg shell thickness.

Table 4: The effect of H₃PO₄ concentration and particle size of egg shell waste on egg shell quality

Variabel	Particle size (mm)	Concentration of H ₃ PO ₄				Average
		Control	3%	4%	5%	
Egg shell weight (g)	1	5.72	5.67	5.72	5.70	5.70
	3	5.85	5.71	5.78	5.92	5.81
Average		5.78	5.68	5.75	5.81	
Egg shell percentage	1	9.52	9.50	9.78	9.59	9.60
	3	9.76	9.62	9.54	9.74	9.66
Average		9.64	9.56	9.66	9.66	
Egg shell thickness (mm)	1	0.377	0.367	0.375	0.371	0.372 ^b
	3	0.382	0.374	0.378	0.387	0.380 ^a
Average		0.379 ^a	0.370 ^b	0.376 ^b	0.379 ^a	

Different superscript at the same raw and column indicate significantly different (P<0.05)

The results correspond with the other studies. Pizzolante *et al.* (2009) reported that limestone particle size have no effect on egg shell weight. The research results of de Witt *et al.* (2009) showed that the particle size have no effect on weight and percentage. Pelicia *et al.* (2011) also stated that limestone the particle size of fine and coarse limestone have no effect on the weight. Phirinyane *et al.* (2011) stated that the limestone particle size have no effect on the character of egg shell. The research results were not in accordance with the research of Ekmay and Coon (2010) who stated that limestone particle size increases the weight of egg shells. Lichovnikova (2007) reported that particle size have effect on egg shell thickness.

The average weight of egg shell, egg shell percentage are in the normal range. According to Nys *et al.* (2011), the egg shell weight about 5.5 g or 8.5-10.5%. Zita *et al.* (2009) stated that egg shell thickness of Isa Brown strain is 0.37 to 0.38 mm.

ACKNOWLEDGEMENT

The experiment project was funded by Directorate General of Higher Education, Ministry of National Education of Republic of Indonesia (Doctor Grant). The authors would like to thank to the Director General of Higher Education, Ministry of National Education of Republic of Indonesia who have provided funding. The authors would also like to the Dean of the Faculty of Animal Science and the Rector of the Gadjah Mada University.

REFERENCES

- Ahmadi, F. and F. Rahimi, 2011. Factors affecting quality and quantity of egg production in laying hens: A review. *World Applied Sci. J.*, 12: 372-384.
- Arpasova, H., M. Halaj and P. Halaj, 2010. Egg shell quality and calcium utilization in feed of hens in repeated laying cycles. *Czech J. Anim. Sci.*, 55: 66-74.
- Davis, A.L., P.A. Curtis, D.E. Conner, S.R. McKee and L.K. Kerth, 2008. Validation of cooking method using shell egg inoculated with salmonella serotypes enteridis and heidelberg. *Poult. Sci.*, 87: 1637-642.

- de Witt, F.H., N.P. Kuleile, H.J. van der Merwe and M.D. Fair, 2009. Effect of limestone particle size on egg production and egg shell quality of hens during late production. *South Afr. J. Anim. Sci.*, 39: 39-40.
- Ekmay, R.D. and Coon, 2010. The effect of limestone particle size on the performance of three broiler breeder poretines. *Int. J. Poult. Sci.*, 9: 1038-1042.
- Guinotte, F., Y. Nys and F. de Monredon, 1991. The effects of particle size and origin of Calcium Carbonat on performance and ossification characteristics in broiler chicks. *Poult. Sci.*, 70: 1908-1920.
- Keshavarz, K. and M.L. Scott, 1993. The effect of solubility and particle size of calcium sources on shell quality and bone mineralization. *J. Applied Poult. Res.*, 2: 259-287.
- Leeson, S. and J.D. Summers, 2005. *Commercial Poultry Nutrition 3rd Edition*. University Books Guelph, Ontario, Canada.
- Lichovnikova, M., 2007. The effect of dietary Calcium source, concentration and particle size on calcium retention, egg shell quality and overall Calcium requirement in laying hens. *Br. Poult. Sci.*, 48: 71-75.
- McDonald, P., R.A. Edwards, J.F.D. Greenhalgh and C.A. Morgan, 2002. *Animal Nutrition*. Sixth edition. Ashford Colour Press, Ltd, Gosport.
- NRC, 1994. *Nutrient Requirement of Poultry*. Ninth revised Edn., National Academy Press. Washington, D.C.
- Nys, Y., M. Bain and Immerseel, 2011. *Improving the Safety and Quality of Eggs and Eggs Products*. Vol 1. Woodhead Publishing Cambridge.
- Ogawa, H., M. Uehara, T Kuwayama, M. Kawashima and K. Tanaka, 2004. Change in Calcium, Magnesium and Phosphorus contents of egg shell during stay in oviduct uterus in the guineafowl and the chicken. *J. Poul. Sci.*, 41: 236-240.
- Pizzolante, C.C., E.S.P.B. Saldanha, C. Lagana, S.K. Kakimoto and C.K. Togashi, 2009. Effect of calcium levels and limestone particle size on egg quality of semi-heavy layers in their second production cycle. *Rev. Braz. Cienc. Avic.*, 11: 79-86.

- Pelicia, K., E.A. Garcia, A.B.G. Faitarone, A.P. Silva, D.A. Berto, A.B. Molino and F. Vercese, 2009. Calcium and available phosphorus levels for laying hens in second production cycle. *Braz. Poult. Sci.*, 11: 39-49.
- Pelicia, K., J.L.M. Mourao, E.A. Garcia, V.M.C. Pinheiro, D.A. Berto, A.B. Molino, A.B.G. Faitarone, F. Vercese, G.C. Santos and A.P., 2011. Effects of dietary calcium levels and limestone particle size on the performance, tibia and blood of laying hens. *Braz. Poult. Sci.*, 13: 29-34.
- Phirinyane, T.B., HJ. Van Der Merve, J.P. Hayes and J.C. Moreki, 2011. Effect of different of ration of coarse and fine limestone particle on production and shell quality of layers peak. *Online J. Anim. Sci.*, 1: 86-91.
- Rodriguez da-Navarro, A.B., A. Yebra, Y. Nys, C. JimenezLopez, J.M. Garcia-Ruiz, 2007. Analysis of avian eggshell microstructure using X -ray area detectors. *Eur. J. of Mineralogy*, 19: 391-398.
- Schaafsma, A., I. Pakan., G.J.H. Hpfstede, F.A.J. Muskiet, E. Van Der Veer and P.J.F. De Vries, 2000. Mineral, amino acid and hormonal composition of chicken egg shell powder and the evaluation of its used in human nutrition. *Poult. Sci.*, 79: 1833-1838.
- Saunders-Baldes, J.L., J.L. Maclsaac, D.R. Korver and D.M. Anderson, 2009. The effect of calcium source and particle size on the production performance and bone quality of laying hens. *Poult. Sci.*, 88: 338-353.
- Squires, E.J., 2003. *Applied Animal Endocrinology*. CABI Publishing, Cambridge, USA.
- Zamani, A., H.R. Rahmani and J. Pourreza, 2005. Supplementation a corn-soybean meal diet with manganese and zinc improves egg shell quality in laying hens. *Pak. J. Biolog. Sci.*, 8: 1311-1317.
- Zita, L., T. Eva and S. Ladislav, 2009. Effet of genotype, age anf their interation on egg quality in brown-egg laying hens. *Atavet, BRNO*, 78: 85-91.

The Microstructure of Egg Shell Waste Treated with H₃PO₄, In vitro Solubility in Different Particle Size and the Using Effect on the Egg Shell Quality of Laying Hens

ORIGINALITY REPORT

12%

SIMILARITY INDEX

6%

INTERNET SOURCES

8%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	agrifoodscience.com Internet Source	1%
2	www.tandfonline.com Internet Source	1%
3	jast.modares.ac.ir Internet Source	1%
4	seer.sct.embrapa.br Internet Source	1%
5	scialert.net Internet Source	1%
6	Submitted to University of New England Student Paper	1%
7	Submitted to Mansoura University Student Paper	1%
8	Submitted to Scottish Agricultural College Student Paper	1%

9

Submitted to University of Queensland

Student Paper

1%

10

S. Ganjigohari, N. Ziaei, A. Ramzani Ghara, S. Tasharrofi. "Effects of nanocalcium carbonate on egg production performance and plasma calcium of laying hens", Journal of Animal Physiology and Animal Nutrition, 2018

Publication

1%

11

Henrieta ARPÁŠOVÁ, PETER HAŠČÍK, Martina HANOVÁ, Jozef BUJKO. "EFFECT OF DIETARY SODIUM SELENITE AND SE-ENRICHED YEAST ON EGG-SHELL QUALITATIVE PARAMETERS OF LAYING HENS EGGS", Journal of Central European Agriculture, 2010

Publication

<1%

12

M. N. Anwar, V. Ravindran, P. C. H. Morel, G. Ravindran, A. J. Cowieson. "Effect of limestone particle size and calcium to non-phytate phosphorus ratio on true ileal calcium digestibility of limestone for broiler chickens", British Poultry Science, 2016

Publication

<1%

13

Hamilton, P. B., and J. D. Garlich. "Aflatoxin as a Possible Cause of Fatty Liver Syndrome in Laying Hens", Poultry Science, 1971.

Publication

<1%

14

S. Swiatkiewicz, A. Arczewska-Wlosek, D. Jozefiak. "Bone quality, selected blood variables and mineral retention in laying hens fed with different dietary concentrations and sources of calcium", *Livestock Science*, 2015

Publication

<1%

15

Submitted to Universiti Teknologi MARA

Student Paper

<1%

16

J. H. Wolford, K. Tanaka. "Factors Influencing Egg Shell Quality—A Review", *World's Poultry Science Journal*, 2019

Publication

<1%

17

Mohamed Ketta, Eva Tumova, Darina Chodova. "Response of three laying hen genotypes to two feed calcium levels", *Czech Journal of Animal Science*, 2019

Publication

<1%

18

muir.massey.ac.nz

Internet Source

<1%

19

repository.lib.ncsu.edu

Internet Source

<1%

20

KAWAZOE, Takayuki, Katsumi YUASA, Kaoru NOGUCHI, Masayoshi YAMAZAKI, and Mikio ANDO. "Effect of Different Sources of Vitamin D on Transfer of Vitamin D to Egg Yolk.", *Nippon Shokuhin Kagaku Kogaku Kaishi*, 1996.

Publication

<1%

21 www.softantenna.com <1%
Internet Source

22 SIM, J. S., L. M. AW-YONG, and D. B. BRAGG. <1%
"Utilization of Egg Shell Waste by the Laying
Hen", Poultry Science, 1983.
Publication

23 Submitted to University of Leeds <1%
Student Paper

24 Submitted to Asian Institute of Technology <1%
Student Paper

25 Submitted to Gyeonggi Academy of Foreign <1%
Languages
Student Paper

26 Submitted to Higher Education Commission <1%
Pakistan
Student Paper

27 TUNÇ EROL, Arzu and CUFADAR, Yusuf. <1%
"Effect of Calcium Sources and Particle Size on
Performance and Eggshell Quality in Laying
Hens", TST, 2015.
Publication

28 Y. Cufadar, O. Olgun, A.Ö. Yildiz. "The effect of <1%
dietary calcium concentration and particle size
on performance, eggshell quality, bone
mechanical properties and tibia mineral contents

in moulted laying hens", British Poultry Science, 2011

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

The Microstructure of Egg Shell Waste Treated with H₃PO₄, In vitro Solubility in Different Particle Size and the Using Effect on the Egg Shell Quality of Laying Hens

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
